



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





600030944Q

6.69. m. 8.



E. BIBL. RADCL.

10.
C. 2.
6.

~~11~~ ~~7~~
~~B. 2~~ ~~2~~
~~6~~ ~~118~~ ~~672~~
C

19149 e. $\frac{114}{3}$





600030944Q

669. m. 8.



E. BIBL. RADCL.

10.
C. 2.
6.

~~11~~ ~~7 1/2~~
~~B. 2~~ ~~118~~ ~~672~~
~~6~~ C

19149 e. $\frac{114}{3}$





**F L O R A
A D E N S I S
A L S A T I C A**

**ET
CONFINIUM REGIONUM
CIS ET TRANSRHENANA**

PLANTAS

**À LACU BODAMICO USQUE AD CONFLUENTEM MO-
SELLÆ ET RHENI SPONTE NASCENTES**

**EXHIBENS
SECUNDUM SYSTEMA SEXUALE**

CUM

ICONIBUS AD NATURAM DELINEATIS

Auctore

CAROLO CHRISTIANO GMELIN

MED. DOCT.

Secret. et Potentiff. Magni Ducis Badenf. Consiliar. Aul. Botan. et Hist.
Nat. Prof. Publ. Ord. Directore Musei Natur. Collegii Med. Bad
Membr. Ord. Acad. Imp. Nat. Cur. Nat. Scrut. Halenf. et Suevic.
Physogr. Götting. Agricult. Scient. et Art. Rheni int. Botan. Ra-
tisbon. et Mineralog. Jenenf. Sodal.

Tom. III.

**CARLSRUHÆ
IN OFFICINA AUL. MÜLLERIANA.**

1 8 0 8.

1. 200.000
2. 100.000
3. 50.000
4. 25.000

5. 12.500
6. 6.250
7. 3.125
8. 1.562
9. 781
10. 390
11. 195
12. 97
13. 48
14. 24
15. 12
16. 6
17. 3
18. 1
19. 0,5
20. 0,25
21. 0,125
22. 0,0625
23. 0,03125
24. 0,015625
25. 0,0078125
26. 0,00390625
27. 0,001953125
28. 0,0009765625
29. 0,00048828125
30. 0,000244140625
31. 0,0001220703125
32. 0,00006103515625
33. 0,000030517578125
34. 0,0000152587890625
35. 0,00000762939453125
36. 0,000003814697265625
37. 0,0000019073486328125
38. 0,00000095367431640625
39. 0,000000476837158203125
40. 0,0000002384185791015625
41. 0,00000011920928955078125
42. 0,000000059604644775390625
43. 0,0000000298023223876953125
44. 0,00000001490116119384765625
45. 0,000000007450580596923828125
46. 0,0000000037252902984619140625
47. 0,00000000186264514923095703125
48. 0,000000000931322574615478515625
49. 0,0000000004656612873077392578125
50. 0,00000000023283064365386962890625
51. 0,000000000116415321826934814453125
52. 0,0000000000582076609134674072265625
53. 0,00000000002910383045673370361328125
54. 0,000000000014551915228366851806640625
55. 0,0000000000072759576141834259033203125
56. 0,00000000000363797880709171295166015625
57. 0,000000000001818989403545856475830078125
58. 0,0000000000009094947017729282379150390625
59. 0,00000000000045474735088646411895751953125
60. 0,000000000000227373675443232059478759765625
61. 0,0000000000001136868377216160297393798828125
62. 0,00000000000005684341886080801486968994140625
63. 0,000000000000028421709430404007434844970703125
64. 0,0000000000000142108547152020037174224853515625
65. 0,00000000000000710542735760100185871124267578125
66. 0,000000000000003552713678800500929355621337890625
67. 0,0000000000000017763568394002504646778106689453125
68. 0,00000000000000088817841970012523233890533447265625
69. 0,000000000000000444089209850062616169452667236328125
70. 0,0000000000000002220446049250313080847263336181640625
71. 0,00000000000000011102230246251565404236316680908203125
72. 0,000000000000000055511151231257827021181583404541015625
73. 0,0000000000000000277555756156289135105907917022705078125
74. 0,00000000000000001387778780781445675529539585113525390625
75. 0,000000000000000006938893903907228377647697925567626953125
76. 0,0000000000000000034694469519536141888238489627838134765625
77. 0,00000000000000000173472347597680709441192448139190673828125
78. 0,000000000000000000867361737988403547205596224069953369140625
79. 0,0000000000000000004336808689942017736027981120349766845703125
80. 0,00000000000000000021684043449710088680139905601748834228515625
81. 0,000000000000000000108420217248550443400699528008744171142578125
82. 0,0000000000000000000542101086242752217003497640043720855712890625
83. 0,00000000000000000002710505431213761085017488200218604278564453125
84. 0,000000000000000000013552527156068805425087441001093021392822265625
85. 0,0000000000000000000067762635780344027125437205005465106964111328125
86. 0,00000000000000000000338813178901720135627186025002275534820556640625
87. 0,000000000000000000001694065894508600678135930125011377674102783203125
88. 0,0000000000000000000008470329472543003390679650625056888370513916015625
89. 0,00000000000000000000042351647362715016953398253125284441852569580078125
90. 0,000000000000000000000211758236813575084766991265626422209262847900390625
91. 0,0000000000000000000001058791184067875423834956328132111046314239501953125
92. 0,00000000000000000000005293955920339377119174781640660555231571197509765625
93. 0,000000000000000000000026469779601696885595873908203302776157855987548828125
94. 0,0000000000000000000000132348898008484427979369541016511388289279937744140625
95. 0,00000000000000000000000661744490042422139896847705082556941463999688720703125
96. 0,000000000000000000000003308722450212110699484238525412784707319998443603515625
97. 0,0000000000000000000000016543612251060553497421192627063923536599992218017578125
98. 0,00000000000000000000000082718061255302767487105963135319617682999961090087890625
99. 0,000000000000000000000000413590306276513837435529815676598088414999805450439453125
100. 0,0000000000000000000000002067951531382569187177649078382990442074999027252197265625
101. 0,00000000000000000000000010339757656912845935888245391914952210374995136260986328125
102. 0,000000000000000000000000051698788284564229679441226959574761051874975681304931640625
103. 0,0000000000000000000000000258493941422821148397206134797873805259374878406524658203125
104. 0,00000000000000000000000001292469707114105741986030673989369026296874392032623291015625
105. 0,000000000000000000000000006462348535570528709930153369946845131484371960163116455078125
106. 0,0000000000000000000000000032311742677852643549650766849734225657421859800815582275390625
107. 0,00000000000000000000000000161558713389263217748253834248671128287109299004077911376953125
108. 0,000000000000000000000000000807793566946316088741269171243355614140546495020389556884765625
109. 0,000000000000000000000000000403896783473158044370634585621677782070273247510194778442378125
110. 0,000000000000000000000000000201948391736579022185317292810838891035136623755097389221190625
111. 0,0000000000000000000000000001009741958682895110926586464054194455175683118775486946105953125
112. 0,00000000000000000000000000005048709793414475554632932320270972275878415593877434730529765625
113. 0,000000000000000000000000000025243548967072377773164661601354861379392077969387173652648828125
114. 0,0000000000000000000000000000126217744835361888865823308006774306896960389846935868263244140625
115. 0,00000000000000000000000000000631088724176809444329116540033871534484801949234679341316220703125
116. 0,000000000000000000000000000003155443620884047221645582700169357672424009746173396706581103515625
117. 0,0000000000000000000000000000015777218104420236108227913500846788362120048730866983532905517578125
118. 0,00000000000000000000000000000078886090522101180541139567504233941810600243654334967664527587890625
119. 0,000000000000000000000000000000394430452610505902705697837521169709053001218271674838322637939453125
120. 0,0000000000000000000000000000001972152263052529513528489187605848545265006091358374191613189697265625
121. 0,0000000000000000000000000000000986076131526264756764244593802924272632503045679187095806594848828125
122. 0,00000000000000000000000000000004930380657631323783821222969014621363162515228395935479032974244140625
123. 0,000000000000000000000000000000024651903288156618919106114845073106815812576141979677395164871220703125
124. 0,0000000000000000000000000000000123259516440783094595530574225365534079062857209898386975824356103515625
125. 0,00000000000000000000000000000000616297582203915472977652871126827670395314286049491934879121780517578125
126. 0,000000000000000000000000000000003081487911019577364888264355634138351976571430247459674395608902587890625
127. 0,0000000000000000000000000000000015407439555097886824441321778170691759882857151237298371978044512939453125
128. 0,00000000000000000000000000000000077037197775489434122206608890853458799414285756186491659890222564697265625
129. 0,000000000000000000000000000000000385185988877447170611033044454267293997071428780932458299451112823486328125
130. 0,0000000000000000000000000000000001925929944387235853055165222271336469985357143904662291497255564117431640625
131. 0,00000000000000000000000000000000009629649721936179265275826111356682349926785719523311457486277820587158203125
132. 0,000000000000000000000000000000000048148248609680896326379130556783411749633928597616557287431389102935791015625
133. 0,0000000000000000000000000000000000240741243048404481631895652783917058748169642988082786437156945514678955078125
134. 0,00000000000000000000000000000000001203706215242022408159478263919585293740848214940413932185784727573394775390625
135. 0,000000000000000000000000000000000006018531076210112040797391319597926468704241074702069660928923637866973876953125
136. 0,0000000000000000000000000000000000030092655381050560203986956597989632343521205373510348304644618189334869379765625
137. 0,00000000000000000000000000000000000150463276905252801019934782989948161717606026867551741523223090946674346898828125
138. 0,000000000000000000000000000000000000752316384526264005099673914949740808588030134337758707616115454733371724494140625
139. 0,0000000000000000000000000000000000003761581922631320025498369574724854042940150671668793538080577273666858622470703125
140. 0,000000000000000000000000000000000000188079096131566001274918478736242702147007533583439676904028863683342931123515625
141. 0,0000000000000000000000000000000000000940395480657830006374592393681213510735037667917198384520144318416714655617578125
142. 0,00000000000000000000000000000000000004701977403289150031872961968406067553675188339585991922600721592083573278087890625
143. 0,000000000000000000000000000000000000023509887016445750159364809842030337768375941697929959613003607960417866390439453125
144. 0,0000000000000000000000000000000000000117549435082228750796824049210151688841879708489649798065018039802089331952197265625
145. 0,000000000000000000000000000000000000005877471754111437539841202460507584442093985424482489903250901990104466597609878125
146. 0,0000000000000000000000000000000000000029387358770557187699206012302537922210469927122412449516254509950522332988049378125
147. 0,00000000000000000000000000000000000000146936793852785938496030061512689611052349635612062247581272549752611664940246890625
148. 0,000000000000000000000000000000000000000734683969263929692480150307563448055261748178060311237906362748763058324701234453125
149. 0,0000000000000000000000000000000000000003673419846319648462400751537817240276308740890301556189531813743815291623506172265625
150. 0,00000000000000000000000000000000000000018367099231598242312003757689086201381543704451507780947659068719076458117530861328125
151. 0,0091835496157991211560018788445431006907718522257538904738295343595382290587654306640625
152. 0,00459177480789956057800093942227155034538592611287694523691476717976911452938271533203125
153. 0,002295887403949780289000469711135775172692963056438472618457383589884557264691357666015625
154. 0,0011479437019748901445002348555678875863464815282192363092286917949422786323456788330078125
155. 0,00057397185098744507225011742778394379317324076410961815461434589747113931617283941650390625
156. 0,000286985925493722536125058713891971896586620382054809077307172948735569658086419708251953125
157. 0,0001434929627468612680625293569459859482933101910274045386535864743677848290432098535409765625
158. 0,007174648137343063403126467847299297414665509551370226932679323718389241452160492677048828125
159. 0,0035873240686715317015632339236496487073327547756851134663396618591946207260802463385244140625
160. 0,00179366203433576585078161696182482435366637738784255673316983092959731036304012316692721953125
161. 0,000896831017167882925390808480912412176833188693921278366584915464798655181520061583461109765625
162. 0,000448415508583941462695404240456206088416594346960639183292457732399327590760030791730548828125
163. 0,0002242077542919707313477021202281030442082971734803195916462288661996637953800153958652744140625
164. 0,00011210387714598536567385106011405152210414858674015979582311443309983189769000769793263720703125
165. 0,0056051938572992682836925530057025761052074293370079897911557216549915948845003848966318603515625
166. 0,00280259692864963414184627650285128805260371466850399489557861082749579744225019244831590017578125
167. 0,0014012984643248170709231382514256440263018573342519974477893054137478987211250962241579500878906

CLASSIS XV.

TETRADYNAMIA.

SILICULOSAE.

- * *Silicula integra, nec apice emarginata.*
- CCCVI. DRABA. *Silicula* valvulis planiusculis. *Stylus* subnullus.
- CCCXIV. LUNARIA. *Silic.* valvulis planis pedicellata. *Stylus* exsertus.
- CCCIV. MYAGRUM. *Silic.* valvulis concavis. *Stylus* persiftens.
- CCCV. ISATIS. *Silic.* lanceolata, monosperma, decidua.
- ** *Silicula emarginata apice.*
- CCCIX. IBERIS. *Petala* duo exteriora majora.
- CCCX. ALYSSUM. *Filamenta* quædam latere interiore dente notata. *Silicula* bilocularis.
- CCCXI. COCHLEARIA. *Silic.* cordata: valvulis obtusis gibbis.
- CCCVII. LEPIDIUM. *Silic.* cordata: valvulis acute carinatis.
- CCCVIII. THLASPI. *Silic.* obcordata: valvulis marginato - carinatis.
- CCCXIII. BISCUTELLA. *Silic.* biloba supra infraque, margine carinato.

SILIKUOSAE.

CCCXII. *CORONOPUS.* *Silic.* rotundato-re-
niformis, tuberculis atque
lamellis versus periphæriam
muricata.

S I L I Q U O S A E.

† *Calyx clausus foliolis longitudinaliter
dehiscens.*

CCCXXV. *RAPHANUS.* *Siliqua* articulata.

CCCXVIII. *ERYSIMUM.* *Siliqua* tetragona.

CCCXIX. *CHEIRANTHUS.* *Siliqua* germine
utrinque glandula notato.

CCCXX. *HESPERIS.* *Glandulae* intra stamina
breviora. *Petala* obliqua.

CCCXXI. *ARABIS.* *Glandulae* 4 intra foliola
calycina. *Stigma* simplex.

CCCXXIII. *BRASSICA.* *Glandulae* 2 intra
stamina breviora; 2 extra
stamina longiora.

CLASSIS XV.
SILICULOSA

CCCIV. MYAGRUM.

Linn. Gen. plant. ed. Schreb. n. 1069.

Calyx tetraphyllus: *foliolis* ovato-oblongis, concavis, dehiscentibus, coloratis, deciduis. *Corolla* tetrapetala cruciformis. *Petala* plana, subrotunda, obtusa: *unguibus* angustis. *Germen* ovatum. *Stylus* filiformis, longitudine calycis. *Stigma* obtusum. *Silicula* obcordata, subcompressa, integra, rigida, apice stylo rigido conico terminata, bivalvis: loculis quibusdam sæpe vacuis. *Semina* subrotunda.

957. **MYAGRUM** *rugosum* filiculis sulcatis pilosis rugosis, foliis oblongis obtusis dentatis. *Linn. Sp. pl. 2. p. 893.*

Myagrum filiculæ articulo primo stricto, altero globoso striato. *Hall. strip. helv. n. 525.*

Rapistrum monospermum. *Rauh. pin. 95. Prodr. 37. Icon.-Mapp. fl. Alf. p. 266. Icon optima.*

Germ. Runzlicher Leindotter.

Gall. Cameline.

Ha b. In agris ficcioribus et ruderatis passim prope *Kehl*. In Brisgovia in agris am *Kaiserstuhl* passim frequens. In Margraviatu superiore prope *Weil* Basileam versus. Circa Basiliam in agris ad Augusta Rauracorum; am *Rothenhaus*, nec non in agris circa *Klein-Hüningen*, in agris arenosis circa *Neudorf*, *Michelfelden* et *Maert* Rhenum versus. In Alsatia prope *Argentina* in agris *vor dem Fischerthor*, retro *der Citadelle* et alibi passim post *Mapum* vidi.

Flor. Majo, Junio. *Semina* Julio matura. ☉

Radix fusiformi-teretiufcula. *Caulis* pedalis, cubitalis, erectiusculus, sulcatus, villosus-hispidus, supra medium ramosus. *Rami* alterni, diffusi. *Folia* alterna, petiolata, dentato-pinnatifida, obtusa; ramea sessilia, ovalilanceolata, dentata, omnia hispida. *Flores* caulem ramosque terminantes, racemosi, numerosi. *Calycis* foliola oblonga, obtusa, viridi-flavescentia. *Petala* patentia, obtusa, subemarginata, lutea, calyce duplo longiora. *Siliculæ* biarticulatæ, rugosæ, articulo inferiore minore; superiore ovato, striato, villosus, stylo rostrato instructo, monospermo.

Obs. Planta omnino annua, non confundenda cum *M. perenni*. *Synonyma* citata, a *Linnaeo* ad *M. perenne* allegata, suadente *Lachenalio* et *Villarsio*. *Flor. Delph.* 3. p. 246, omnino hic pertinent.

958. *MYAGRUM perfoliatum* siliculis obcordatis subsessilibus, foliis amplexicaulibus. *Linn. Sp. pl.* 2. p. 893.

Myagrum loculo secundo conico, sterili biloculari. *Hall. stirp. helv.* n. 524.

TETRADYM. SILICULOSA.

7

Myagrum monospermon latifolium. *Bauh. Prodr.* 51. *Icon.* — *Mapp. fl. Als.* p. 206.

Germ. *Durchwachsener Leindotter.*

Gall. *Cameline perſévillée.*

Hab. In Agro Baſileenſi circa *Bielbencken* in agris t. *Stachelino* et *Lachenalio*; in Voſgeſi agro prope *Sanctum Mauritium* t. *Joh. Bauhino.*

Flor. Majo, Junio. *Semina* Julio matura. ☉

Caulis cubitalis et altior, erectus, teres, ſubſtriatus, glaber, glaucus, ramosus. *Rami* axillares, erectiusculi. *Folia* alterna, ſeſſilia, cordatolote lanceolata, amplexicaulia obtuſa ſubſinuato remotiuscule denticulata, glabra, glauca. *Flores* caulem ramosque terminantes, racemoſi. *Corollæ* parvulae, luteæ. *Siliculæ* obovato-cordatæ, adpreſſæ, ſtylo recto præditæ, monospermæ.

Nota. Ad rariores omnino Floræ noſtræ cives pertinet, in regionibus Galliæ meridionalis, Italiæ et Hiſpaniæ inter ſegetes vulgaris.

959. **MYAGRUM Bauhini** ſiliculis obovatis glabris polyspermis, ſoliis amplexicaulibus ſeſſilibus lineari-lanceolatis remote dentatis, glabris.

Myagrum dentatum. ſiliculis obovatis pedunculatis polyspermis, ſoliis amplexicaulibus glabris dentatis. *Willd. Linn. Sp. pl.* 3. p. 408.

Myagrum fativum. var. γ . *Linn. Sp. pl.* 2. p. 894. — *Pollich. fl. pal. n.* 602. var. γ .

Alyſſum ſoliis glabris, auriculatis, profunde dentatis. *Hall. ſtirp. helv. n.* 489. var. γ

3 **TETRADYN. SILICULOSA.**

Alysson foetidum. *Mapp. fl. Alf. p. 21.*

Myagrum foetidum. *Bauh. pin. 109. Prodr. p. 51.* — *Lindern. Tournesf. Alsat. p. 94. t. 1. (Icon mala. Folia sagittata margine integra ibi expressa.)*

Germ. Bauhins Leindotter.

Gall. Cameline de Bauhin.

Hab. In arenosis humidiusculis ad fossulas. In Margraviatu superiore ad pontem *Wiesæ* t. *C. Bauhino.* In Palatinatu inter *Weinheim* et *Gernsheim* locis cultis arenosis. t. *Pollichio.* In Alsatia retro *Obernheim* versus *Girrbaden* ad marginalia arida copiose, t. *Mappo.*

Flor. Majo, Junio. *Semina* Julio Augusto matura. ☉

Radix perpendicularis, tenuis, fibrosa. *Caulis* fesquipedalis et altior, erectus, teres, glaber, laete viridis, ramosus. *Rami* alterni, erecti, glabri. *Folia* alterna, sessilia, linearilanceolata, semiamplexicaulia, margine sinuato-dentata, 4 ad 5 uncias longa 4 ad 7, lineas lata, infima superioribus profundius inciso-dentata, omnia glabra, laete viridia. *Flores* caulem ramosque terminantes, racemoso-corymbosi. *Pedunculi* longiusculi, uniflori, erecti. *Corollæ* parvæ, pallide luteæ. *Capsulæ* orbiculato-obovatæ, stylo terminatæ, glabræ, biloculares, bivalves polyspermæ.

Nota. Specie omnino diversa a *Myagro* sativo.

Obs. Planta ultra quindecim jam annos ex seminibus in horto botanico Carlsruhano culta, multationi non succubuit.

TETRADYN. SILICULOSA.

960. **MYAGRUM** *sativum* filiculis obovatis polyspermis, foliis sessilibus amplexicaulibus subsagittatis, integris, utrinque villoso-scabris.

Myagrum sativum filiculis obovatis pedunculatis polyspermis. *Linn. Sp. pl.* 2. p. 894. — *Pollich. fl. pal. n.* 602.

Alyssum foliis sagittatis, siliquis cordiformibus. *Hall. stirp. helv. n.* 489.

Mœnchia sativa floribus racemosis, petalis integerrimis, siliculis obovatis. *Roth. fl. germ.* 2. p. 75.

Alysson. Bot. Paris. Mapp. fl. Alf. p. 20.

Myagrum sylvestre. Bauh. pin. 109.

1. *Myagrum sativum. Bauh. pin.* 109. — *Pollich. l. c. var. β.*

Icon. Cavanill. ic. 1. t. 66. — *Oeder. fl. Dan.* t. 1038. — *Schkuhr. bot. Handb. t.* 178.

Germ. Gemeiner Leindotter. Dotterkraut. Dort.

Gall. Sefame batard. Cameline cultivé.

Hab. Utrinque passim inter segetes, præsertim in Lini agris. *β.* colitur passim.

Flor. Majo, Junio. *Semina* Julio Augusto matura. ☉

Caulis pedalis, sesquipedalis et altior, erectus, obsolete angulatus, scaber, pilosus, ramosus. *Rami* alterni, erecti. *Folia* alterna, sessilia, sagittato-ovato-lanceolata, integra quandoque subdenticulata, semper utrinque villosa scabra. *Flores* terminales, racemose.

corymbosi. *Pedunculi* erecti, tenues, glabri uniflori. *Corollæ* pallide luteæ. *Petala* oblonga, integra, obtusa, calyce longiora. *Capsulæ* obovatæ, glabræ, stylo terminatæ, biloculares, bivalves, polyspermæ.

β. Differt *Foliis* manifestius denticulatis, minus villosis, *Capsulis Seminibusque* majoribus.

Pharm. MYAGRI f. CAMELINÆ Herba; Semen.

Ex feminibus paratur oleum mite edule, aurei fere coloris, facile rancescens. Semina aviculis, præsertim fringillis acceptissima.

961. *MYAGRUM paniculatum* filiculis lentiformibus orbiculatis, punctato rugosis. *Linn. Sp. pl.* 2. p. 894. — *Pölich. fl. pal. n.* 603.

Rapistrum foliis amplexicaulibus, auriculatis, acutis. *Hall. stirp. helv. n.* 522.

Rapistrum arvense, folio auriculato, acuto. *Tournef. inst. p.* 211. — *Mapp. fl. Als.* p. 267.

Myagro similis filicula rotunda. *Bauh. pin.* 109. *Prodr. p.* 52. *Icon.*

Icon. Oeder. fl. Dan. t. 204. — *Schkuhr. bot. Handb. t.* 178.

Germ. Rispenblütiger Felddotter.

Gall. Cameline paniculée.

Hab. Utrinque passim inter segetes. Inter *Carlsruhe* et *Beurtheim*; in agris et arvis submontosis auf dem *Thurnberg*, prope *Groezingen* et alibi non infrequens.

Flor. Junio, Julio. *Semina* Augusto matura. ☉

Caulis 2 — 3 pedalis, erectus, obsolete angulosus, scaber, superne ramosus. *Rami* inordinati, sparsi, erecti. *Folia* alterna, sessilia, amplexicaulia, utrinque scabra, margine integrifuscula, aut remote subdenticulata. *Flores* racemosi, caulem ramosque terminantes. *Racemi* simplices, erecti, longiusculi, nudi. *Corollæ* petala lutea, obtusa, parva, calyce sublongiora. *Siliculæ* lenticulari-rotundæ, punctato-rugosulæ, stylo præditæ, monopermæ.

962. *MYAGRUM saxatile* filiculis lentiformi-obovatis glabris, foliis radicalibus petiolatis ovato-lanceolatis subdenticulatis, caulinis sessilibus lineari-lanceolatis integris scabris.

Myagrum saxatile filiculis lentiformi-obovatis glabris, foliis oblongis serratis scabris, caule paniculato. *Linn. Sp. pl.* 2. p. 894.

Alyssum foliis scabris, radicalibus petiolatis, ovatis, dentatis, caule subnudo flexuoso. *Hall. stirp. helv. n.* 490.

Thlaspi alpinum minus capitulo subrotundo. *Bauh. pin.* 107. — *Prodr.* p. 48. *Icon. bona.*

Icon. Jacqu. *Fl. Austr.* 2. t. 128.

Germ. Weisblütiger Felsendotter..

Gall. Cameline des Roches.

Bab. In Ditione Basileensi in monte *Wasserfall* t. C. Bauhino; in montibus *Ramstein*,

penheimensi copiose post *Polichium* vidi. In Sponhemia in rupibus Basalticis prope *Martinstein* et alibi passim.

Flor. Majo, Junio. *Semina* Julio matura. ☞

Radix fusiformi-cylindrica, descendens. *Caulis* 2-3-4 pedalis, erectus, teretiusculus, glaberrimus, superne ramosus. *Folia* radicalia infimaque caulina subpetiolata, ovato-lanceolata, dentata, villosiuscula, reliqua alterna, sessilia, amplexicaulia, sagittato-lanceolata, integriuscula, glaberrima, ex viridi-glaucescencia. *Flores* caulem ramosque terminantes, copiosi, racemoso-corymbosi. *Corollæ* patentes, parvæ, luteæ. *Siliculæ* maturæ pendulæ, nigræ, splendentes, deciduæ, submonospermæ.

Pharm. ISATIS FOLIA.

Folia amariuscula, subadstringentia, hodie in medicina exoleta.

Ex foliis recentibus plantae maceratis, in pastam redactis, color coeruleus obtinetur, qui in arte tinctoria ad lanas tingendas adhibetur. Hinc in diversis Germaniæ, Galliæ, Angliæ regionibus colitur. De methodo Isatin colendi et præparandi, vide Astruc mem. pour. L'hist. de Languedoc p. 523. sq.

recta, integra, stylo subdestituta, bilocularis: *dissepimento* valvis parallelo: *valvulis* plano-concavis. *Semina* plura, parva, subrotunda.

† *Caule nudo.*

964. *DRABA aizoides* scapo nudo simplici, foliis ensiformibus, carinatis ciliatis. *Linn. Mant. gr.*

Draba cauliculis nudis, foliis linearibus ciliatis. *Hall. stirp. helv. n. 498.*

Moenchia aizoides, *Roth. fl. germ. 2. p. 72.*

Alyssum alpinum hirsutum luteum. *Tournef. inst. p. 217.*

Sedum alpinum hirsutum luteum. *Bauh. pin. 284.*

con. Jacqu. fl. Austr. 2. t. 20. f. 2. — Mill. Icon. 20. f. 2.

germ. Gelbblühendes Felsenhungerblümlein.

hall. Drave aizoides.

lab. In Ditione Basileensi, in rupibus calcareis montis Dornach, Vogelberg, circa Wallenburg, Langenbruk, Kallenstuh, Schaffmatt. In monte Wasserfall, frequens.

hor. Martio, Aprili, Majo. Semina Majo, Junio matura. ☞

Radix ramosa, multiceps. *Scapus* uncialis, digitalis et altior, erectus, teres, glaber, simplex. *Folia* radicalia numerosa, in rosulas congesta, sessilia, lineari-lanceolata, integra,

marginē utrinque, albide ciliata, plana, carina glabra, quinque ad novem lineas longa, lineam lata, firma. *Flores* scapum terminantes racemosi. *Calycis* foliola concava, patentia ex viridi-luteola. *Corollæ* petala oblonga, obtusa, quandoque submarginata, patentia, lutea, calyce duplo longiora. *Siliculæ* lanceolatae, acutæ, compressæ, flavicantes, villosiusculæ, *Stylō* erecto, glabro præditæ. *Semina* 5 — 9, ovata, compressa, flavescencia.

Nota. Sub *Draba alpina*. Linnaei Sp. pl. 2 p. 836. militabat.

Planta omnino pulchella, rupes præsertim altiores calcareos sequens; facile tamen in hortorum rupibus cicuratur.

965. *DRABA verna* spapis nudis, foliis lanceolatis subserrulato-denticulatis, petalis corollæ bipartitis.

Draba verna scapis nudis, foliis lanceolatis subincilis. Linn. Sp. pl. 2. p. 896. — Pollich. fl. pal. n. 604.

Draba cauliculis nudis, foliis subhirsutis, subdentatis. Hall. stirp. helv. n. 496.

Alysson vulgare Polygoni folio, caule nudo. Tournef. inst. p. 217. — Mapp. Alf. p. 20.

Bursa pastoris minor, loculo oblongo. Bauh. pin. 108.

Aline quinta minima. Tabernaemont. p. 1087. Icon.

Icon. Curtis. fl. Lond. — Oeder. fl. Dan. t. 983. — Schkuhr. bot. Handb. t. 179.

Germ. Hungerblümchen. Klein Gänsekrout. Gall. Drave printanniere.

Hab.

12

13

14



Hab. Utrunque in agris, pascuis macilentis, cultis, hortis, muris, ad viarum marginalia passim abunde.

Flor. Febuario, Martio, Aprili et seq. *Semina* Majo, Junio matura. ☉

Scapus uncialis, digitalis et altior, erectus, nudus, debilis. *Folia* radicalia in orbem congesta, terræ adjacentia, ovato-lanceolata, integrifuscula, quandoque plus minusve profunde serrato-dentata, villifuscula. *Flores* scapi terminantes, racemoso-corymbosi, demum racemosi. *Corollæ* exiguae, albæ. *Petala* calyce longiora, obtusa. *Siliculæ* ovaes, obtusæ, compressæ, integræ, *Stylo* destitutæ, glabræ, polyspermæ. *Semina* ovalia, compressa, fusca.

Ubi abundanter crescit, solum fere semper macilentum, sabulosum indicat. Arenam figit.

†† *Caule folioso.*

66. *DRABA muralis* caule ramoso, foliis cordatis dentatis, amplexicaulibus. *Linn. Sp. pl.* 2. p. 897. — *Pollich. fl. pal.* n. 605. — *Hall. stirp. helv.* n. 499.

Bursa pastoris major loculo oblongo. *Bauh. pin.* 108. — *Prodr.* p. 50. *Icon.*

Bursa pastoris sublongo loculo affinis planta. *Bauh. hist.* 2. p. 938.

Icon. propria.

Ger. m. Mauerhungerblümchen.

Call. Drave de murs.

Hab. Retro Græzingen ad viam asperam lapidosam calcaream dumetosam rarius; prope Dietlingen et Stein ad vias lapidasas cum

Thlaspi perfoliato non infrequens. In Marggraviatu superiore in asperis lapidosis, dumetosis prope *Weil* et *Crenzach* ad vineas. Basileæ extra portam *Spahlenthor* im *Fuhrengäfslein* t. *Zeyhero*; circa *Burglingen* ad *Birsam*, ad *St. Jacob*, ad Rheni ripam am *Rheinschänzlein*: prope *St. Margreth*, circa *Friedlingen*, *Reichen*, et inter portam *Steinenthor*, et viam *Mönchenstein*; in ipso pago *Binningen*, ad sepes umbrosas passim t. *Lachenalio*. In Alsatia prope *Barr* ad vinearum vias lapidosas dumetosas vidi. In Palatinatu in montibus altis virgultosis ac sylvosis. circa *Steinbach* ad rupes, t. *Pollichio*. In Sponhemia rupibus schistoso-argillaceis prope *Winnigen* ad *Mosellam*, prope *Burgsponheim*, *Naumburg*, *Winterburg*, *Martinstein* et *Oberstein*, non infrequens.

Flor. Aprili, Majo. *Semina* Junio, Julio matura. ☉

Caulis digitalis, palmaris, pedalis, erectus, hirsutus, simplex aut ramosus. *Folia radicalia* petiolata, ovata dentato-ferrata, *caulina* alterna, amplexicaulia, sessilia, cordato-ovata, dentata, omnia hirsuta. *Flores* caulem ramosque terminantes, racemosi. *Corollæ* parvæ, albidae. *Petalæ* subrotunda, calyce longiora. *Siliculæ* lineari-ellipticæ, glabræ, compressæ, 6-8-10 spermæ.

Obs. Hujus plantæ specimina sesquipedalia, ramosa, ad Pyrenæorum radices in Catalonia vidi.

CCCVII. L E P I D I U M.

Linn. Gen. pl. ed. Schreb. n. 1077.

Calyx tetraphyllus: *foliolis* ovatis, concavis, deciduis. *Corolla* tetrapetala, cruci-

lms. *Petala* obovata, calyce duplo longiora, unguibus angustis. *Silicula* cordata, emarginata, compressa, margine acuta, bilocularis: *calculis* navicularibus, carinatis, *dissepimento* lanceolato contrariis. *Semina* nonnulla, ovato-acuminata, basi angustiora, nutantia.

967. *LEPIDIUM sativum* floribus tetradynamis, foliis oblongis multifidis. *Linn. Sp. pl.* 2. p. 899.

Lepidium foliis varie divisis incisisque. *Linn. Hort. Cliff.* 331.

Nasturtium hortense vulgatum. *Bauh. pin.* 103.

Nasturtium. *Tabernæmont. p.* 841. *Icon.*

β. *Nasturtium hortense crispum angustifolium* et *latifolium.* *Bauh. pin.* 104. *Prodr. p.* 43. 44. *Icon.*

Nasturtium crispum. *Tabernæmont. p.* 842. *Icon.*

Icon. Regnault. Bot. — Blackw. herb. t. 23.

Ger. Garten-Kresse. Zahme Kresse. β. Krausse Garten-Kresse.

Call. Le Cresson Alenois.

Hab. Colitur in hortis culinariis et vineis passim frequens. In ruderatis ad urbes et pagos ex hortorum rejectamentis quasi spontanea.

Flor. Majo, Junio, Julio. *Semina* Julio, Augusto matura. ☉

Variat. Foliis angustioribus et latioribus plus minusve divisis, planis et crispatis.

Pharm. *NASTURTII HORTENSIS* *Hertl* recens; *Semen.*

Planta irritans, acriuscula, antiscorbuticas et di-
reticas vires exercet, in culinis notissima.

Semina quovis anni tempore facillime germinant.

968. *LEPIDIDIUM latifolium* foliis ovato-lanceolatis integris serratis. *Lin.*
Sp. pl. 2. p. 899. — *Hall. stirp. he*
n. 505.

Lepidium latifolium. *Bauh. pin.* 97.

Lepidium Aeginetæ. *Tabernemont. p.* 84
Icon.

Icon. *Regnault. bot.* — *Blakw. t.* 448.

Germ. *Pfefferkraut.* *Bitterkraut.* *Fleisch-*
kraut.

Gall. *La Passerage.*

Hab. In Principatus superioris vicinia *Für-*
stenbergensi. In rupibus *Fürstenbergi*
vulcanicis præaltis præruptis ad *Hoheneg-*
prope *Engen* ubi maximam ejus abundan-
tiam vidi. Colitur in hortis culinariis.

Flor. Junio, Julio, Augusto. *Semina* Augusto
Septembri maturant.

Caulis cubitalis, tripedalis et altior, erectus, glaber, ramosus. *Folia* radicalia caulinaque ovato-lanceolata, obtusa, margine serratis utrinque glabra, superiora rameaque lanceolata, acuta, integra. *Flores* caulem ramosque terminantes, paniculati. *Corollæ* parvae, albæ. *Siliculæ* subrotundæ, compressæ, submarginatæ.

Pharm. LEPIDII LATIFOLII Radix. Folia recentia.

Folia mafficata acris. amara, antiscorbuticas et emeticas vires exercent. Sale culinari et aceto condita in uentis cum carnibus comeduntur; Appetitum florum excitant. Radix ac folia pro Sinapismo contra dolorem ischiadicum laudantur.

969. **LEPIDIUM graminifolium** foliis linearibus: superioribus integerrimis, caule paniculato virgato, floribus hexandris. *Linn. Sp. pl. 2. p. 900.*

Lepidium Iberis Poll. fl. pal. n. 607.

Lepidium Pollichii foliis inferioribus lato-lanceolatis ferratis; superioribus linearibus subintegerrimis, filiculis orbiculatis emarginatis. *Roth. fl. germ. 2. p. 91.*
— *Willden. Linn. Sp. pl. 3. p. 441.*

Masturtium foliis imis pinnatis incis, superioribus linearibus integerrimis. *Hall. firs. helv. n. 507.*

Iberis Cammerar. Epitom. p. 184. (cum foliis infimis bene expressis.)

Iberis secunda. Tabernæmont. p. 848. Icon optima, (absque foliis infimis.)

Ger. Schmalblättrige Kresse.

Call Passerage graminiforme.

Hab. In Palatinatu Badensi prope *Heidelberg* frequens. In Palatinatu transrhenano ad vias ac in plateis desertis circa *Dürkheim, Ungstein, Callstatt* abunde; circa *Oppenheim, Nierstein, Guntersblum, Worms* et alibi post *Pollichium* vidi. In *Sponheim* ad pagorum vias prope *Winningen* ad *Mosellam*; prope *Weiler, Martinsstein, Sprendlingen* et alibi passim abunde.

Flor. Junio, Julio, Augusto, Septembri et Octobri. *Semina* ab Augusto ad Novembrem maturantur. 7

Radix perennis fusiformis, subramosa, extus fusco-lutea, intus albida. *Caulis* 2-3 pedalis, adscendens, erectus, obsolete angulosus, glaber, ramosus. *Rami* alterni, numerosi, divergentes. *Folia radicalia* infimaque passim caulina, longe petiolata, pinnatifida lacinia terminali maxima, ovata, aut ovali lanceolata, quandoque non pinnatifida, sessivalia, seu ovali-lanceolata serrata. *Caulina* reliqua *ramea* alterna, sessilia, linearia, integerrima, glabra, patentia, aut deflexa, læte viridia. *Flores* racemosi, caulem ramosque terminantes. *Racemi* erecti, aphylli. *Calyx* tetraphyllus, *foliolis* purpurascens, binis latiusculis et binis angustioribus. *Corolla* *Petal* *tala* quatuor, aequalia, subrotunda, obtusiuscula, alba, calyce majuscula. *Stamina* sex, quatuor longiora et duo breviora, omnia *Antheris* subrotundis, luteis perfectis instructa. *Silicula* elliptico-subrotunda, compressa, glabriuscula, bilocularis, bisperma. *Stylo* breviter obtuso persistente coronata. *Semen* ovatum, ex flavicante fuscum.

Obs. Specimina abs me in locis citatis aliisque regionibus Gallie meridionalis examinata, semper staminibus sex instructa fuerunt.

Individua ex Seminibus in horto Botanico ultra quindecim jam annos culta, perennia, mutationi non succubuerunt.

Folia radicalia pinnatifida in plantis spontanei tempore inflorescentie et fructificationis, facile evanescent, tunc omnia sæpe linearia, integerrima sunt.

Nota. Non pro peculiari specie a L. graminea Linnaei diversa declaranda, nec pro L. lbe

ris Linn. habenda, quæ est planta annua, pedalis, minus ramosa, Foliis radicalibus caulinisque omnibus et passim rameis lanceolatis ferratis instructa. Præterquam Floribus dian-dræis, et Siliculis emarginatis Stylo destitutis, prædita est.

Radix plantæ nostræ masticata, acris, saporis Cochleariæ Armoraciæ Linn. Forſan Iberis veterum Botanicorum, et Medicorum, cujus Radice contusa morbis Ischiadicis mirum in modum medebantur. Servilius Damocrates plantam invenit, quam carmine celebratam dixit Iberida, circa parietinas. Semper floret, foliis Nasturtii, caule cubitali, semine minimo, radice nasturtii odore, ad coxendicum rheumatismos utilis, Plin.

970. *LEPIDIUM ruderale* floribus di-andris apetalis, foliis radicalibus dentatopinnatis; ramiferis linearibus integerrimis. Linn. *Sp. pl.* 2. p. 900. — *Poll. fl. pal.* n. 606.

Nasturtium apetalum foliis pinnatis, radicalium pinnis semipinnatis, caulinarum simplicibus. *Hall. stirp. helv.* n. 508.

Nasturtium sylvestre tenuissime divisum. *Mapp. fl. Als.* p. 209.

Nasturtium sylvestre, Oxyridis folio *Bauh. pin.* 105.

Thlaspi minus germanicum. *Tabernæmont.* p. 853. *Icon.*

Icon. Oeder. fl. Dan. t. 184. — *Schkuhr, bot. Handb. t.* 180.

Germ. Stink-Kresse. Klein Besenkraut.

Gall. Passerage ruderale.

La b. Utrunque ad urbes et pagos, in ruderalis, locis et plateis desertis fabulosis, passim frequens.

Flor. Junio, Julio, Augusto. *Semina* Septembri, Octobri matura. ☉

Caulis palmaris, pedalis et altior, erectus, teres glaber ramosus. *Rami* alterni, copiosi, tenues, virgati. *Folia* radicalia infimaque passim caulina pinnata seu pinnatifida, plus minusque incisa, *caulina* reliqua et *ramea* alterna, sessilia, linearia, integerrima, glabra, laete viridia. *Flores* caulem ramosque terminantes, racemosi. *Racemi* erecti, aphylli desflorati elongati. *Calyces* tetraphylli, erecti, conniventes, exigui. *Corollæ Petala* nulla, quandoque minutissima, vix conspicua. *Stamina* duo, rarius quatuor ad sex. *Siliculæ* subrotundæ, compressæ, glabrae, apice emarginatæ, *Stylo* destitutæ, bivalvès, bispermæ. *Semina* ovata, compressa, flavicantia.

Loca sabulosa præsertim ad urbes et pagos sequitur, ad sabulum figendum apta. Odor totius plantæ fortis, sapor acris. Cimices gravi odore pellere fertur. Siccata ad Scopas minores, hinc inde adhibetur.

CCCVIII. T H L A S P I.

Linn. Gen. pl. ed. Schreb. n. 1078.

Calyx tetraphyllus: *foliolis* ovatis concavis, erecto patulis, deciduis, *Corolla* tetrapetala, cruciformis: *Petala* obovata, calyce duplo longiora, unguibus angustis. *Silicula* compressa obcordata, emarginata *Stylo* instructa, bilocularis: *dissepimento* lanceolato, *valvulis* navicularibus, marginato - carinatis. *Semina* plurima nutantia, futuris affixa.

971. *THLASPI arvense* filiculis orbiculatis, foliis oblongis dentatis glabris.

Linn Sp.pl. 2. p. 901. — Pollich. fl. pal. n. 608. Hall. stirp. helv. n. 511. (sub Nasturtio.)

Thlaspi arvense filiquis latis. Bauh. pin. 105. — Mapp. fl. Alf. p. 301.

Thlaspi majus. Tabernæmont. p. 849. Icon. Icon. Curtis fl. Lond. — Blackw. t. 68.

Germ. Ackertäschelkraut.

Gall. Tabouret des Champs.

Hab. Utrinque in cultis, agris, vineis, hortis, inter segetes passim nimis copiose.

Flor. Majo, Junio. Semina Julio Augusto matura. ☉

Caulis palmaris, pedalis, erectus, angulato-fulcatus, glaber, ramosus. Rami alterni. Folia radicalia petiolata, ovata, integriuscula, caulina alterna, sessilia, subsemiamplexicaulia, sagittata, ovata, sinuato-dentata, omnia glabra, virescentia, quandoque subflavescentia. Flores racemosi, caulem ramosque terminantes. Corollæ parviusculæ, albæ. Siliculæ orbiculatæ, compressæ, margine membranaceo lato cinctæ, emarginatæ, glabræ, Stylo brevi instructæ. Semina plurima, ovata, striata, flavicantia.

Pharm. THLASPEOS. Semina.

Planta ut omnes es hac Classe, antiscorbutica. Vaccinum lac. butyrum, carnes sapore ingrato inficit. Agrorum sæpe vitium. In granariis servata vermes frumento infestos fugat. Semina hodie in medicina exoleta.

972. *THLASPI campestre* filiculis subrotundis, foliis sagittatis dentatis incanis.

Linn. Sp. pl. 2. p. 902. — Pollich. fl. pal. n. 690.

Nasturtium foliis imis petiolatis ovatis, caulinis sagittatis dentatis. Hall. stirp. helv. n. 509.

Thlaspi vulgatius. Bauh. hist. 2. p. 921. — Mapp. fl. Alf. p. 301.

Thlaspi arvense Vacariæ incano folio majus et minus. Bauh. pin. 106. Prodr. p. 47.

Thlaspi secundum. Tabernæmont. p. 849. Icon.

β. Folio glabro minus ferrato.

Thlaspi arvense acetosæ folio. Bauh. pin. 105.

Thlaspi majus tertium. Tabernæmont. p. 850. Icon.

Icon. α. Curtis, fl. Lond. t. 303. β. Blackw. t. 407.

Germ. Feldtäfelkraut.

Gall. Tabouret vélue.

Hab. In ruderatis ad pagos et urbes passim; in sabulosis virgultosis ad rivos præsertim in Brigovia prope Freyburg abunde. In campestribus argillosis et in vinearum colibus hinc inde non infrequens.

Flor. Majo, Junio. Semina Julio, Augusto matura. ♂

Caulis palmaris, pedalis cubitalis et altior, erectus, teres, superne ramosus, utrinque pubescens, incanus, mollis. Rami erecti, sparsi. Folia radicalia numerosa, petiolata, ovata, integra, sæpius subpinnatifida, caulina alterna, sparsa, sessilia, semiamplexicaulia, sagittata, obtusiuscula, denticulata, tomentoso-pubescen-

tia, incana, mollis. *Flores* caulem ramosque terminantes, racemosi. *Racemi* erecti, densi. *Corollæ* Petala parva, albida, vix calycæ majora. *Silicula* subrotunda, compressa, emarginata, *Styl*o recto, longitudine emarginaturæ. *Semina* oblonga, nigricantia.

Planta ad sabulum figendum apta, præsertim in locis inundationibus expositis.

975. *THLASPI montanum* filiculis obcordatis, foliis glabris, radicalibus carnosis obovatis integerrimis; caulinis amplexicaulibus, corollis calycæ majoribus. *Linn. Sp. pl. 2. p. 902.* — *Pollich. fl. pal. n. 610.*

Lepidium caule erecto, foliis radicalibus petiolatis ovatis, caulinis amplexicaulibus sagittatis. *Hall. stirp. helv. n. 518.*

Bursa pastoris montana Globulariæ folio. *Tournef. in fl. — Mapp. fl. Alf. p. 46.*

Thlaspi alpinum bellidis cœruleæ folio. *Bauh. pin. 106.*

Thlaspi undecimum album supinum. *Tabernaemont. p. 852. Icon.*

Icon. Jacq. fl. Austr. t. 237.

Germ. Berg-Täschelkraut.

Gall. Tabouret des montagnes.

Hab. In montosis sylvaticis dumetosis subgraminosis lapidosis. In Margraviatu superiore ad montem prope *Krenzach* rarius. In Ditione Basileensi in montibus umbratis lapidosis *Muteto*, *Callen*, *Vogelberg* et supra *Dornach*, nec non in monte *Wasserfall*. In Principatus superioris vicinia *Fürstbergen* prope *Engen* ad *Krügenthal* cum *Viola* mirabili abundanter hanc

plantam vidi florentem. In Palatinatu circa *Steinbach* ad montem *Donnersberg* post *Pollichium* eandem observavi.

Flor. Aprili, Majo. *Semina* Junio, Julio matura. ¶

Radix ramosa, fibrosa. *Caules* ex radice sæpe plures, digitales, palmares, et altiores, erecti, ad basin curvuli, teretes, glabri, simplices, herbacei. *Folia radicalia* in rosulam congesta, in petiolum desinentia, obovata, obtusa, integra, quandoque subdentata, crassiuscula, glabra, viridia; *caulina* alterna, minora, sessilia, amplexicaulia, cordato sagittato-aut ovato-lanceolata, integra, aut obsolete crenulata, glabra, glauco-viridia. *Flores* corymboso-umbellati, demum racemosi caulem terminantes. *Calyces* ex viridi-purpurascens, margine albi. *Corollae Petala* quatuor, obovata, obtusa, plana, æqualia, calyce duplo longiora. *Silicula* obcordata, compressa, glabra, *Stylo* recto persistente prædita, inbetrasperma. *Semina* elliptico-orbiculata, compressa, glabra, fusco-lutescentia.

Obs. Specimina nostratia omni ex parte conveniunt cum Icone ab Ill. Jacquino l. c. exhibita.

In Horto culta et cicurata, mutationi non succubuit.

Ad rariores Floræ nostræ cives pertinet.

974. *THLASPI perfoliatum* siliculis obcordatis, foliis caulinis cordatis glabris subdentatis, petalis longitudine calycis, caule ramoso. *Linn. Sp. pl. 2. p. 902.* — *Pollich. fl. pal. n. 611.*

Nasturtium foliis imis ovatis, reliquis amplexicaulibus cordatis. *Hall. stirp. helv. n. 510.*

TETRADYN. SILICULOSA. 29

Thlaspi arvense perfoliatum majus. *Bauh. pin.* 106. — *Mapp. fl. Alf. p.* 302.

Thlaspi oleraceum. *Tabernæmont. p.* 851. *Icon.*

Icon. Jacqu. fl. Austr. t. 337.

German. Großes ästiges durchwachsenes Tackschelkraut.

Gall. Tabouret persévillée.

Hab. Utrinque in agris et vineis calcareo - argillaceis præsertim collicolis et montosis passim frequens. In agris et vineis prope *Durlach*, *Græzingen*, *Bruchsal*, *Heidelberg* et alibi. In Margraviatu superiori passim abunde. In Alsatia vineis et agris lutosi calcareis prope *Barr.* In Palatinatu in vineis circa *Dürkheim*, in aggeribus Rhenum versus inter *Oppenheim* et *Gernsheim* abundanter post *Pollichium* vidi.

Flor. Aprili, Majo. *Semina* Majo, Junio maturæ. ☉ ➤

Caulis palmaris, pedalis, erectus, basi inæquus, teres, glaber, ramosus. *Rami* alterni, simpliciusculi, quandoque divisi. *Folia radicalia* in petiolum desinentia, ovali-subrotunda, subdenticulata, glabra; *caulina* alterna, sessilia, amplexicaulia, cordata, integra, quandoque subdenticulata, glabra, omnia ex viridi glauca. *Flores* racemosi, caulem ramosque terminantes. *Racemi* laxi, erecti. *Corollæ* *Petalæ* calyce majuscula, albida. *Siliculæ* obcordatæ, glabræ, suboctospermæ.

975. *THLASPI alpestre* siliculis obcordatis, foliis subdentatis: caulinis amplexicaulibus, petalis longitudine calycis, caule simplici. *Linn. Sp. pl.* 2. p. 903.

30 TETRADYN. SILICULOSA.

Lepidium caule erecto, foliis radicalibus ovatis petiolatis, caulinis ovatis amplexicaulibus. *Hall. stirb. helv. n. 519.*

Thlaspi perfoliatum minus. *Bauh. pin. 106.*
— *Mapp. fl. Alf. p. 302.*

Thlaspi pumilum tertium. *Clus. hist. 2. p. 131. Icon.*

Thlaspi Clusii minimum. *Tabernæmont. p. 854. Icon ex Clusio.*

Germ. Kleines einfaches durchwachsenes Täschelkraut.

Gall. Tabouret perfévillée petite.

Hab. In rupestribus et dumetosis lapidosis asperis calcareis præsertim. In dumetis ad viarum lătera lapidosa auf dem *Thurnberg*, retro *Græzingen*, prope *Stein*, *Elmendingen* et *Dietlingen* cum *Draba murali*. In rupestribus præaltis prope *St. Goar* ad *Rhenum* non infrequens.

Flor. Aprili, Majo. *Semina* Majo, Junio matura. ☉

Differt a præcedente specie affini, *Radice* minore; *Caule* digitali, palmari, vix altiore, graciliore, magis stricto, simplici quandoque tantum ramo unico instructo; *Foliis* minoribus, *Racemo* minore, pauci floro, *Floribus* minoribus, *Siliculis* subtetraspermis.

Obs. Planta ex seminibus in horto culta, constans, non in præcedentem speciem abiit.

976. *THLASPI Bursa Pastoris* filiculis obcordatis, foliis radicalibus pinnatifidis. *Linn. Sp. pl. 2. p. 903.* — *Pollich. fl. pal. n. 612.*

Nasturtium filiquis triangularibus. *Hall. stirp. helv. n. 514.*

TETRADYN. SILICULOSA. 31

Bursa pastoris major folio sinuato. Bauh. pin. 108. — Mapp. fl. Alf. p. 46.

Bursa pastoria major. Tabernamont. p. 512. Icon.

2. *Bursa pastoris eleganti folio, instar Coronopi repentis. Camerar. hort. 32. Mapp. fl. Alf. p. 46.*

3. *Bursa pastoris major folio non sinuato. Bauh. pin. 108. Mapp. l. c.*

4. *Bursa pastoris media. Bauh. pin. 108. — Mapp. l. c.*

Bursa pastoria minor. Tabernamont. p. 512. Icon.

Icon. Curtis. fl. Lond. — Bulliard. Herb. t. 223. — Oeder. fl. Dan. t. 729. — Reg. mult. Bot.

Germ. Täschelkraut. Hirten-Täschelkraut. Hirtentafche.

Call. Le Tabouret. Tabouret bourse à pasteur.

Hab. Utrinque in cultis, hortis, oleraceis, æris, vineis ruderatis et viis frequens.

Flor. per totam æstatem. Semina a Junio ad Novembrem maturat. ☉

Caules ex radice plures, palmares, pedales, cubitales, erecti, basi sæpe curvuli, obsoletæ angulosi, plus minusve pubescentes, ramosi. Folia radicalia terræ incumbencia, in petiolum desinentia, pinnatifida, segmentis lanceolatis, aut ovatis integrusculis aut denticulatis, caulina alterna, sessilia, amplexicaulia, sagittato-lanceolata, dentata, aut integra, omnia plus minusve hirsuta, viridia, quandoque incana. Flores caulem ramosque terminantes, racemosi. Corollæ parvæ albidæ. Siliculæ obcordatæ, triangulares, compressæ, margine de-

32 TETRADYN. SILICULOSA.

stitutæ, glabræ, quandoque villosiusculæ, polyspermæ.

Planta polymorpha, foliorum structura maxime varians.

Memorabilis degeneratio Foliis omnibus crispatis, undulatis, hirsutissimis, incanis, Calycibus monstrosis, magnis, dense tomentosis, a Cynipe quodam producta, ad muros hinc inde circa Carlsruhe crescit.

Pharm. BURSÆ PASTORIS Herba.

Hodie fere in medicina exoleta, olim præsertim in hæmorrhagiis et profusionibus sanguinis pulmonum et uteri maxime in usu fuit.

Herba recens una cum oleosis, sanat Hæmaturiam Bobus et Tauris, si tempestive adhibita fuerit. Scopol. fl. Carn. 2. p. 17.

Planta tenella ovibus accepta.

CCCIX. IBERIS.

Linn. Gen. plant. ed. Schreb. n. 1080.

Calyx tetraphyllus: *foliolis* obovatis, concavis, patentibus, æqualibus, parvis, deciduis. *Corolla* tetrapetala, inæqualis. *Petala* obovata, obtusa, patentia, *unguibus* oblongis, erectis: horum *duo exteriora proxima Petala* longe majora, inter se æqualia; *duo interiora minima*, reflexa. *Silicula* erecta, suborbiculata, compressa, emarginata, margine acuto cincta, bilocularis; *Dissepimento* lanceolato; *valvulis* navicularibus, compressis, carinatis. *Semina* nonnulla subovata.

977. *IBERIS amara* herbacea, foliis lanceolatis acutis subdentatis, floribus racemosis. Linn. Sp. pl. 2. p. 906. — Pollich. fl. pal. n. 614.

Iberis

TETRADYN. SILICULOSA. 33

Iberis foliis dilatatis, dentatis, floribus umbellatis. *Hall. stirp. helv. n. 520.*

Thlaspi umbellatum arvense amarum. *Bauh. hist. 2. p. 925. — Mapp. fl. Als. p. 301.*

Thlaspi amarum. Tabernamont. p. 851. Icon.

Ger. Bitterer Bauernsenf.

Gall. Iberide amère.

Hab. In agris et arvis præsertim calcareo-argillaceis. In arvis retro dem *Thurnberg* et prope *Grözingen* non infrequens. In *Marggraviatu* superiore inter segetes prope *Lippurg* frequens, nec non inter *Weil Haltingen* et alibi. In arvis cis et transrhenanis passim. In Palatinatu in agris circa *Esselborn*, *Alzey*, *Oppenheim*, prope dem *Holzhof* t. *Pollichio*.

Flor. Julio, Augusto. *Semina* Septembri, Octobri matura. ☉

Radix fusiformis, fibrosa, tenuis, albida. *Caulis* palmaris, pedalis, et altior, erectus, angulatus, villosiusculus, ramosus. *Rami* sparsi, alterni, oppositique, diffusuli. *Folia* alterna, sessilia, cuneiformi-lanceolata, obtusa, antice denticulata, *ramea* superiora integra, omnia crassiuscula, glabra. *Flores* caulem ramosque terminantes, corymbofo-racemosi, umbellam imitantes. *Corollæ Petala* duo exteriora, interioribus duplo longiora, nivea, *Siliculæ* ellipticæ, emarginatæ, margine alatæ, bispermæ.

Planta florens ob copiosos flores pulchella, verum horum decus.

978. *IBERIS nudicaulis* herbacea, foliis sinuatis, caule nudo simplici. *Linn. Sp. pl. 2. p. 907. — Pollich. fl. pal. n. 615.*

Iberis foliis pinnatis, pinnis ovatis acutis
Hall. stirp. helv. n. 521.

Nasturtium petræum, foliis Bursæ pastoris
Bauh. pin. 104. — Mapp. fl. Alf. p. 209.

Nasturtium petræum. Tabernæmont. p. 842.
Icon.

Icon. Oeder. fl. Dan. t. 323. — Schkuhr. bot.
Handb. t. 179.

Germ. Sand- Iberkraut.

Gall. Iberide nudicaule.

Hab. Utrunque in pascuis arenosis sterilibus
apricis, locisque sabulosis sylvaticis in viis.
Circa *Carlsruhe* im *Hardtwald* passim fre-
quens. Inter *Mühlberg* et *Dachsland* in
arenosis collibus abunde. In Margraviatu
superiore prope *Weil* et *Haltingen* in
agris sabulosis.

Flor. Martio, Aprili, Majo. *Semina* Majo
Junio matura. ☉

Caulis digitalis, palmaris, erectus, teres,
nudus, glaber tenuis. *Folia radicalia* in roset-
tam congesta, terræ incumbencia, in petiolum



plana, calyce subbreviora, patentissima, integra aut emarginata. *Filamenta* breviora denticulo introrsum basi inserto. *Silicula* in aliis emarginata, in aliis integra, compressa aut ventricosa, subglobosa, *Stylo* longitudine filiculæ instructa, bilocularis: *Dissepimento* elliptico; *valvulis* elliptico-hemisphæricis. *Semina* nonnulla, orbiculata.

979. *ALYSSUM montanum* ramulis suffruticosis diffusis, foliis punctato-echinatis. *Linn. Sp. pl.* 2. p. 907.

Alyssum suffruticosum foliis ellipticis scabris. *Hall. stirp. helv. n.* 492.

Alyssum ramulis suffruticosis diffusis, foliis obverse lanceolatis tomentosis. *Jacqu. Vind.* 258.

Adyseton montanum. *Scopol. fl. Carn.* 2. p. 14.

Alysson perenne, montanum, incanum. *Tournef. inst. — Mapp. fl. Als.* p. 20.

Thlaspi montanum luteum. *Bauh. hist.* 2. p. 928. — *Bauh. prodr. p.* 49. *Icon.*

Icon. Iacqu. fl. Ausir. t. 37.

Germ. Bergsens. Felsensens.

Gall. Alysse des montagnes.

Hab. In Brisgovia monte *Kaiserstuhl* in rupibus præaltis vulcanicis ad arcem *Limburg*, nec non in rupibus prope *Sponeck* et *Büdingheim*. In Margraviatu superiore ad rupes prope *Rhenum* infra *Efringen*. In Ditione Basileensi in rupibus circa *Birsek* t. *C. Bauhino* et *Lachenalio*. In vicinia Principatus superioris im *Hegau* ad rupes et muros

Castelli *Hohentwyl* abunde vidi. In Alsatia ad rupes erga *Niedeck im Hochgebürg t. Mappo*.

Flor. Majo. *Semina* Junio, Julio matura. *Suj frutex*.

Radix ramosa, fibrosa, tenax. *Caules* e radice plures, palmares, et altiores, basi decumbentes, adscendentes, simpliciusculi, quandoque ad basin ramosi, suffruticosi, incani. *Folia* caulina alterna, numerosa, sessilia, obovato-lanceolata, basi attenuata, superiora sublinearilanceolata, obtusa, crassiuscula, integra incana, micantia, scabra, semiunciam ad unciam longa. *Flores* cauliculos terminantes corymbofo-racemosi, demum racemosi. *Calyces* pallide flavescentes, scabriusculi, decidui. *Corollae Petala* emarginata, lutea, calyce longiora. *Filamenta* aliqua ad basin squama subtridentata instructa. *Siliculae* ovato-orbiculatae compressae, *Stylo* praeditae, glabriusculae, bispermae.

Planta in rupibus peramoena, tempore inflorescentiae ob copiosos flores luteos e longinquo conspicua.

Obs. In Horto culta mutationi non succubuit.

Individua in altioribus regionibus Helvetiae et in Pyrenaeis observata, nostratibus demissiora, et magis incana sunt.

Folia per lentem spectata, stellulis villosis micantibus pulcherrimis instructa sunt.

980. *ALYSSUM arenarium* caulibus suffruticosis tortuoso diffusis, foliis subcuneiformi-lanceolatis, obtusis incanis, flaminibus subdentulis.

Alyssum campestre. *Pollich. fl. pal. n. 618. exclusis Synonymis.*

Adyseton montanum? *Wetterauische flor.*
2. p. 425.

Moenchia campestris. *Roth. fl. germ.* 2.
p. 74.

Alyssum supinum luteum. *Lobel. Icon.* p.
220.

Germ. Gelbbühender Sandfensf.

Gall. Alysse de sable.

Hab. In collibus arenosis apricis sterilibus sic-
cis. In Margraviatu superiore ad montem
Kaiserstuhl Brisgoviae prope *Sasbach* ver-
sus *Limburg* frequens. In Palatinatu ad
vias arenosas inter Mainz et *Finden*; pro-
pe *Nieder-Engelheim*: inter *Benzheim* et
Darmstadt in locis arenosis post *Pollichium*
vidi. In Sponhemia prope *Martinstein* cum
Stipa pennata et *Melica ciliata*, nec non
in collibus aridis *auf der Abtey*.

Flor. Majo, Junio. *Semina* Junio, Julio ma-
tura. *Suffructex.*

Maxime affinis praecedenti a qua differt,
Caulibus magis basi ad terram diffusis, tortuo-
sis, gracilioribus, *Racemis* tenuioribus. *Corollis*
subminoribus. *Petalis* obtusis, minus emargi-
natis. *Siliculis* minoribus, magis orbiculatis,
obtusis, subemarginatis.

Differt ab *Alyssō campestri* *Linnaei*, *Radice*
perenni lignosa, *Caulibus* a basi ad medium tenaci-
bus lignosis altioribus, *Racemis* longioribus. *Co-*
rollis majoribus, *Petalis* calyce duplo longiori-
bus. *Filamentis* saepe denticulatis, absentia *Se-*
tarum receptaculo insertarum.

Nota *Alyssum* campestre *Linn.* in regionibus
campestribus Galliae et Hispaniae calidioribus, vi-
x unquam in Germania crescit.

Non confundendum cum *Alyssō calycino* *Linn.*,
a quo differt, *Radice* non annua, *Caulibus* non

herbaceis, Calycibus non persistentibus, Corollis luteis, tabescentibus non albis, calyce duplo longioribus.

Planta ad arenam in collibus figendam apta.

981. ALYSSUM calycinum caulibus herbaceis, calycibus persistentibus, corollarum petalis luteis, demum albidis.

Alyssum calycinum caulibus herbaceis, staminibus omnibus dentatis, calycibus persistentibus. *Pollich. fl. pal. n.* 617.

Alysson foliis ellipticis scabris, calycibus persistentibus. *Hall. stirp. helv. n.* 494.

Adyseton calycinum. *Scopol. fl. Carn. 2.* p. 13.

Alysson incanum serpylli folio minus. *Tournef. infl. p.* 217. — *Mapp. fl. Alf. p.* 20.

Thlaspi XVIII. clypeatum minus secundum. *Tabernaemont. p.* 853. *Icon.*

Icon. Jacqu. fl. Austr. t. 338.

Germ. Gemeines Steinkraut.

Gall. Alysse caliculée

Hab. Utrunque in locis arenosis sterilibus, elevatis et demissioribus apricis, aridis frequens.

Flor. Aprili, Majo, Junio. *Semina* Julio, Augusto matura. ☉

Caules ex radice plures, digitales, palmares et altiores, basi decumbentes, demum erectiusculi, teretes, firmi, asperi, canescentes, simplices aut subramosi. *Folia* alterna, subsessilia, sparsa, lanceolata, obtusa, integra, crassiuscula, aspera, ex viridi incana. *Flores* caulem terminantes, corymbum primo imitantes, demum racemosi. *Calyces* persistentes,

lipiduli, incani. Corollæ Petala calyce paulo longiora, parva, primo lutea, demum albida. *Filamenta* quatuor longiora (in nostris saltem individuis) nullis denticulis instructa, duo autem breviora denticulata. *Siliculæ* orbiculatæ compressæ, obtusæ, subemarginatæ, *Stylo* breviter instructæ, 3-4 spermæ, piloso-asperæ.

Nota. Variat in montosis altioribus rupestribus *Caulibus* uncialibus, biuncialibus subtilitariis, *Foliis* minoribus angustioribus cinereis. *Corollis* minoribus, luteis, mox albescentibus. *Hæc* variatio pusilla, sine dubio apud *Linnaeum* sub *Alyssso* minimo allegata est.

982. **ALYSSUM incanum** caule erecto, foliis lanceolatis incanis integerrimis, floribus corymbosis petalis bifidis. *Linn. Sp. pl. p. 590. — Pollich. fl. pal. n. 616.*

Moenchia incana. Roth. fl. germ. 2. p. 73.

Thlaspi capsulis oblongis, incanum. Bauh. hist. 2. p. 929,

iberis prima. Tabernæmont. p. 848.

Germ. Weisblühendes Steinkraut.

Gall. Alysse blanchatre.

Hab. In arenosis sterilibus ad frutices *am Landgraben* inter *Carlsruhe* et *Mühlburg* infrequens; prope *Graben*; circa *Mannheim* ad *Rheni* ripam abunde; ad vias arenosas inter *Ladenburg* et *Lorsch*; nec non inter *Oppenheim* et *Worms* post *Pollichium* vidi.

Flor. Junio, Julio, Augusto. *Semina* Septembris matura. ♂

Radix fusiformis, fibrosa, albida. *Caulis* sesquipedalis, bipedalis et altior, erectus, obsolete angulatus, villosus-incanus, scaber, ramulosus. *Rami* alterni, sparsi, erecti. *Folia* al-

40 TETRADYN. SILICULOSA.

terna, sessilia, lanceolata, integriuscula, villosa, incana. *Flores* caulem ramosque terminantes, racemosi, quasi corymbum longum densum imitantes. *Corollæ Petala* calyce triplo fere longiora, femibifida, alba. *Siliculæ* ovatae, acutiusculæ, compressæ, glabriusculæ, *Stylo* longo instructæ, bivalves, 7-8 spermæ.

Flores apibus mel præbent.

Planta tenella ovibus accepta et conveniens. *Ad arenam* in sterilibus figendam apta.

CCCXI. COCHLEARIA.

Linn. Gen. pl. ed. Schreb. n. 1079.

Calyx tetraphyllus: *foliolis* ovatis, concavis, dehiscentibus, deciduis. *Corolla* tetrapetala, cruciformis. *Petala* obovata, patentia, calyce duplo majora, unguibus angustis calyce brevioribus, patulis. *Silicula* cordata, gibbifolia, turgida, *Stylo* instructa, bilocularis; *valvulis* gibbis obtusis. *Semina* quatuor circiter in singulo loculo.

983. *COCHLEARIA officinalis* foliis radicalibus cordato-subrotundis, caulinis oblongis subsinuatis. *Linn. Sp. pl. 2. p. 903.* — *Hall. stirp. helv. n. 503. sub Nasturtio.*

Cochlearia batava. *Blackw. herb. t. 227.*

Cochlearia folio subrotundo. *Bauh. pin. 110.*

Cochlearia. *Tabernæmont. p. 847. Icon.*

Icon. Oeder. fl. Dan. t. 135. — Regnault. bot.

Germ. Aechtes Löffelkraut.

Gall. Cochleaire. l'herbe aux cuilliers.

Hab. In Europæ borealis littoribus marinis. In Ditione Basileensi prope *Pierre pertuis* in rupibus irrigatis t. *Zeyhero*. *Colitur* passim in hortis. Circa *Carlsruhe* ad viam *über der Schiefswiese* ante aliquot annos eam vidi, sine dubio ex hortorum reiectamentis.

Flor. Majo, Junio. *Semina* Julio Augusto matura. ☉ ♂.

Caules ex radice plures, semipedales, pedales, erecti, angulati, glabri, simplices. *Folia* alterna, radicalia longe petiolata, cordato-orbiculata, integriuscula vel obsolete sinuato-angulata; *caulina* ovato-oblonga, dentata, obtusa, superiora sessilia, amplexicaulia, omnia glabra, laete viridia, succulenta. *Flores* cauliterminantes, racemoso-corymbosi. *Corollæ* *Petala* ovata, obtusa, unguiculata, patentia, calyce duplo longiora, alba. *Siliculæ* ovatæ, inflatæ, rugosulæ, *Stylo* brevi instructæ, bilocularibus, 8-10 spermæ.

Pharm. COCHLEARIA *Herba recens.*

Herba masticata, amariuscula, acris, antiscorbutica, incidens, mundificans, usitatissima.

Herbam cum lacte acidulato vel ejus sero parant Islandi; condiunt eam etiam sale culinari in magnis doctis et per hiemem servant. Oves avidè quidem illam depascunt valde pinguescunt, sed caro nauseoso inde sapore inficitur. Eggert Olaffen. *Reise in dieuene Island*. I. p. 483.

984 COCHLEARIA *Armoracia* foliis radicalibus lanceolatis crenatis, caulinis incisis. *Linn. Sp. pl.* 2. p. 904. — *Hall. stirp. helv.* n. 504. *sub Nasturtio.*

Raphanus ruscifolius. *Bauh. pin.* 96.

Raphanus maritimus. *Tabern. p.* 799. *Icon.*

*Icon. Blackw. t. 415. — Regnault. bot. —
Germ. Meerrettig.*

Gall. le Raifort sauvage.

Hab. In fossis et ad rivulos Europae passim.
Circa *Carlsruhe* ad fossam versus *Durlach*.
forſan ex hortorum adjacentium rejecta-
mentis. Colitur in oleraceis.

Flor. Junio, Julio. *Semina* Augusto, Sep-
tembri maturat. 7

Radix cylindrica, craſſa, ramosa, alba,
firma carnoſa. *Caulis* cubitalis, tripedalis et
altior, erectus, glaber ramosus. *Folia radicalia*
petiolata, ovato-lanceolata, quandoque corda-
to-ovata, crenata, *caulina*, alterna ſubſeffilia,
lanceolata, incifo dentata, aut profundius in-
ciſa, omniaglabra, ruſoſo-venoſa. *Flores* ter-
minales, racemoſi, laxi. *Corollae Petala* calyce
duplo longiora, alba. *Siliculae* ovatae, ſubun-
dulatae, bivalves, 8-10 ſpermæ.

O b ſ. Individua ſpontanea in Gallia meridionali obſer-
vata. Ceſſiſſime foliis radicalibus cordato-ovatis.



CCCXII. CORONOPUS.

Hall. stirp. helv. p. 217.

Calyx tetraphyllus, *Corolla* tetrapetala, cruciformis: *Petalis* ovatis obtusis. *Stylus* brevis, capitatus, deciduus. *Silicula* rotundato-re-niformis, tuberculis atque lamellis radiantibus versus peripheriam muricata et in ambitu quasi cristata, in medio glabriuscula, compressa, bilocularis, evalvis. *Loculamenta* undique clausa, et ad axem inter se connata ut pericarpium ex duabus siliculis partialibus compositum esse videtur. *Semina* solitaria, ovata, lenticulari-compressa. *Gart. fruct pl. 2. p. 293.*

985. **CORONOPUS *Ruellii*** caule depresso, foliis pinnatifidis, filiculis rotundato-re-niformibus cristato-muricatis.

Coronopus Ruellii. *Hall. stirp. helv. n. 502.*

Cochlearia Coronopus foliis pinnatifidis.

Linn. Sp. pl. 2. p. 904.

Nasturtium sylvestre capsulis cristatis. *Tourn. inst. — Mapp. fl. Als. p. 209.*

Coronopus Ruellii sive *Nasturtium verrucosum.* *Bauh. hist. 2. p. 919.*

Ambrosia campestris repens. *Bauh. pin. 138.*

Coronopus repens. *Tabernaemont p. 302. Icon. bona.*

Icon. Oeder. fl. Dan. t. 202. — Blackw. t. 120.

Schkuhr. bot. Handb. t. 181.

Germ. Ruellischer Kraehensufs.

Gall. Cranson de Mr. Ruellius.

Hab. Circa Carlsruhe auf der Schiefswiese in-frequens. Prope Graben ad vias apricas humidiusculas. In Margraviatus superioris

subudis lutosis inter *Mullheim* et *Neuenburg*, t. *Vulpio* rarius. Circa *Basileam am Spahlen Thor auf dem Davids - Boden*, versu *Mænchenstein*, inter pontem *Birlæ* et *Mutenz*; inter *Dornach* et *Arlesheim* in viis lulosis. In Principatu superiori prope *Petershausen* et *Constanz* ad vias abundanter videtur. In *Alsatia* prope *Argentina* *vor dem Spital* et *Kronenburgerthor* ac alibi passim. In *Palatinatu* ad vias inter *Lambsheim* et *Frankenthal*; inter *Alzey* et *Alwig* post *Pölichium* vidi.

Flor. Junio, Julio, Augusto. *Semina* Augusto Septembri matura. ☉

Radix fibrosa albida. *Caules* ex radice plures, palmares, pedales, ad terram depressi, teretes, ramosi. *Rami* decumbentes, alterni. *Folia* alterna, in petiolum brevem desinentia pinnatifida: *Pinnis* alternis, cuneiformi-lanceolatis, subintegris, ex glauco-viridibus, glabris. *Flores* racemosi, laterales. oppositifolii sessiles, breves, pauciflori. *Corollæ* parvæ, albidæ. *Siliculæ* subcordatæ, cristatæ, bispermae.

Pharm. NASTURTII VERRUCOSI Herbar. recens.

Planta acris nasturcina ob *Johannam Stephens* famosa, quae suo ad calculum medicamentum usum admiscuit. Hall. l. c. in Pharmaciis non confundenda cum *Plantagine Coronopo* Linn.

Planta tenella inter *Acetaria* praesertim in Angliā ahibetur.

986. *CORONOPUS didyma* caule procumbente piloso, foliis pinnatifidis, floribus subdiandris, siliculis didymis reticulato-rugosis, emarginatis.

TETRADYN. SILICULOSA. 45

Coronopus didyma filiculis emarginatis didymis reticulato-rugosis, stylo obsoleto, corymbis multifloris. *Smith Comp. fl. Britann. p. 208.*

Lepidium didymum. *Linn. Mant. 92. Ait. Hort. Kew. 2. p. 374.*

Lepidium anglicum floribus diandris tetrapetalis, foliis pinnatifidis, caule procumbente piloso. *Hudson. fl. Angl. p. 280.*

Germ. Englischer Krähenfuß.

Gall. Cranfon d'Angeltaire.

Hab. In Angliae ruderatis. Circa *Carlsruhe* in ruderatis *am Holzmagazin* quasi spontanea sine dubio ex horto Botanico emigravit.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caules ex radice plures, pedales et longiores procumbentes, teretes, glaces, pilosi, ramosi. *Rami* alterni, procumbentes. *Folia* alterna, profunde pinnatifida: *Pinnis* alternis lanceolatis acutiusculis, glabriusculis, subintegris. *Flores* racemosi, laterales, oppositifolii. *Corollæ* minutissimæ, albido-viridulæ. *Stamina* sæpe tantum duo conspicua. *Siliculæ* didymæ, cordatæ, emarginatæ, reticulato-rugolæ, dispersæ.

Planta pulchella, habitus Lepidii, Silicula Coronopi.

CCCXIII. BISCUTELLA.

Linn. Gen. plant. ed. Schreb. n. 1084.

Calyx tetraphyllus: *Foliolis* ovatis acuminatis, basi gibbis, coloratis, deciduis. *Corollæ*

tetrapetala, cruciformis. *Petala* oblonga, obtusa, patentia. *Silicula* erecta, compresso plana; semibifida, lobis subrotundis, bilocularis: *dissepimento* lanceolato, in *Stylum* rigidum terminato; loculis bivalvibus, *dissepimento* affixis. *Semina* solitaria, subrotunda, compressa in medio loculi posita.

987. *BISCUTELLA laevigata* filiculis glabris, foliis, lanceolatis, ferratis. *Linn Mant.* 255.

Biscutella *silicula* laevi, foliis asperis dentatis. *Hall- stirp. helv. n.* 501.

Biscutella didyma. *Scopol. fl. Carn.* 2. p. 14. an?

Icon. Jacq. fl. Austr. t. 339.

Germ. Glatte Brillenschote.

Gall. Lunetier lisse.

Hab. In Margraviatu superiore in collibus vulcanicis rupestribus praeruptis apricis graminosis *am Kaiserstuhl* Brisgoviae inter *Sasbach* et *Limburg* rarius.

Flor. Aprili, Majo. *Semina* Junio, Julio maturat. ☞

Radix praelonga, teretiuscula, subramosa, *Caules* ex radice saepe plures, sesquipedales et altiores, erecti, teretes, rariter hirsuti, superne ramosi. *Folia* inferiora oblonga, obtusiuscula, basi attenuata, in petiolum desinentia. remote dentato-ferrata, caulina superiora alterna, sessilia, integriuscula, minora, omnia pilosa, hirsuta, canescentia, *Flores* caulem ramosque terminantes, corymboso umbellati. *Calycis* foliola flavescentia, obtusa, patula, decidua, glabra. *Corollae* petala oblonga, ob-

tas, patentia, lutea, calyce duplo longiora. *Silicula* didymæ, planæ, utrinque glabræ, *Stylo* prælongo terminatæ.

Obs. Planta in alpinis Helvetiæ et in Pyrenæis ubi abundanter crescit observata, minor, pilis albis longioribus instructa est. In Horto ex seminibus culta, æquino major, perennis.

Ad rariores Floræ nostræ cives pertinet.

CCCXIV. L U N A R I A.

Linn. Gen. pl. ed Schreb. n. 1085.

Calyx tetraphyllus: *foliis* ovato-oblongis, obtusis, conniventibus, deciduis, quorum duo alterna basi gibba. *Corolla* tetrapetala, cruciiformis. *Petala* integra, plana, obtusa, magna, in ungues longitudine calycis desinentia. *Stylus* brevis. *Stigma* obtusum, integrum. *Silicula* elliptica, compresso-plana, integra, marginata, pedicellata, stylo terminata, bilocularis, bivalvis, *Dissepimento* valvulis parallelo, æquali, plano. *Semina* nonnulla, reniformia, compressa, marginata, in medio filiculæ posita: *Receptaculis* filiformibus, longis, futuris lateralibus insertis.

988. *LUNARIA biennis* radice bienni foliis cordatis dēntatis filiculis ellipticis obtusis.

Lunaria annua foliis oppositis. *Linn. Sp. pl. 2. p. 911.*

Lunaria major filiqua rotundiore. *Bauh. pin. 203. — Mapp. fl. Alf. p. 181.*

*Viola lunaris prima. Tabernæmont. p. 696
Icon.*

Icon. Schkuhr. bot. Handb. t. 182.

Germ. Zweyjährige Mond-Viole. Mondkraut.

Gall. Lunaire bisannuelle.

Hab. In adscensu montis Thurnberg retro Durlach in dumetis ad vinearum vias in frequens. In Alfatia ad vallem retro Teutsch-Rumbach; retro Haslach in den Niedecker Thal t. Mappo. Colitur passim in hortis.

Flor. Majo, Junio. Semina Julio, Augusto maturat. ♂

Radix biennis. Caulis 2-3 pedalis, erectus, striatus, pilosus, ramosus. Folia inferiora opposita, superiora alterna, petiolata, cordata, dentata, hirsuta. Flores caulem ramisque terminantes racemosi. Racemi laxi. Corollæ ex albedo-violaceæ, inodoræ. Siliculæ ellipticæ, utrinque obtusæ.

Nota. Facile a sequente cum qua sæpe confunditur Siliculis ellipticis subrotundis, nec elliptico-lanceolatis distinguitur.

Obs. In Horto culta, mutationi non succubuit.

Flores apibus mel largiuntur.

989. LUNARIA *perennis* radice perenni foliis profunde cordatis, inæqualiter sub sinuato-dentatis, siliculis elliptico-lanceolatis acutis.

Lunaria rediviva foliis alternis *Linn. Sp. pl. 2. p. 911,*

Lunaria perennis filiquis ovatis lanceolatis *Hall. stirp. helv. n. 500.*

TETRADYN. SILICULOSA. 49

Lunaria major filiqua longiore *Bauh. hist.*
2. p. 882. — *Mapp. fl. Alf.* p. 181.

Viola Lunaria secunda. *Tabernæmont.* p.
696. *Icon.*

Germ. Wohlriechendes Mondkraut. Wohl-
riechende Mondviole.

Gall. Lunaire odorante.

Hab. In Principatu Fürstenbergenſi in rupi-
bus calcareis inter frutices ad rivulos et
ivos paſſim frequens cum Aconito Na-
pello. In Alſatia retro Markirch in mon-
toſis. t. Mappo. In Sponhemia prope Kirch-
berg in lapidoſis dumetoliſ, umbroſiſ hu-
midis non inſrequens. Colitur paſſim in
hortiſ.

Flor. Majo, Junio. Semina Julio Auguſto
matura. ☞

Planta ſpecioſa. Radix perennis. Flores
ſarveolentes. Corollæ albedo-violaceæ. Si-
licule ovali lanceolatae, utrinque attenuatae.

Obſ. Mutationi in Horto culta, non ſuccubuit.

Pharm. VIOLÆ LUNARIÆ Semina.

Semina amariuſcula acria olim in Epilepſia com-
mendabantur.

Flores apibus mel concedunt.

TETRADYN. SILIQUOSA.
CCCXV. DENTARIA.

Linn. Gen. pl. ed. Schreb. n. 1087.

Calyx tetraphyllus: foliolis ovato oblon-
gis, longitudinaliter conniventibus, obtuſiſ, de-
ciduiſ. Corolla tetrapetala cruciformiſ. Pe-
tala ſubrotunda, obtuſa, vix emarginata,
piana, deſinentia in ungues longitudine ca-
lyciſ. Siliqua longa, teres, biloculariſ, bi-

50 TETRADYN. SILIQUOSA.

valvis, elastice diffiliens, valvulis revolutis
dissepimento valvis paulo longiore. *Semina*
 plura subovata.

990. *DENTARIA bulbifera* foliis inferioribus pinnatis, summis simplicibus, bulbis in alis foliorum inferioribus.

Dentaria bulbifera foliis inferioribus pinnatis, summis simplicibus. *Linn. Sp. pl.* 2912. — *Hall. stirp. helv.* n. 470.

Dentaria heptaphyllos baccifera. *Bauh. pin.* 322.

Dentaria quarta. *Tabernæmont. p.* 324 *Icon.*

Icon. Oeder. fl. Dan. t. 361. — *Schkuhr. bot. Handb. t.* 183.

Germ. Bollentragende Zahnwurz. Korallenwurz.

Gall. Dentaire bulbifere.

Hab. In Sponhemix sylvis montosis apricis prope *Winningen* ad Mosellam non infrequens.

Flor. Majo. *Semina* Julio matura. 7

Radix corallina, dentato-squamosa, alba. *Caulis* sesquipedalis et altior, erectus, teretis, glaber, simplex. *Folia* inferiora opposita, petiolata, pinnata, foliolis septem, lanceolatis serratis, scabriusculis, caulina media quinque superiora subterna, suprema alternà, simplicibus. *Bulbilli* in axilla foliorum præsertim inferiorum sessiles, subrotundi, maturitate atro-fusci, decidui. *Flores* caulem terminantes, corymbosæ racemosæ. *Corollæ* ex albido dilute purpureæ violaceæ, speciosæ. *Semina* sæpissime abortientia.

Pharm. DENTARIÆ f. ANTIDYSENTERICÆ Radix.

Radix acris, saporis ingrati olim contra infantum tormina et Dysenteriam laudata. Non confundenda cum Radice DENTELLARIÆ quæ ex Plumbagine europaea Linn. in Gallia meridionali et Hispania obvia desumenda est, quæ alba, longa, crassiuscula, carnosa, acris et fervida atque salivam elicit, hanc ob rationem in dentium doloribus præsertim usitata, a qua virtute sine dubio nomen obtinuit.

Nota. Bulbuli Dentariæ, terræ commissi prolem propagant.

991. *DENTARIA heptaphyllos* foliis inferioribus septenis, superioribus quinis. *Hall. stirp. helv. n. 469. β*

Dentaria foliis omnibus pinnatis, foliolis quinis septenisque oblongis, ferratis. *Lamarck. Encycl. 2. p. 268. — Aiton. hort. Kew. 2. p. 386.*

Dentaria pentaphylla foliis summis digitatis, *Linn. Sp. pl. 2. p. 912.*

Dentaria heptaphyllos. *Bauh. pin. 322. — Mapp. fl. Als. p. 93.*

Dentaria secunda. *Tabernæmont. p. 324. Icon.*

Germ. Siebenblättrige Zahnwurz.

Gall. Dentaire à sept folioles.

Hab. In Margraviatu superiore in sylvis montosis prope *Candern* non infrequens. In Ditionis Basileensis umbrosis sylvaticis montosis montis *Mutet*, in monte *Wasserfall*, *Vogelberg*, *Kallen*, *Belgen* et alibi frequens. In Alsatia sylva *der Dambacher Wald*, in montosis prope *Wangenburg t. Mappo.*

Flor. Aprili, Majo. *Semina*, Junio, Julio maturatione. ¶

Radix corallina, alba. *Caulis* cubitalis et altior, erectus, simplex. *Folia* omnia pinnata caulina inferiora petiolata, foliolis septem, superiora foliolis quinque, ovali-lanceolatis serratis, glabris. *Flores* caulem terminantes corymbofo-racemosi. *Corollæ* albæ, violaceæ mixtæ, speciosæ.

Obs. Planta in horto culta, mutationi non succubuit
Flores apibus accepti.

992. *DENTARIA pentaphyllos* foliis omnibus quinato digitatis. *Gerard. fl. Gallopr.* p. 356.

Dentaria digitata foliis digitatis serratis
Lamarck. Encycl. 2. p. 268.

Dentaria pentaphyllos β. γ. *Linn. Sp. pl.* 2 p. 912.

Dentaria foliis quinatis digitatis. *Hall. stirp. helv.* n. 469. β

Dentaria pentaphyllos foliis mollioribus
Bauh. pin. 322. — *Mapp. fl. Als.* p. 92

Dentaria tertia. *Tabernaemont. p.* 324. *Icon bona.*

Germ. Zahnwurz mit gefingerten Blättern.

Gall. Dentaire digitée.

Hab. In Ditione Basileensi in monte Rechaas versus Wallenburg rarius t. Zeyhero. In Alsatia montosis retro Baar et Markirch in sylvis umbrosis; in montosis prope Stein ad Bergzabern t. Mappo.

Flor. Aprili, Majo. *Semina* Junio, Julio maturatione. ¶

Facile a præcedente affini specie distinguitur, *Foliis* omnibus digitatis quinatis.

Obi. Planta in horto culta constantissima permanfit, hinc omni jure pro peculiari specie declaranda.

Ex hac etiam atque ex præcedente specie Radix *DITTARIÆ* officinarum promiscue defumitur.

CCCXVI. C A R D A M I N E.

Linn. Gen. plant. ed. Schreb. n. 1088.

Calyx tetraphyllus: *foliis* ovato-oblongis, obtusis, subpatulis, gibbis, parvis, deciduis. *Corolla* tetrapetala, cruciformis. *Petala* oblonga, obovata, patentissima, desinentia in ungues erectos, calyce duplo longiores. *Siliquæ* longa, cylindraceo compressa, bilocularis, bivalvis; *valvulis* dehiscendo spiraliter revolutis. *Semina* plurima, subrotunda.

995. *CARDAMINE impatiens* foliis pinnatis incisis stipulatis, floribus apetalis. *Linn. Sp. pl. 2. p. 914. — Pollich. fl. pal. n. 619.*

Cardamine foliis pinnatis, imis obtuse lobatis, caulinis acute dentatis, floribus apetalis. *Hall. stirp. helv. n. 471.*

Cardamine annua exiguo flore? *Mapp. fl. Alf. p. 53.*

Sisymbrii Cardamines species quædam insipida. *Bauh. hist. 2. p. 886.*

Icon. Oeder. fl. Dan. t. 735.

Germ. Flüchtigblühende Cardamine. Spring-Schaumkresse.

Gall. Cardamine apétale.

Hab. In montosis umbrosis sylvis roridis, præfertim cum Chrysofplenis passim non infrequens.

Flor. Aprili, Majo, *Semina.* Junio, maturat. ♂

Caulis pedalis et altior, erectus, ramosus; vel simpliciusculus, acutangulus, glabriusculus, *Folia radicalia* pinnata: *pinnis* petiolatis, ovatis, obtusis, integriusculis, femi bi- aut trilobis, glabris; *caulina* alterna, pinnata, pinnulis acute incisis, integriusculisve, ad basin utrinque *Stipula* lineari, pilosa, aucta. *Flores* caulem ramosque terminantes, racemosi. *Racemi* laxi. *Calycis* foliola concava, obtusa, ex viridi-fuscescentia. *Corollæ* petala sæpissime nulla, quandoque quatuor, parva, calyce longiora, fugacia. *Stamina* filamenta sex, quatuor longiora, duo subbreviora. *Siliquæ* tenues, tumidiusculæ, longæ, glabræ, maturæ cum impetu elastice diffilientes. *Semina* subrotunda.

Obs. Variat quandoque præsertim in locis minus umbrosis roridis *Staminibus* quatuor, idem observavit Cel. Schkuhr. l. c. p. 268.

994. *CARDAMINE hirsuta* caule foliisque hirsutis pinnatis, floribus tetrandris, quandoque hexandris.

Cardamine hirsuta foliis pinnatis, floribus tetrandris. *Linn. Sp. pl.* 2. p. 915. — *Pollich. fl. pal.* n. 620.

Cardamine foliis pinnatis hirsutis, pinnis subrotundis, staminibus quaternis. *Hall. stirp. helv.* n. 472.

Cardamine quarta *Dalechampii.* *Mapp. fl. Als.* p. 53.

Sisymbrium aquaticum alterum. Camerar. epit. 270. Icon bona.

β. *Glabra, foliis brevissimis pilis fere inconspicuis hirta, flaminibus quatuor quandoque sex.*

Cardamine. Hall. stirp. helv. n. 472. var. β

Cardamine pratensis parvo flore. Mapp. fl.

Alf. p. 53. an? et Cardamine pratensis flore minore albo. Mapp. l. c.

Cardamine sylvestris minor italica. Barrel. Icon. 455.

Nasturtium aquaticum minus. C. Bauh. Cat. pl. Basil. p. 30.

Icon. α. Curtis. fl. Lond. t. 277. — Schkuhr. bot. Handb. t. 187. — Oeder. fl. Dan. t. 148.

Germ. Rauchhaarige Cardamine.

Gall. Cardamine velue.

Hab. Utrunque in umbrosis sylvis humidis, alnetis, populetis, salicetis et pratis udis cum variatione β. passim frequens, nec sicciora tamen respuit.

Flor. Aprili, Majo. Semina Majo, Junio matura. ☉ ♂

Radix fibrosa, tenuis. Caulis palmaris, pedalis, erectus, angulosus, sulcatus, subleuosis, inferne praesertim hirsutus, superne saepe glabriusculus, quandoque totus glabriusculus, simpliciusculus, saepe ramosus. Rami alterni, patentiusculi, glabriusculi. Folia pin-nata, radicalia caulinaque infima foliola petio-lata opposita, alternaque, subrotunda, aut ob-longa, obtusa, subdenticulata: impari majori, obtusiori, subtrilobo; caulina superiora foliola sessilia, lineari-lanceolata, integriuscula, om-nia utrinque hirsuta, quandoque glabriuscula Flores caulem ramosque terminantes, corym-

bofo-racemofi. *Calyx* tetraphyllus, patulus, *foliolis* oblongis, obtusis, villosiusculis, margine rubellis, *Corolla* tetrapetala, *petalis* calyce duplo longioribus, obverse lanceolatis, obtusis, albis. *Stamina* filamenta sæpe quatuor, æqualia, *Antheris* flavis instructa, rarius quinque, quandoque sex, tetradynama. *Siliquæ* erectæ, lineares, compressæ, glabræ, unciam circiter, longæ, elasticæ. *Semina* ovata, compressa.

Obf. Planta pro diverso loco umbroso, aquoso, humido, demissiore, aut elatiore montoso sicciore differt majori et minori hirsutie, quandoque fere glabra occurrit. Specimina glabra staminibus sæpe sex instructa. b. Linnaeus forsan sub *Cardamine* parviflora comprehendebat.

Individua staminibus sex tetradynamis instructa, ab illis, staminibus saltem quatuor prædita sub pecuriari specie *Cardamines* sylvaticæ descripsit Cel. Link. in phytogr. Blættern. 1803. p. 50.

Stamina nobis sæpe quaterna in aliis sena, in nonnullis, rarius quina. Idem observaverunt Cel. Schkuhr. et Pollichius. Intra unum annum individua vidit Schkuhrus, quorum omnes fere

TETRADYN. SILIQUOSA. 57

1. *Cardamine pratensis* magno flore pleno.
Mapp. l. c.

Icon. n. *Curtis fl. Lond. t. 175. Regnault bot.*
— *Oeder. fl. Dan. t. 1039.*

Germ. *Wiesen-Cardamine. Wiesen-Kresse.*

Gall. *Le Cresson des prés. Cardamine des prés.*

Hel. Utrinque in pratis humidis, udis, ad rivulos passim frequens.

Flor. Martio, Aprili, Majo. Semina Majo, Junio matura. 7.

Radix fibrosa, albida. *Caules* ex radice sæpe plures, pedales, sesquipedales, erecti, teretes, glabri, simpliciusculi. *Folia* pinnata: *radicalia* majora, *foliis* subpedicellatis, alternis oppositisve, ovali-subrotundis, dentatis, *caulina* alterna: *foliis* lineari-lanceolatis, integrisculis, omnibus glabris. *Flores* terminatis, racemoso-corymboli. *Calycina* foliola quatuor, ovalia, obtusa, erecta, margine membranacea, albida, quorum duo opposita basifixa. *Corollæ* speciosæ. *Petala* quatuor, obovata, obtusa, albida aut dilute ex purpureo-violacea, venis reticulatis violaceo-cæruleis picta. *Siliquæ* lineares, subcompressæ, glabræ.

Pharm. *CARDAMINES* f. *NASTURTI PRATENSIS* Herba recens; Flores.

Planta amaricans, acriuscula, antiscorbutica, vires exercet antispasmodicas. In Epiphia, Chorea St. Viti, aliisque morbis convulsivis, sæpe magno cum fructu adhibetur.

Flores apibus mel largiuntur.

996. *CARDAMINE amara* foliis radicalibus caulinisque pinnatis conformibus,

foliolis ovalibus obtusis dentatis, caulibus stolonifero.

Cardamine amara foliis pinnatis, axillis stoloniferis. *Linn. Sp. pl. 2. p. 915.* — *Pollich. fl. pal. n. 622.*

Cardamine foliis pinnatis subrotundis angustatis. *Hall. stirp. helv. n. 474.*

Cardamine flore majore elatior. *Tournefort. inf. Mapp. fl. Als. p. 53.*

Nasturtium aquaticum majus et amarum. *Bauh. prodr. p. 45. Icon.*

Icon. Curtis fl. Lond. t. 158.

Germ. Bittere Cardamine.

Gall. Cardamine amère.

Hab. Utrunque in sylvis udis aquosis, praesertim alnetis et salicetis passim frequenter. Circa *Carlsruhe* in sylva *Hauaker*; inter *Rintheim* et *Durlach* abunde.

Flor. Aprili, Majo. *Semina* Junio matura.

Distinguitur a praecedente specie, *Radice* repente stolonifera, *Caulibus* saepe altioribus crassioribus, magis viridibus, succulentioribus. *Foliis* omnibus tam radicalibus quam caulibus pinnatis: *foliolis* conformibus, omnibus ovalibus, obtusis, dentatis, glabris, laete viridibus succosioribus. *Corollis* majoribus lacteis, ventris aquosis pictis, *Antheris* violaceis.

Nota. Stolones axillares quibus b. *Linnaeus* speciem distinxit, in individuis nostratibus desunt. Ex radice et parte infima caulis omnino Stolones proveniunt.

Pharm. CARDAMINES AMARÆ s. NASTURTII MAJORIS AMARI Herba recens.

Planta egregia antiscorbutica, convenit cum *Silybri* *Nasturtio*, sed in acetariis magis amara est. Flores apibus mel praebent.

CCCXVII. SISYMBRIUM.

Linn. Gen. pl. ed. Schreb. n. 1089.

Calyx tetraphyllus: *foliolis* lanceolato-linearibus, patentibus, coloratis, deciduis. *Corolla* tetrapetala, cruciformis. *Petala* oblonga, *patentia*, calyce sæpe minora, quandoque majora, unguibus minimis. *Germen* oblongum. *Stylus* vix ullus. *Stigma* obtusum. *Siliqua* ~~sub~~incurva, gibba, teres, bilocularis, bivalvis, ~~extralis~~ dehiscendo rectiusculis: *dissepimento* valvis paulo longiore. *Semina* plurima parva.

† *Siliquis* *declinatis* *brevibus*.

997. **SISYMBRIUM** *Nasturtium* filiquis *declinatis*, foliis pinnatis: foliolis subcordatis. *Linn. Sp. pl. p. 916.* — *Pollich. fl. pal. n. 623.*

Sisymbrium foliis pinnatis, pinnis subrotundis, brevibus racemis. *Hall. stirp. helv. n. 482.*

Sisymbrium aquaticum. *Mapp. fl. Alsp. 187.*

Sisymbrium aquaticum primum. *Tabernæmont. p. 846. Icon.*

Icon. Curtis. fl. Lond. — Bulliard. Herb. t. 302.

Oeder. fl. Dan. t. 690. — Blackw. t. 260.

Germ. Brunn-Kresse. Wasser-Kresse.

Gall. Cresson d'eau. Cresson de fontaine.

Hab. Utrunque in rivulis, fossis aquaticis, ad scaturigines passim abunde.

Flor. Majo, Junio, Julio. *Semina* Julio, Augusto matura. ♀

Radix fibrosa, repens. *Caulis* pedalis et longior, radicans, decumbens, sulcatus,

glaber, ramosus. *Rami* alterni. *Folia* alterni impari-pinnata: *foliola* sessilia, opposita, ovata, obtusa, impari majore, subcordato subrotundo, glabra, læte viridia, succulenta. *Flora* caulem ramosque terminantes, racemosi. *Calyx* foliola corolla breviora, margine alba. *Corollæ* petala integra, obtusa, alba. *Antheræ* flavæ. *Siliquæ* breves, gibbulæ, compressiculæ.

Variat Caulibus plus minusve decumbentibus quandoque adscendentibus, erectiusculis.

Pharm. NASTURTII AQUATICI Herbar. recens.

Planta egregia antiscorbutica, masticata acris amariscula.

Ad acetaria cum saccharo mixta, accepta, præsertim tempore colligitur. Cavendum ne confundatur cum Cardamine amara quæ caule erecto altiore, foliis multo majoribus, magis ablongis, corollis multo majoribus, sapore porro magis amaro dignoscitur.

998. *SISYMBRIUM sylvestre* filiquis declinatis, foliis pinnatis: foliolis lanceolatis ferratis. *Linn. Sp. pl.* 2. p. 916. *Pollich. fl. pal.* n. 624.

Sisymbrium foliis pinnatis, pinnis dentatis. *Hall. stirb. helv.* n. 485.

Brachiolobus sylvestris. *Allion. fl. Pedem.* t. 56. f. 2.

Sisymbrium palustre repens, Nasturtii foli. *Mapp. fl. Als.* p. 289.

Eruca palustris Nasturtii folio filiqua oblonga. *Bauh. pin.* 98.

Eruca palustris minor. Tabernamont. p. 836. *Icon bona.*

en. Curtis fl. Lond. t. 153.

*erm. Waldkresse. Wald - Rauke. Waf-
serfensf.*

ab. Utrunque ad fossas aquaticas, vias humi-
das, paludosas circa urbes et pagos. Fre-
quentissima in Rheni regionibus planta
agris sæpe molesta.

lor. Junio, Julio, Augusto. *Semina* Au-
gusto, Septembri matura. **z**

Radix quammaxime repens, tenuis, albida,
caules ex radice plures, pedales et altiores,
andoque demissiores, adscendentes, erectius-
si, debiles, angulato-firiat, glabriusculi,
moli. *Rami* alterni villulosi. *Folia* radi-
ca petiolata, profunde pinnatifida, glabra,
immo subovatis, dentato-ferratis, *caulina* al-
ma, subpinnata, pinnis lanceolatis ferratis,
tegrisve. *Flores* caulem ramosque terminan-
s, racemoso-corymboli. *Calycis* foliola ova-
ta, concava, flavescentia. *Corollæ* petala par-
ta, obtusa, patentia, lutea, calyce duplo fere
longiora. *Siliquæ* oblongæ, curvulæ, paten-
s, demum declinatæ, polyspermæ.

Nota. Ob nimiam luxuriam in agris inundationibus
Rheni expositis magnum sæpe vitium, difficiliter
diminuendum.

89. *SISYMBRIUM palustre* filiquis de-
clinatis oblongo-ovatis, foliis pinnati-
fidis, petalis calyce brevioribus. *Leyff.*
fl. hal. n. 679, — Poll. fl. pal. n. 625.

Radicula foliis pinnatis, pinnis dentatis,
petalis calyce brevioribus. *Hall. stirp.*
helv. n. 487.

Sisymbrium aquaticum, foliis in profundas
lacinias divisis, filiqua breviori. *Mapp.*
fl. Als. p. 289.

Raphanus aquaticus foliis in profundas lac-
nias divisilis. *Bauh. Prodr. p. 38. Icon.*

Icon. Oeder. fl. Dan. t. 931.

*Germ. Sumpf-Kresse. Sumpf-Rauke. Sumpf-
Senf.*

Gall. Sifymbre des marais.

Hab. Utrinque ad fossas, aquas, paludes pa-
sim copiose præsertim in Rheni vicini-
inundata.

Flor. Junio, Julio, Augusto. *Semina* Augusto
Septembri matura. ☉ ♂

Radix fusiformis, alba. *Caulis* pedali
bipedalis et altior, erectus aut decumbens, an-
gulosus, glaber, ramosus. *Rami* alterni. *Folli-*
alterna, petiolata, amplexicaulia, profunde
pinnatifida: segmentis alternis oppositisve, ova-
li-lanceolatis, obtusiusculis, ferratis, glabris
læte viridibus. *Flores* caulem ramosque ter-
minantes, racemosi. *Calycis* foliola, margini
lutescentia, corolla longiora. *Corollæ* petala
obtusa, flava, parva. *Siliquæ* ovato-cylindricæ,
breves, turgidæ, rugosulæ, polyspermae.

1000. *SISYMBRIUM amphibium* fili-
quis declinatis oblongo-ovatis, foliis
oblongo-lanceolatis pinnatifidisve ferra-
tis, calyce longioribus. *Linn. Sp. pl. ed.*
Willden. 3. p. 491.

Sisymbrium amphibium filiquis declinatis,
oblongo-ovatis, foliis pinnatifidis ferra-
tis. *Linn. Sp. pl. 2. p. 917. Pollich. fl.*
pal. n. 626. (sub *S. aquatico* petalis ca-
lyce longioribus.)

Radicula foliis integris aut pinnatifidis, pe-
talis calyce longioribus. *Hall. stirp. helv.*
n. 486

- α. *Sisymbrium palustre* foliis infimis in pin-
nas setaceas sectis, superioribus lanceo-
latis ferratis. *Pollich. l. c. β.*

Sisymbrium foliis infimis capillaceis, sum-
mis pinnatifidis. *Linn. Hort. Cliff. 337.*

- β. *Sisymbrium aquaticum* foliis simplicibus
dentatis ferratis. *Linn. l. c. β. Pollich. l. c.*

Radicula foliis oblongis ferratis integris.
Hall. stirp. helv. n. 436.

Sisymbrium aquaticum, Raphani folio, fili-
qua breviori. *Mapp. fl. Alf. p. 289.*

Raphanus aquaticus alter. *Bauh. Prodr.*
p. 38. Icon.

- γ. *Sisymbrium terrestre* γ. *Linn.*

Sisymbrium aquaticum foliis variis. *Vail-
lant. bot. parif. p. 185.*

Radicula foliis inferioribus ovatis ferratis, su-
perioribus semipinnatis. *Hall. stirp. helv.*
n. 486. β

Icon. Oeder. fl. Dan. t. 984.

Ger. Land- und Wasser-Rauke.

Gall. Sisymbre amphibe.

Hab. Utrique cum variationibus, in humi-
dis, udis, aquosis, fossis, inundatis præ-
sertim Rheno vicinis passim copiose.

Flor. Junio, Julio. *Semina* Augusto, Sep-
tembri matura. ☉ ☿

Facile a *S. palustri* distinguitur, *Radice* re-
pente fibrosa, *Corollarum* petalis calyce duplo
fere longioribus, *Siliquis* brevioribus, silicu-
lis magis referentibus.

Flores apibus mel largiuntur.

1001. *SISYMBRIUM pyrenaicum* sili-
quis subovatis, foliis inferioribus lyratis,

superioribus bipinnatifidis amplexicaulibus, stylis filiformibus. *Linn Sp. pl.* 3. p. 916. — *Vill. fl. Delph.* 3. p. 341. t. 38.

Sisymbrium foliis pinnatis imorum pinnis oblongis, superiorum linearibus, integris, siliquis ovalibus oligospermis. *La Chenal. Obs. bot. in Act. helv. Vol. IV.* p. 291.

Brachiolobus pyrenaicus. *Allion. fl. Pedem.* n. 1013. t. 18. f. 1.

Alyssum foliis imis subrotundis, caulini pinnatis filiformibus, amplexicaulibus. *Hall. stirp. helv.* n. 488.

Alyssum foliis pinnatis multiformibus, floribus racemosis luteis. *Allion. Spec. Pedem.* 40. t. 7.

Icon. De Lachenal. l. c. tab. 15. optima.

Germ. Pyrenäische Rauke.

Gall. Sisymbre des Pyrénées.

Hab. In Margraviatu superiore in pratis inter *Emmendingen* et *Freyburg*, nec non in pratis ad montem *Kaiserstuhl* prope *Ihringen*; prope Basileam in pratis inter *Neuhaus* et *Friedlingen* abunde; inter *Klüber* et *Kleinhuningen*, nec non versus *Weil*, et in pratis ad *Wiesam* post *Lachenalium* vidi.

Flor. Majo, Junio. *Semina* Julio matura. 7

Radix multiceps, obliqua, fibrosa, albida.

Caules ex una radice plures, pedales, sesquipedales et altiores, erecti, teretes, glabri vel subpubescentes, ramosi. *Rami* alterni subflexuosi, subdivisi. *Folia* glabra, vel rariter pilosula, *radicalia* profunde pinnatifida: *pinnulis* versus folii basin sensim decrefcentibus, oppositis,

lvs, alternisve rectangulis, ovali-oblongis, integrifusculis: foliolo extremo plerumque latiori; *caulina* superiora *ramaeque* amplexicaulia, pennata: *foliulis* linearibus angustis alternis oppositisve, longioribus, integris, vel una alteraque lacinia auctis, aliquantulum recurvata. *Flores* caulem ramosque terminantes, racemosi. *Calyx* tetraphyllus: *foliulis* ovali-linearibus, acutis, patentibus, ex viridi-luteis, glabris. *Corolla* tetrapetala, cruciformis, *petala* calyce tertia parte longioribus, patentibus, luteis. *Siliqua* ovato-oblonga, extrorsum gibbula, sesquilineam, ad duas lineas longa, glabra, *Stylo* tenui persistente praedita. *Semina* 6-8-12, quandoque plurima, subrotunda.

Floribus pulcherrime luteis, in pratis gregatim per magnos cespites se diffundentibus e longinquo dignoscitur. De La Chenal. l. c.

Obs. Individua in Pyrenæorum pratis abs me lecta, cæratibus simillima sunt.

Planta ex seminibus in Horto culta, mutatione non succubuit.

†† *Siliquis erectis longioribus.*

1002. *SISYMBRIUM tenuifolium* foliis infimis bipinnatifidis, superioribus pinnatifidis, supremis simplicibus, linearilanceolatis integris, glabris.

Sisymbrium tenuifolium foliis integerrimis: infimis tripinnatifidis; supremis integerrimis. Linn. *Sp. pl.* 2. p. 917. — Pollich. *fl. pal.* n. 627.

Eruca foliis glabris pinnatis, pinnis linearibus difformibus. Hall. *fl. alp. helv.* n. 461.

Brassica muralis foliis lanceolatis sinu-
ferratis læviusculis, caule erecto gracili
Huds. fl. angl. 1. p. 290.

Eruca sylvestris. *Blackw. herb. t. 266.*

Eruca tenuifolia perennis flore luteo. *Bo-
hist. 2. p. 861. — Mapp. fl. Als. p. 10*

*Icon. Curtis. fl. Lond. t. 158. (sub Brassica
murali) — Bulliard herb. t. 335. — F.
nault bot. t. 254.*

*Ger. m. Wilde stinkende Rauke. Dünneblättrige
Rauke.*

*Gall. La Roquette sauvage. Sisymbre b.
lante.*

*Hab. Inter Carlsruhe et Mühlberg am La-
graben in locis elevatis arenosis frequen-
ter. In Brisgovia ad muros et rupes prope
Breisach et alibi abunde, In Marggravia
superiore ad Wieslam t. C. Bauhino.
Palatinatu copiose ad vias et muros prope
Mannheim, Worms, Oppenheim, Mayentz
et alibi post Mappum vidi. In Alsatia prope
Argentoratum in muris nec non ad vineas
passim. In Sponhemia ad muros, in cul-
tibus et vineis passim copiose.*

*Flor. Junio, Julio, Augusto. Semina Augusto
Septembri matura. ♀*

Radix fusiformis, dura, extus flavica
Caulis bipedalis et altior, erectus, inferne
cumbens, teres, subsiriatus, glaber, ramosus
Rami alterni, copiosi. *Folia* alterna, petio-
lata, inferiora profunde pinnatifida: *pinnis*
neari-lanceolatis, simplicibus, divisisque,
tenuibus; *suprema* integra, lineari-lanceola
omnia ex glauco-viridia, glabra. *Flores* ca-
ulem ramosque terminantes, laxo racemo

Calycina foliola erecto-patentia, concava, haecolato-linearia. *Corolla* tetrapetala, cruciformis. *Petala* speciosa, calyce duplo longiora. obtusa, patentia, lutea. *Siliqua* erecta, uncialis et longior, compressa. *Stylo* brevi, tetragono, et *Stigmate* capitato, instructa.

Pasta digitis trita foetens, masticata ardens. acris.

Not. In regionibus sabulosis praecipitibus egregie arenam figit; ad Rhenum muros et rudera arcium antiqua passim condecorat,

Pasta in horto culta, constantissima permanfit.

Floues apibus mel praebent.

1003. **SISYMBRIUM** *Erucastrum* foliis runcinato-pinnatifidis, subhispidis, caule hispido. *Poll. fl. pal. n.* 628.

Sisymbrium *Erucasirum* subacaule, foliis ly-rato-runcinatis laevibus, caule hispido.

Gouan. Obs. bot. p. 42. *t.* 26.

Sisymbrium murale. *Villars. fl. Delph.* 3. *p.* 351.

Germ. Acher-Rauke.

Gall. Sisymbre des champs.

Hab. Utrinque in locis cultis Rhenum versus passim copiose. In agris inter *Lichtenau* et *Kehl* abunde. Circa *Ladenburg* ad aggeres gramineos et vias cum *Valantia cruciata*. In vinearum collibus siccis circa *Dürkheim*; in agris prope *dem Holzhof*, *Frankenthal*, *Worms*, *Oppenheim*, *Alzey*, *Lambsheim*, *Niederulm*, *Mainz* et alibi passim frequens post *Pollichium* vidi.

Flor. Majo, Junio, Julio. *Semina* Augusto matura. 2

Radix subfusiformis, fibrosa. *Caulis* digitalis. palmaris, pedalis, saepe sesquipedalis,

erectus, ad basin quandoque subdecumbens, teres inferne præsertim villoso-hirsutus, foliosus, rarius subaphyllus, ramosus. *Rami* alterni, villosiusculi. *Folia radicalia caulinaque* petiolata ad costam mediam fere pinnatifida, subruncinata, segmentis ovali-oblongis, obtusiusculis, subsinuato-dentatis, villosiusculis glabrisve, læte viridibus. *Flores* caulem ramisque terminantes, racemosi. *Racemi* simplices, longi, laxi. *Calyx* tetraphyllus, *foliolis* erectiusculis, pilosiusculis. *Corollæ* petala quatuor cruciformia, calyce duplo longiora, obovata, submarginata, pallide lutescentia, ætate, sæpe ex luteo-albida. *Siliquæ* oblongo-teretiusculæ, patulæ, glabræ, compressiusculæ, unciam circiter longæ, *Stylo* tereti, crassiusculo, et *Stigmate* subcapitato terminatæ.

Nota. Linnaeus sine dubio hanc speciem cum *S. murali* comprehendebat.

Individua nostra cum illo, quod Gouanus descriptum exhibuit, omni ex parte fere conveniunt. *Corollæ* tamen petala pallidiora in nostris observantur.

Non confundenda cum *Brassica Erucastro* Linn. quæ corollas habet fere duplo majores.

1004. *SISYMBRIUM arenosum* caule subfolioso ramoso, foliis lyratis rectangulo dentatis hispidis pilis ramosis *Linn. Sp. pl.* 2. p. 919.

Eruca foliis scabris semipinnatis pinnis rectangulis rarius dentatis. *Hall. stirp. helv.* n. 462.

Arabis arenosa, *Scop. fl. Carn. ed.* 2. p. 32. t. 40. — *Roth. fl. germ.* 3. p. 111.

Eruca coerulea in arenosis crescens. *Bauh. pin.* 99. — *Prodr.* p. 40. *Icon. Mapp. fl. Alf.* p. 101.

Icon. Barrelier. Icon. n. 196.

Germ. Sand-Rauke.

Gall. Sifymbre des sables.

Hab. In Margraviatu superiori in campestribus sabulosis prope Basileam, Rhenum versus. In Ditionis Basileensis arenosis ad Birsam frequens. In præaltis jugis et petris circa *Delsperg* et *Lauffen t. C. Bauhino*. In muris circa Castellum Befortienſe *t. Joh. Bauhino*; nec non in confiniis Sundgovia. In Sponhemia arvis prope *Wunningen* ad Mosellam passim non infrequens.

Flor. Aprili, Majo. *Semina* Junio matura. ☉

Radix fusiformis, fibrosa, fusca. **Caules** ex radice sæpe plures, palmares, pedales et altiores, erecti, teretes, siriati, hirsuti, ramosi. **Rami** alterni. **Folia** radicalia plurima, in orbem digesta, petiolata, ad costam mediam sere pinnatifida, lyrata: *pinnis* rectangulis ovali-oblongis subdentatis, suprema majori ovata dentata: *caulina* alterna, minora, minus profunde secta, omnia hirsuta: pilis ramosis, bifidis trifidisque instructa. **Flores** caulem ramosque terminantes, racemosi, laxiusculi. **Calycina** foliola glabra. **Corollæ** petala quatuor, obovata. calyce duplo longiora, pallide ex cæruleo violacea, venis saturatioribus picta, quandoque albida. **Siliquæ** tenues, subfiliformes, compressiusculæ, glabræ, *Stylo* lineari, obtuso, terminatæ, sesquiunciam ad binas uncias longæ.

Uſ. Individua pusilla, vix digitalia, incana. hirsuta, simpliciuscula, vidi prope Aranjuez in Castilia.

Flores apibus mel præbent.

1005. **SISYMBRIUM** *Loeselii* foliis run-
cinatis, acutis, hirtis, caule retrorsum
hispido. *Linn. Sp. pl.* 2. p. 921. — *Pol-
lich. fl. pal.* n. 630.

Erysimum foliis pinnatis, extima pinna trian-
gula, petiolis longis hirsutis. *Hall. Goett.*
p. 248.

Erysimum angustifolium majus. *Bauh. pin.*
107.

Erysimum hirsutum filiqua *Erucæ.* *Loesel. fl.*
pruss. p. 69. t. 14.

Icon. Jacqu. fl. Austr. t. 324.

Germ. Loeselifche Rauke.

Gall. Sifymbre de Mr. Loesel.

Hab. In Palatinatu transrhenano circa *Worms*
in aggeribus gramineis sub falicibus; inter
Oppenheim et *Guntersblum* ad vias post
Pollichium vidi.

Flo r. Junio, Julio. *Semina* Augusto matura. ☉

Radix fusiformi cylindrica fibrosa alba

1006. **SISYMBRIUM** *Sophia* petalis calyce minoribus, foliis decomposito pinnatis. *Linn. Sp. pl.* 2. p. 920. — *Pollich. fl. pal.* n. 629.

Sisymbrium pinnis et pinnulis foliorum pinnatis, laciniis linearibus subhirsutis. *Hall. stirp. helv.* n. 484.

Sisymbrium annuum Absynthii minoris folio. *Mapp. fl. Als.* p. 290.

Nasutium sylvestre tenuissime divisum. *Bauh. pin.* 105.

Sophia Chirurgorum. *Lobel. Icon.* 738.

Herba *Sophia* latifolia et angustifolia. *Tabernamont. p.* 26. *Icon.*

Icon. Bulliard. herb. t. 271. — *Blackw. t.* 440.

German. Sophienkraut. Wurmkraut.

Call. Sisymbre parviflore.

Hab. Utrinque in ruderatis, muris, arvis, ad viarum marginalia passim frequens.

Flor. a Majo in Augustum. *Semina* Augusto, Septembri matura. ☉

Radix fibrosa, albida. *Caulis* 2-3 pedalis et altior, erectus, teres, subpubescens, inermis, ramosus. *Rami* alterni. *Folia* alterna, elliptica, tripinnata: *pinnulis* linearibus, angustis, acutis. *Flores* caulem ramosque terminantes, racemoli. *Racemi* longi, multiflori. *Calycis* foliola erecta, linearia, obtusiuscula, margine lutescentia. *Corollae* petala angusta, obtusa lutea, parva, calyce subbreviora. *Siliuae* teretes, graciles, obsolete tetragonae, subcapitulosa, semiunciam longae, polyspermae. *Semina* ovata, obtusa, rufa, parva.

Pharm. SOPHIAE *Semina*; *Herba.*

Semina acria, stimulantia, hodie in medicina fore exoleta.

Flores apibus mel præbent.

Nota. SISYMBRIUM lrio foliis runcinatis dentatis nudis, caule lævi, filiquis erectis. Linn. et S. strictissimum foliis lasceolatis, (obsolète) dentato-serratis caulinis Linn. hucusque frustra in regionibus nostris quæsi.

CCCXVIII. E R Y S I M U M.

Linn. Gen. pl. ed. Schreb. n. 1090.

Calyx tetraphyllus: foliolis ovato-oblongis, parallelo-conniventibus, coloratis, deciduis. *Corolla* tetrapetala, cruciformis. *Petalum* oblonga, plana, apice obtusa, unguibus longitudine calycis, erectis. *Nectarium* glandula duplex intra filamentum brevius. *Germen* lineare, tetragonum, longitudine staminum. *Stylus* brevissimus. *Stigma* capitatum, persistens parvum. *Siliqua* longa, linearis, stricta, exacte tetragona, bivalvis, bilocularis. *Semina* plura, parva, subrotunda.

1007. *ERYSIMUM officinale* filiquis spicæ adpressis, foliis runcinatis. Linn. Sp. pl. 2. p. 922. — Pollich. fl. pal. n. 631.

Erysimum foliis pinnatis, pinnis rectangulis acutis, extrema triangulari maxima, filiquis adpressis. Hall. stirp. helv. n. 478.

Erysimum vulgare. Bauh. pin. 100. — Mapp. fl. Als. p. 102.

Erysimon primum. Tabernæmont. p. 840. Icon.

Sisymbrium officinale. Scop. fl. Carn. ed. 2. n. 824. — Roth. fl. Germ. 3. p. 131.

Icon. Curtis. fl. Lond. — Bulliard. herb. t. 259. — Regnault. bot. — Blackw. t. 28.

Germ. Gelber Wegsensf. Gelbes Eisenkraut.

Gall. Le Velar, ou la Tortelle.

Hab. Utrunque in ruderatis ad vias, muros, hortos, sæpes passim abunde.

Flor. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix fusiformi-cylindrica, fibrillosa, alba. *Caulis* sesquipedalis, cubitalis et altior, erectus, teres, striatus, pubescens, scaber, ramulosus. *Rami* simpliciusculi, alterni, divaricati. *Folia* alterna, petiolata, subpubescentia, subtus scabrisuscula, pinnatifida, laciniis oppositis, sursum dentato-lobatis, extima triangulari, majore, trifida. *Flores* caulem ramosque terminantes, racemosi. *Racemi* floriferi subrotundi, fructiferi elongati, nudi. *Corollæ* petala quatuor, cuneiformi-obtusa, lutea, calyce longiora. *Siliquæ* cylindrico-subulatæ, octangulæ, apicem versus tetragonæ, pubescentes, vix unciam longæ, erectæ, cauli adpressæ, polyspermæ. *Semina* ovalia, bruna, splendentia.

Pharm. ERYSIMI Herba; Semen.

Planta acriuscula, incidens, resolvens, in pectoris morbis, raucedine, asthma laudabatur.

Flores melliferi sunt.

1008. *ERYSIMUM Barbarea* foliis lyratis, extimo subrotundo. *Linn. Sp. pl. 2. p. 922. — Pollich. fl. pal. n. 632.*

Erysimum foliis pinnatis, pinna extrema maxima. *Hall. stirp. helv. n. 479.*

Sisymbrium Erucae folio, glabro, flore luteo. *Tournef. infl. 226. — Mapp. fl. Alf. p. 288.*

74 TETRADYN. SILIQUOSA.

Barbarea mas. *Tabernæmont. p. 843. Icon.*
β. *Sifymbrium erucæ folio glabro, minu*
et præcocius Tournef. inst. 226.

Erysimum præcox foliis inferioribus lyratis
 superioribus pinnatifidis, laciniis linearibus
 oblongis integerrimis. *Smith. fl. bri*
2. p. 707.

Icon. Regnault. bot. — Oeder. fl. Dan. t. 98.
Germ. St. Barbaren - Kraut.

Gall. Herbe de St. Barbe.

Hab. Utrunque ad fossas aquaticas passim fre
 quens. **β.** in graminosis, pratis ad fossas
 passim etiam in hortis colitur.

Flor. Majo, Junio. *Semina* Julio maturantur
 7 var. **β** ♂

Radix fusiformi-cylindrica, fibrosa, alba
Caulis pedalis, sesquipedalis et altior, erectus
 sulcato-angulatus, glaber, superne ramosus
Rami alterni, numerosi, virgati. **Folia** alte
 ra, amplexicaulia, glabra, lineata, subclaud
 lyrata, laciniis lineari-lanceolatis, obtusis, in
 tegris, terminali majore, ovata, obtusa, re
 panda. **Flores** caulem ramosque terminantes
 racemosi. **Racemi** florentes ovati, fructiferi
 elongati. **Calyx** tetraphyllus, foliolis linearibus,
 obtusis, lutescentibus, duobus oppositis
 basi gibbulosis. **Corolla** cruciformis, petalobus
 cuneiformibus, obtusis, luteis, calyce dupli
 longioribus. **Glandulae** geminae, subrotundae
 utrinque inter germen et filamen brevius; glandula
 solitaria, conica, utrinque inter filamen
 longiora et calycem. **Siliquae** unciam circiter
 longae, obsolete tetragonae, glabrae, polyspermae.
Semina ovali-orbicularia, flavescentia
 bruna.

Variatio 8. Differt Radice bienni Corollis pallidioribus. Siliquis longioribus. Ob variationes intermedias prostantes, vix peculiarem speciem constituit.

In hortis etiam floribus plenis elegantissimis observatur.

PARM. BARBAREÆ Folia viridia.

Sapor Nasturtii, acris, Antiscorbuticas vires exercet. Pro acetariis apta.

Flores apibus mel præbent.

1009. ERISYMMUM *Alliaria* foliis cordatis. Linn. Sp. pl. 2. p. 912. Pollich. fl. pal. n. 633. Hall. stirp. helv. n. 480.

Silybrium *Alliaria*. Scop. fl. Carn. ed. 2. n. 825. — Roth. fl. germ. 3. p. 131.

Hesperis allium redolens. Monif. hist. 2. p. 252. — Mapp. fl. Alf. p. 142.

Alliaria Bauh. pin. 110. Tabernæmont. p. 1144. Icon.

Icon. Curtis fl. Lond. t. 144. Oeder. fl. Dan. t. 935. — Bulliard. herb. t. 338. — Blackw. t. 372.

Germ. Knoblauchskraut.

Gall. *Alliaire*.

Hab. Utrinque in ruderatis, sepibus, ad vias umbrasas ageres, vineas, sylvas, passim frequens.

Flor. Aprili, Majo. *Semina* Junio, Julio maturantur. ♂

Radix fusiformi-cylindrica, fibrosa, alba. *Caulis* sesquipedalis, cubitalis et altior, erectus, teres, subsiriatus, inferne pubescens, superne glaber, subramosus. *Rami* alterni, quandoque nulli. *Folia* alterna, petiolata, cordata,

76. TETRADYN. SILIQUOSA.

glabra, venosa, laete viridia, inæqualiter sinu-
dentata. *Flores* terminales, racemosi. *Calyx*
foliola lanceolata, obtusa, albido-viridescen-
tia, caduca. *Corolla* tetrapetala, crucifera.
Petala obovata, obtusa, nivea, calyce lon-
ga. *Siliqua* tetragona, polysperma.

Pharm. ALLIARIÆ Herba recens.

Tota planta, digitis trita, Allium redolet.
Hildano et Boerhaavio ad Gangrænam com-
mendam laudabatur.

Lac vaccarum odore alliaceo inficit.

Flores apibus mel concedunt.

1010. ERYSIMUM *Cheiranthoides*

foliis lanceolatis subdenticulatis, sca-
usculis, siliquis patulis.

Erysimum foliis lanceolatis integerrimis.
Linn. Sp. pl. 2. p. 923. — *Pollich. fl.* 1
n. 634. — *Hall. stirp. helv.* n. 477.

Turritis Levcoji folio. *Tournef. inst. p.* 1
Mapp. fl. Alf. p. 314.

Myagrum siliqua longa. *Bauh. pin.* p. 1

Erysimon tertium. *Tabern. p.* 840. *Id.*

Icon. Iacqu. fl. Austr. t. 23. — *Schkuhr.*
handb. t. 183.

Germ. Levcojenartiger Hederich.

Gall. Vêlar giroflier.

Ha b. Utrique in ruderalis, cultis, arvis, a-
vineis passim frequens.

Flor. a Majo, ad Septembrem. *Semina* 8
tembri, Octobri matura. ☉ ♂

Radix fusiformi-cylindrica, fibrosa, alb-
Caulis pedalis, sesquipedalis et altior, erec-
tus versus basin quandoque incurvus, angula
scabriusculus, ramosus. *Rami* alterni. *F.*

parva, alterna, sessilia, lanceolata, acuta, scaberrima, inferiora rariter denticulata, superiora integra, viridia. Flores caulem ramosque terminantes, racemoso-corymbosi, demum elongati. Calyx tetraphyllus, luteus. Corollae inodora, parva, lutea. Stigma capitatum, interiusculum. Siliquae tetragonae, patulae, uncinatae et longiores.

Nota. Variat Caulibus demissioribus et altioribus, plus minusve ramosis, quandoque ramosissimis. Foliis latioribus et angustioribus, interiusculis et subrepando remote denticulatis. Corollarum petalis plus minusve obtusis et emarginatis.

In sabulosis arenam figit.

Flores melliferi.

1011. *ERYSIMUM hieracifolium* caule stricto subramoso, foliis lanceolatis remote argute sinuato denticulatis, filiquis erectis.

Erysimum hieracifolium foliis lanceolatis serratis. Linn. Sp. pl. 2. p. 923. — Pollich. fl. pal. n. 635.

Erysimum hieracifolium foliis lanceolatis remote dentatis, filiquis erectis. Willden. Linn. Sp. pl. 3. p. 511.

Erysimum strictissimum. Wetterauische flora. 2. p. 451.

Turritis Levcoji folio angustiori, filiquis longis erectis? Mapp. fl. Alf. p. 314.

Levcojum luteum sylvestre hieracifolium. Bauh. Prodr. p. 102.

Icon. Oeder. fl. Dan. t. 923.

Germ. Habichtskrautartiger Hederich.

Gall. Vêlar a févilles des épervière.

Hab. In Palatinatu in montibus ac collibus asperis lapidosis circa Bingen et Creuzenach

78 TETRADYN. SILIQUOSA.

abunde post *Pollichium* vidi. In Spon-
mia in collibus præruptis lapidosis ad l-
sellam frequens.

Flor. Junio, Julio. *Semina* Augusto m-
ra. ♂

Radix fusiformi-cylindrica, fibrosa, a
Caulis sesquipedalis, cubitalis, erectus, an-
latus, pilosiusculus, ramosus. *Rami* alte
Folia radicalia numerosa, terrae incumben-
profundius sinuato-dentata, caulinis angustior
caulina alterna, sessilia, lanceolata, rem-
sinuato-denticulata, pilosiuscula, scabriuscu-
Flores racemosi, caulem ramosque termi-
tes. *Calycina* foliola quatuor, conniven-
lanceolata, ex lutescente pallide viridia,
briuscula. *Corollæ* petala speciosa, pater-
obtusula, cum unguibus fere semunciam lor-
integra, pallide lutea, subinodora. *Siliquæ*
tragonæ, erectæ, subpilosulæ, duas fere
cias longæ, stigmate capitato subemargin-
terminatæ.

A præcedente facile distinguitur, Corollis
to majoribus, pallidioribus, Siliquis erectis de-
longioribus.

Nota Planta non confundenda cum *ERYSI-*
hieracifolio Jacquini fl. Austr. t.

Obs. Variat Caulibus simpliciusculis saltem in p-
superiore ramos paucos habentibus, et caul-
a basi ad summitatem usque quammaxime ramo-
Planta ob flores speciosos elegans.
Flores apibus mel præbent.

CCCXIX. CHEIRANTHUS.

Linn. Gen. pl. ed. Schreb. n. 1091.

Calyx tetraphyllus compressus, foliolis
æolatis concavis, erectis, parallelo-conniv-

tes, deciduis, quorum duo exteriora basi
 subba. *Corolla* tetrapetala, cruciformis. *Petala*
 subrotunda, calyce longiora, unguibus longi-
 tudine calycis. *Nectarifera* glandula cingens
 basin staminis minoris utrinque. *Germen* tu-
 berculo utrinque notatum. *Stigma* bipar-
 titum, reflexum, persistens. *Siliqua* lon-
 ga, compressa, bilocularis, bivalvis, Stigmate
 bifido instructa. *Semina* plura, plana, subova-
 ta, margine membranacea.

1012. **CHEIRANTHUS** *Cheiri* foliis lan-
 ceolatis acutis glabris, ramis angulatis,
 caule fruticoso. *Linn. Sp. pl.* 2. p. 924.

Leucojum foliis glabris lanceolatis integer-
 rimis. *Hall. stirp. helv.* n. 443.

Leucojum luteum vulgare. *Bauh. pin.* 202.

Viola petraea lutea. *Tabern. p.* 688. *Icon.*

Icon. Regnault. bot. — *Bulliard. herb. t.* 349.

— *Blackw. t.* 179. (*planta fativa.*)

Germ. Gelbe Violen. Goldlack.

Gall. Giroflier jaune.

Hab. Passim in muris vetustis urbium et pago-
 rum. Muros urbis *Freyburg* *Brigoviae*
 tapetis aurei inslar tegit. In *Basileæ* muris
 non infrequens. Multifariæ variationes in
 hortis hospitantur.

Flor. Majo, Junio. *Semina* Julio, Augusto
 matura. ♂ ♀ 5

Pharm. CHEIRI Flores.

Flores suaveolentes, aromatici, amari. adstring-
 entes, incidentes vires exercent.

80 TETRADYN. SILIQUOSA.

In hortis adhuc præcipue hospitantur CHEIRANTHUS incanus Linn. Die Winter-Levcoje. CH. annuus. Linn. Die Sommer-Levcoje. et CH. maritimus. Linn. Die kleine Levcoje, quæ hortorum areas egregie decorat.

Nota. Frustra hucusque CHEIRANTHUM simoidem Linn. in regionibus rhenan vicinis quæsi.

CCCXX. HESPERIS.

Linn. Gen. pl. ed Schreb. n. 1093.

Calyx tetraphyllus: foliolis lanceolatis, nearibus, parallelo-conniventibus, deciduorum duo opposita basi gibba. *Corolla* trapetala, cruciformis. *Petala* oblonga, reflexa, desinentia in unguem attenuatum longitudine calycis. *Glandula* mellifera, terminata, inter flamen brevius et germen. *Stylus* bipartitus, oblongus, erectus, apice connivens. *Siliqua* longa, compresso-plisfricta, bilocularis, bivalvis. *Semina* plurimovata, compressa.

1013. *HESPERIS matronalis* caule stipulicili erecto, foliis ovato-lanceolatis, denticulatis, petalis mucrone emarginatis. Linn. Sp. pl. 2. p. 927. — Roth germ. 2. p. 107.

Hesperis hortensis. Bauh. pin. 202.

Viola hyemalis flore albo et purpureo. Bernamont. p. 692. Icon 1. 2.

β. Floribus plenis niveis.


Icon. Kniph. cent. 7. n. 32.

Gei

Germ. Nacht-Viole. Pfingst-Viole.

Gall. La Julienne. Cassiole.

Hab. In Sibiria. Colitur cum variatione β . praesertim in rusticorum hortis ex quibus in agros et vineas passim migravit, vix ullibi in Germania spontanea.

Flor. Majo, Junio. *Semina* Julio, Augusto matura. 

Radix cylindrica, fibrosa. *Caules* ex radice saepe plures, cubitales et altiores, erecti, teretes, striati, hirsuti. *Folia* alterna, ovato-lanceolata, dentata, hirsuta, inferiora petiolata, superiora sessilia, semiamplexicaulia. *Racemus* caulem terminans, longus, speciosus. *Calyx* rubello-fuscus, hirsutus. *Corollae* petala emarginata, cum acumine parvo interjecto, pallide violacea, aut purpurea, in variatione β saepissime nivea.

Flores suaveolentes. β . egregium horticorum, praesertim rusticorum, decus.

Arbus maxime melliferi. In hortis hospitatur. *HESPERIS* tristis Linn. caule hispido ramis patente. Die Nacht-Viole omnium odoratissima, cum primis tempore vespertino,

1014. *HESPERIS inodora* caule simplici erecto, foliis subhastatis dentatis, petalis obtusis. *Linn. Sp. pl.* 2. p. 927.

Hesperis caule hirsuto, foliis subasperis, ovato-lanceolatis, serratis, petiolatis. *Hall. stirp. helv. n.* 448.

Hesperis sylvestris inodora. Bauh. pin. 202.
— *Hall. ad Rupp. fl. Jen. p.* 78. t. 1.

Hesperis tertia Clus. hist. p. 297.

Icon. Jacqu. fl. Austr. t. 347.

Germ. Wald-Nachtviole.

Gall. Cassiolette des bois.

Hab. In ditione Basileensi ad Augustiam l
racorum versus Rheni ripam in dum
prope balneum *Bubendorf* ad dextram
Regiæ; in monte *Franspurg* et *Sissac*

Flor. Majo, Junio, Julio. *Semina* Julio,
gusto matura. ♂ ♀.

Radix ramosa, fibrosa, albida. *Caulis*
quipedalis, tripedalis, erectus, teretiuse
aut sulcato-angulatus, plus minusve hirs
simplex aut ramosus. *Rami* alterni, ex
Folia alterna, subhastata, aut cordato-lan
lata, acuminata, argute subsinuato-den
ferrata, inferiora longius, superiora brev
petiolata, aut subsessilia, utrinque villosa, s
riuscula. *Flores* caulem ramosque terminat
racemosi, suaveolentes. *Calyx* fusco-pu
rascens, quandoque viridis, pilosohirs
Corollae Petala obtusa, integra, subemarg
tate, albida, incarnata, purpurea, purpu
violacea. *Siliquae* teretiuseculae, obsolete
tetragonae, striatae, glabrae, subtorulo
uncias binas ad quatuor longæ, polysper
Semina oblonga, fusca.

Nota. Non confundenda cum præcedente sp
a qua differt, Foliis majoribus cordatis
hastatis, acuminatis, argute subsinuato-den
ferratis, minus scabris.

Flores non inodori, sed suave odorati, et id
fertim in umbrosis.

Apibus melliferi.

Ad rariores omnino Floræ cives pertinet. Ad
dumeta ad arcium vetustarum rudera,

TETRADYN. SILIQUOSA.

23

CCCXXI. A R A B I S.

Linn. Gen. plant. ed. Schreb. n. 1094.

Calyx tetraphyllus, deciduus: *foliolis* parallelis-conniventibus: duobus oppositis majoribus, ovato-oblongis, acutis, basi prominulis: duobus reliquis linearibus, erectis. *Corolla* tetrapetala, cruciformis. *Petala* ovata, patentia, desinentia in ungues longitudine calycis. *Nectaris* quatuor: singulum ex *squamula* intra *fundum* folioli calycini receptaculo affixa, reflexa, persistente. *Stigma* obtusum, integrum. *Stylus* compressa, longissima, linearis, torulis *asymmetria* inaequalis: *Valvulis* longitudine ferme *aequalibus*. *Semina* plurima, subrotunda, compressa.

1015. **ARABIS** *alpina* caule ramoso, ascendente, foliis amplexicaulibus cordato-hastatis sinuato-dentato-ferratis hirsutis.

Arabis alpina foliis amplexicaulibus dentatis.

Linn. Sp. pl. 2. p. 928.

Arabis caule ramoso diffuso, foliis amplexicaulibus, dentatis, incanis. *Hall. stirp. helv. n. 451.*

Draba secunda. *Clus. hist. 2. p. 125. Icon. bona.*

Icon. Oeder. fl. Dan. t. 62.

Ger. m. Alpen-Gaensekraut.

Gall. Arabette des alpes.

Hab. In montosis rupestribus vallis *Höllenthal* *Brissgovia*. In Ditione *Basilicensi* in rupe-

stribus altioribus prope *Dornach*, *Ramstein*, *Wasserfall*, *Vogelberg*, *Neunbrunn*, *Langenbruck* et alibi abunde. In montis calcareis prope *Schaffhausen* vidi.

Flor. Aprili, Majō, *Semina* Julio, matura.

Radix fibrosa, luteo-albida. *Caulis* digitalis, palmaris, pedalis, sesquipedalis, ascendens, teretiusculus, hirsutus, simpliciusculus, sæpissime ramosus. *Rami* alterni, diffusifolia radicalia numerosa, in petiolum breviter decurrentia, ovata, caulina alterna, sessilia amplexicaulia, cordato-aut hastato ovata, sinuato-dentato-ferrata, obtusa, omnia hirsuta laete viridia, quandoque ex viridi incana, lutescentia. *Flores* caulem ramosque terminantes, corymbofo-racemosi, demum elongati racemosi. *Calycis* foliola bina, ovata, deorsum gibba, pallide viridia, hirsuta. *Corollæ* petala ovata, obtusa, alba, ungue viridia, calyce duplo longiora, *Siliquæ* tenues, curvæ, compressæ, torulosæ, glabræ, polyspermæ, binas ad tres uncias longæ. *Semina* ovilia, compressa, fusca.

Obs. Planta in alpinis altioribus sæpe nana, caule simplicissimo, foliis minoribus incanis prædita.

In supalpinis præsertim calcareis, major, sesquipedalis, maxime ramosa, Ramis ex basi caulis præsertim copiosis ad terram diffusis.

Semina ex individuis pusillis simplicibus alpinis Helvetiæ et Pyrenæorum lecta, in Horto Botanico tunc commissa, reddiderunt individua pedalia sesquipedalia quammaxime ramosa, nostratibus similia.

1016. *ARABIS Turrita* caule erecto simpliciusculo, foliis amplexicaulibus dentatis, filiquis prælongis, maturis unguis recurvatis.

Arabis Turrita foliis amplexicaulibus, siliquis decurvis planis linearibus margine incrassatis. *Smith. fl. Brit.* 2. p. 714.

Arabis Turrita foliis amplexicaulibus, siliquis decurvis planis, linearibus, calycibus subrugosis. *Linn. Sp. pl.* 2. p. 930.

Arabis umbrosa. siliquis compressis latis, uno versu pendulis. *Crantz. Austr.* p. 39. t. 3. f. 2.

Leucojum foliis acute dentatis, amplexicaulibus, siliquis nutantibus incanis. *Hall. stirp. helv.* n. 444.

Turritis major, Plateau. *Clus. hist.* 2. p. 126. *Icon.*

n. Jacqu. fl. Austr. t. 11.

m. Thurmkrautartiges Gänsekraut.

l. Arabette Tourelle.

b. In Ditionis Basileensis rupestribus umbrosiscirca *Wallenburg, Langenbruck, Ramstein, Dornach* et alibi frequens; Basileæ in muris Collegii Erasimiani.

r. Sub fine Aprilis, Majo. *Semina* Junio, Julio matura. *z*

Radix ramoso-fibrosa, extus pallide fusca. *stems* sesquipedalis, cubitalis, erectus, teretulus, simplex, rarius ramo uno alterove ditus, villosus-tomentosus. *Folia* radicalia, linæque infima in petiolum decurrentia, oblongo-lanceolata, acuta, margine inæquasi-nuato-dentata, subundulata; *caulina* remota alterna, sessilia, amplexicaulia, cordato-ovata, remote denticulata, omnia villosa-tomentosa, pallide viridia. *Flores* terminales, cymosi. *Calycina* foliola alterna basi gibbula, parviscula aut pilis raris munita, pallide viridia.

Corollæ petala oblonga, submarginata, alba aut pallide ochroleuca, calyce longiora. *Stigmata* linearis, compressa, quatuor ad quinque uncias longæ, obtusa, glabræ, feminibus torulosa, maturæ uno versu decurvata, polysperma. *Semina* elliptica, submarginata, compressa margine membranacea, fusca.

Nota. Non confundenda cum *A. alpina*, a qua differt, Caule altiori, crassiori, semper simpliciter erecto. Foliis longioribus, latioribus exacte dentatis, nec dentato-serratis. Siliculis longioribus, latioribus, maturis uno versu decurvatis.

1017. *ARABIS thaliana* caule erecto ramoso basi hirsuto, foliis radicalibus petiolatis ovali-lanceolatis integris denticulatisve, caulinis alternis sessilibus lanceolatis integriusculis, superioribus lanceolato linearibus integris, omnibus ciliatis-hispidis, petalis calyce duplo longioribus.

Arabis thaliana foliis petiolatis lanceolatis integerrimis. *Linn. Sp. pl.* 2. p. 929. — *Pollich. fl. pal.* n. 636. cum *Icone*.

Arabis foliis radicalibus petiolatis ovatis dentatis, caule subnudo ramoso. *Hall. stirp. helv.* n. 452.

Turritis vulgaris ramosa. *Mapp. fl. Alf.* 313.

Bursa pastoria five *Pilosella filiquosa*. *Bauh. hist.* 2. p. 870.

Icon. *Curtis. fl. Lond.* — *Oeder. fl. Dan.* t. 1106 — *Schkuhr. bot. handb.* t. 185.

Germ. Gänsekraut. *Thurnsenf.*

Gall. Arabette rameuse.

Hab. Utrunque in agris et arvis arenosis aliisque passim frequens. Circa *Carlsruhe* et alibi abunde.

Flor. Aprili, Majo, Junio. *Semina* Augusto, Septembri matura. ☉

Radix fibrosa, albida. *Caulis* palmaris, pedalis, sesquipedalis et altior, erectus, teres, rore glauco tectus, inferne hirsutus, superne glabrescens, ramosus aut simpliciusculus. *Rami* alterni, superne subnutantes. *Folia* radicalia petiolata, ovato-oblonga, versus basin præsertim dentata, setuloso-hirsuta, punctis proximalis utrinque scabra; *caulina* alterna, sessilia, inferiora lanceolata, denticulata aut integra, superiora lineari-lanceolata, integra, setuloso-ciliata. *Flores* caulem ramosque terminantes, racemosi. *Racemi* deflorati elongati. *Calicis* foliola ovata, concave, hirsuta. *Corollæ* petala quatuor, obtusa, integra, alba, ungue viridia, calyce duplo longiora. *Glandulæ* nectariferae sex, minutæ. *Siliquæ* cylindrico-subtetragonæ, graciles, patentes, septem ad novem lineas longæ, glabræ, polyspermæ. *Semina* parva, subrotunda, fusco-flavicantia.

Obs. Bis sæpe eodem anno floret, Aprili et Septembri.

In muris, rupibus, lapidosis sæpe major cubitalis fere, magis ramosa; in sterilibus arenosis minor fructuoscula.

CCCXXII. T U R R I T I S.

Linn. Gen. pl. ed. Schreb. n. 1095.

Calyx tetraphyllus: *foliolis* ovato-oblongis, parallelo-conniventibus, deciduis. *Corolla* tetrapetala, cruciformis. *Petala* ovato-oblonga, obtusa, erecta, integra: *unguibus* erectis. *Styl-*

lus nullus. *Stigma* obtusum. *Siliqua* omnium longissima, stricta, tetragona, angulis oppositis alternis obsoletis et subcompressis, bilocularis, bivalvis: *Valvulis* dissepimento vix æqualibus. *Semina* numerosa, rotundiuscula, emarginata.

1018. *TURRITIS glabra* foliis radicalibus dentatis hispidis, caulinis integerrimis amplexicaulibus glabris. *Linn. Sp. pl.* 2. p. 930. — *Pollich. fl. pal. n.* 637. — *Hall. stirp. helv. n.* 455,

Turritis foliis inferioribus cichoraceis, cæteris perfoliatis. *Mapp. fl. Alf. p.* 315.

Brassica sylvestris foliis circa radicem cichoraceis. *Bauh. pin.* 112.

Turritis Tabernæmont. p. 849. *Icon.*

Icon. Curtis fl. Lond. t. 253.

Germ. Glattes Thurmkraut. Thurm Kohl.

Gall. Tourelle glabre.

Hab. Utrique passim in dumetosis arenosis

TETRADYN. SILIQUOSA.

89

Herb. lineari-lanceolata, obtusiuscula, convexa, erecta, ex viridi-albicantia, glabra. *Corollæ* petala lanceolata, obtusa, ex viridi albida, calyce longiora. *Siliquæ* prælongæ, lineares, erectæ, cauli approximatae, glabræ, polyspermæ.

Fæta arenam præfertim ad præcipitia figit. Flores quibus melliferi.

1019. **TURRITIS** *hirsuta* foliis omnibus hispidis, caulinis amplexicaulibus. *Linn. Sp. pl.* 2. p. 930. — *Pollich. fl. pal.* n. 638. — *Hall. stirp. helv.* n. 456.

Arabis hirsuta. *Scop. fl. Carn.* ed. 2. n. 835. *Roth. fl. germ.* 2. p. 112.

Turritis. *Lob. Icon.* 220. *Mapp. fl. Alf.* p. 313

Turritis minor. *Mapp. fl. Alf.* p. 313.

Erysimo similis *hirsuta alba.* *Bauh. prodr.* p. 42. *Icon.*

Icon. *Oeder. fl. Dan.* t. 1040. — *Schkuhr. bot. Handb.* t. 185.

Ger. Rauchhariges Thurmkraut.

Gall. Tourelle vélue.

Herb. Utrunque passim in dumetis gramineis, pratisque inæqualibus apricis et collibus dumetosis frequens. Circa *Carlsruhe* ad vias prope *Beurtheim* in dumetis graminosis.

Flor. Majo, Junio. *Semina* Julio matura. ♂

Radix fibrosa, albida. *Caulis* palmaris, perisetalior, erectus, teres, hirsutus, simplex, rarius ramosus. *Folia* radicalia in rosulam congesta, petiolata, ovata, caulina alterna, sessilia, amplexicaulia, basi subsagittata, omnia lanata, hirsuta. *Flores* terminales, corymbosomæ. *Calycina* foliola albide marginata,

Corollæ petala parva, albida. *Siliquæ* longæ, graciles, subtetragonæ, juniores cauli adpressæ, fere, adultiores nutantes.

Herba tenella ovibus accepta. Flores apibus melliferi.

Nota. Semper hirsuta, quandoque foliis integriculis instructa est.

CCCXXIII. B R A S S I C A.

Linn. Gen. plant. ed. Schreb. n. 1096.

Calyx tetraphyllus: foliolis lanceolatis, nearibus, concavo-canaliculatis, basi gibbi erectis, parallelis, deciduis. *Corolla* tetrapetala, cruciformis. *Petala* subovata, plana, persistentia, integra, in ungues sensim attenuata, longitudine fere calycis. *Glandulæ* nectariferæ quatuor, ovatæ, quatum una utrinque interfiamina longiora et calycem posita. *Siliqua* longior, teretiuscula, depressa utrinque, *Stylo* longiore obtuso, in quibusdam ensiformi coronato, dissepimento apice prominenti tereti, bilocularis, bivalvis: *valvulis* dissepimento brevioribus. *Semina* plura globosa.

† *Stylo* obtusiusculo.

1020. *BRASSICA orientalis* foliis cordatis amplexicaulibus glabris: radicalibus scabris integerrimis, siliquis tetragonis. Linn. Sp. pl. 2. p. 931. — Pollich. fl. pal. n. 639.

Eruca foliis ovatis integerrimis amplexicaulibus. Hall. stirp. helv. n. 457.

Brassica Turrita foliis oblongis cordato-sagittatis amplexicaulibus, siliquis tetragonis caulem superantibus strictissimis.
Weig. obs. bot. p. 32.

Brassica campestris perfoliata, flore albo.
Bauh. pin. 112. — Mapp. fl. Alf. p. 42.

Icon. Jacqu. fl. Austr. t. 282.

Germ. Orientalischer Kohl.

Gall. Chou d'Orient.

Hab. Utrunque in agris et arvis passim non infrequens.

Flor. Junio, Julio. *Semina* Augusto maturata. ☉ ♂

Radix fusiformi-cylindrica, fibrosa, albida. *Caulis* pedalis, sesquipedalis, cubitalis, et altior, erectus, teres, glaberrimus, glauco-viridis, simplex, vel ramosus. *Folia* radicalia sessilia, obovata, obtusa, caulinis minora, caulina alterna, amplexicaulia, perfoliata, ovalia, obtusa, omnia integerrima, ex glauco-viridia, crassiuscula, glaberrima. *Flores* terminales, racemosi. *Calyx* quandoque subpurpureus, erectus. *Corollæ* petala calyce longiora, ex viridi-albida. *Siliquæ* exacte tetragonæ, prælongæ, erecto-patentes, glabræ.

1021. *BRASSICA alpina* foliis caulinis sessilibus cordato-sagittatis amplexicaulibus, radicalibus petiolatis ovatis, omnibus integerrimis, petalis erectis.

Brassica alpina foliis caulinis cordato-sagittatis amplexicaulibus, radicalibus ovatis, petalis erectis. *Linn. Mant. 1. p. 95. — Poll. fl. pal. n. 640. —*

Turritis foliis glaberrimis, curvis, ovatis, rotundis, petiolatis, superioribus am-

plexicaulibus, sagittatis obtusis. *Hal stirp. helv. n. 454.*

Turritis *Brassica* foliis omnibus glabris: caulinis cordato-sagittatis amplexicaulibus radicalibus ovatis petiolatis. *Leers. fl. Herb. n. 518.*

Brassica alpina perennis. *Mapp. fl. Alf. p. 41*
Icon. Villars fl. Delph. 3. t. 36.

Germ. Alpenkohl.

Gall. Chou des alpes.

Hab. In Palatinatu in sylvis montosis lapidosi
circa *Steinbach* ad montem *Donnersberg*
cum *Turritide glabra* copiose t. *Poll*
chio. In sylvis circa *Lautercken.* In A
satia in montosis sylvis versus *Niedel*
retro *Haslach t. Mappo.*

Flor. Majo, Junio. *Semina* Augusto matura.

Radix fibrosa, albida. *Caulis* sesquipedali
cubitalis et altior, erectus, teres, glaber, si
plex. *Folia* radicalia petiolata, ovata, obtus
integra, caulina alterna, sessilia, amplexica
lia, cordato-sagittata, acuminata, integerrim
omnia glabra, ex glauco-viridia. *Flores* ca
lem terminantes, racemoso-corymbosi. *Cal*
cis foliola erecta, pallide flavescentia, a
purpurascentia; quorum duo latiora co
vexa, basi gibbula. *Corollæ* albæ, parvin
culæ. *Petala* lineari-oblonga, obtusa, c
lyce duplo longiora. *Staminum* filamenta d
breviora; basi glandulis duabus instructa. *A*
theræ sagittatæ, flavæ. *Siliquæ* tetragonæ, line
res, glabræ, ultra duas uncias longæ, erect
in pedunculo patente, *Stylo* brevissimo obtu
instructæ. *Semina* numerosa, obovata, co
pressa, fusco-flavicantia.

Nota. Non confundenda cum Turritide glabra Linn. cui fere similis est.

Ad rariores omnino Floræ cives pertinet.

1022. *BRASSICA arvensis* foliis amplexicaulibus spathulatis repandis: summis cordatis integerrimis. *Linn. Mant.* 95.

Brassica campestris perfoliata, flore purpureo. *Bauh. pin.* 112. *Mapp. fl. Alf. p.* 42.

Perfoliata siliquosa purpurea. *Clus. Joh. Bauh.* 2. p. 836.

Brassica campestris purpureo flore. *Clus. hist.* 2. p. 127.

Germ. Purpurrother Feldkohl.

Gall. Chou à fleur pourpré.

Hab. In Alsatia? (Circa Argentinam in campestribus prope *Oswald t. Mappo.*)

Flor. Aprili, Majo, Junio. *Semina*, Julio matura. ☉

Corollis amoene violaceo-purpurascensibus facile dignoscitur.

Nota. *Brassica arvensis* vera. vix unquam in Germania sponte crescit. Plantam a *Mappo. l. c.* allegatam, frustra prope Argentinam quæsi, quæ abunde in campestribus Hispaniæ, præsertim in regno Valentino et Murciano occurrit. Hinc non immerito ex catalogo civium germaniæ delenda est.

Brassicam campestris Linn. hucusque frustra in regionibus rhenanis et confinibus quæsi.

1023. *BRASSICA Napus* radice caulescente fusiformi, foliis lævibus, superioribus cordato-lanceolatis amplexicaulibus, inferioribus lyratis dentatis. *Smith. fl. brit.* 2. p. 719.

Brassica Napus radice caulescente fusiformi. *Linn. Sp. pl. 2. p. 931.*

Napus sylvestris. Bauh. pin. 95. — Mapl fl. Alf. p. 208.

β. *Napus sativa. Bauh. pin. 95.*

Icon. Regnault. bot. Blackw. herb. t. 224.

Germ. Wilde Rübsaat. Rübſen. Rebs. wild Steckrüben.

β. *Zahme Rübsaat. Rebs. Sommer oder Winterrübsaat oder Rebs.*

Gall. Navet sauvage. β. Navet cultivée.

Hab. In agris praesertim sabulosis, ad agrorum marginalia passim non infrequens. β. copiose colitur.

Flor. Aprili, Majo. *Semina* Julio matura. ☉

Planta culta notissima, agros deaurans, flores odorata. Seritur semen primo vere in agris bene cultis et stercoretis, tunc versus autumnum floret, et Septembri semina maturat. Haec vulgo denominatur Sommerrebs. Aut seritur autumnno, tunc sequenti Aprili et Majo floret, et Junio Semina maturat, haec Winterrebs audit, et ejus cultura praefertur.

Pharm. NAP1 Semina.

Hodie fere in medicina exoleta.

Ex seminibus oleum unguinosum pressione obtinetur.

Flores melliferi, apibus quammaxime accepti.

Planta adhuc tenella, oleracea.

1024. *BRASSICA Rapa* radice caulescente orbiculari depresso, carnosa, folii radicalibus lyratis scabris, caulinis integerrimis laevibus. *Smith. fl. brit. 2. p. 719*

Brassica Rapa radice caulescente orbiculari depresso, carnosa. *Linn. Sp. pl. 2. p. 931.*

Rapa fativa rotunda. *Bauh. pin.* 89.

Rapum. *Camerar. epit. p.* 218.

3. Rapa fativa oblonga f. femina. *Bauh. pin.* 89.

Icon. Regnault. bot. — Blackw. t. 226.

Germ. Rübe. Gemeine Rübe. Weisse Rübe.

Gall. Rave.

Hab. In arvis Angliæ et Belgii. Colitur passim copiose.

Flor. Majo, Junio, Julio. *Semina* Augusto Septembri matura. ☉ ♂

Hujus multæ variationes majoribus minoribusque radicibus, rotundis, fusiformibus, coluntur.

Pharm. RAPÆ f. RAPI Radix recens.

Radix culinaria. Decoctum raparum saccharo edulcoratum in aphthis medicina domestica præstans. Rapæ assatæ, calidæ pernionibus impositæ, celebrantur.

1025. *BRASSICA oleracea* radice caulescente tereti carnosa, foliis omnibus glabris glaucis repandis lobatisve. *Smith. fl. brit. 2. p.* 790.

Brassica oleracea radice caulescente tereti carnosa. *Linn Sp. pl. 2. p.* 932.

Brassica (oleracea sylvestris) maritima arboresca f. procerior ramosa. *Morif. hist. 2. p.* 208.

1. *Brassica (oleracea viridis)* alba vel viridis. *Bauh. pin.* 111.

2. *Brassica (oleracea rubra)* capitata rubra. *Bauh. pin.* 111.

3. *Brassica (oleracea capitata)* capitata alba. *Bauh. pin.* 111.

4. *Brassica (oleracea sabauda) alba crisp*
Bauh. pin. 111. undulata. Renealm.
pl. 113.
 5. *Brassica (oleracea laciniata) laciniata* r
bra. *Bauh. hist. 2. p. 832.*
 6. *Brassica (oleracea selenisia) angustio ap*
folio. *Bauh. pin. 112.*
 7. *Brassica (oleracea fabellica) fimbriat*
Bauh. pin. 112.
 8. *Brassica (oleracea botrytis) cauliflor*
Bauh. pin. 111.
 9. *Brassica (oleracea asparagoides.) Baul*
pin. 111.
 10. *Brassica (oleracea Napobraffica) radi*
napiformi. *Tournef. inst. 219.*
Napo - brassica. *Bauh. pin. 111. prodr. 54.*
 11. *Brassica (oleracea gongylodes) gongy*
lodes. *Bauh. pin. 111.*
Brassica caulo Rapa. Bauh. hist. 2. p. 832.
Germ. Kohlkraut. Großer Kohl.
1. *Weisser oder grüner Kohl. Weis*
kraut.
 2. *Rother Kopfkohl. Rother Kappes.*
 3. *Weisser Kopfkohl. Weisser Kappes.*
 4. *Weisser Wirsing.*
 5. *Rother Wirsing.*
 6. *Brauner Kohl. Blaukohl. Federkohl.*
 7. *Grüner Wirsing. Herzkohl.*
 8. *Blumenkohl. Käsekohl.*
 9. *Broccoli. Weisser, grüner und schwar*
zer Broccoli oder Spargelkohl.
 10. *Erdkohlraben. Kohlraben unter der*
Erde.
 11. *Kohl-*

11. Kohlraben. Kohlraben über der Erde.
 Gall. 1. Chou blond ou Chou verd.
 2. Chou-rouge.
 3. Chou-blanc.
 4. Chou-frisé blanc.
 5. Chou-frisé rouge.
 6. Chou-Plumage. Chou-brun.
 7. Chou-fimbrié.
 8. Chou-fleur.
 9. Broccoli.
 10. 11. Chou-raves.

Hab. In Angliæ maritimis. Colitur cum multis variationibus.

Flor. Aestate. Semina Julio, Augusto matura. ☉ ♂

Planta notissima culinaris. Brassica alba capitata laevis, in nostris regionibus et alsaticis muria condita: Sauerkraut audit.

Pharm. BRASSICA CAPITATA ALBA et RUBRA.
 Muria condita, nautis egregium antiscorbuticum.

†† *Erucæ siliquis stylo ensiformi.*

1026. BRASSICA *Erucastrum* foliis dentato-pinnatifidis, caule hispido, siliquis laevibus. Linn. *Sp. pl.* 2. p. 932. — *Pollich. fl. pal.* n. 641.

Erucæ caule hirsuto, foliis hirsutis semipinnatis, pinnis subrotundis dentatis. Hall. *fl. helv.* n. 459.

Erysimum Erucastrum. Scop. *fl. Carn. ed.* 2. p. 27. n. 827. — Roth. *fl. Germ.* 2. p. 104.

Eruca sylvestris major lutea, caule aspe
Bauh. pin. 98. — *Mapp. fl. Als p.* 101

Eruca sylvestris. *Tabernæmont.* p. 837. *Icon. Bulliard herb.* t. 331.

Germ. Rauenkohl. Wilde Rauke.

Gall. Roquette sauvage.

Hab. Utrunque in ruderatis, ad agrorum n-
gines et vias, in hortis et vineis passim
quens.

Flor. Junio, Julio, Augusto. *Semina* Augusto
Septembri matura. ☉

Radix cylindrico-fusiformis, fibrosa, al
Caules ex radice plures, palmares, pedalesquipedales et altiores, erecti, adscenden
 quandoque decumbentes, teretes rariter l
 futi, quandoque glabri, ramosi. *Rami alter*
Folia radicalia numerosa, caulina alterna, g
 briuscula, aut rariter pilosula, pinnatifida, l
 mentis alternis oppositisve lineari-lanceola
 obtuse dentatis: extimo trifido. *Flores* caul
 ramosque terminantes, racemosi. *Corollæ* lut
Siliquæ patentes, obtuse tetragonæ, cornu
 apice capitatae, læves.

Nota. Differt in locis humidioribus arenosis, Corollis altioribus, sulcatis, erectis, infernalis raris obfistis, Foliis majoribus, Corollis saturate luteis, majoribus. Haec varietas sub Sifymbrio obtusangulo recepta in Linn. Sp. pl. ed. Willdenowii. 3. 504. Huc pertinet a Taberaea montana cuncta planta, nec non Eruca inodora. Joh. Bauh. hist. pl. 2. p. 862. Icon.

CCCXXIV. S I N A P I S.

Linn. Gen. plant. ed. Schreb. n. 1097.

Calyx tetraphyllus, patens: *foliolis* linearibus, concavo-canaliculatis, deciduis. *Coro*

Stipula, cruciformis. *Petala* subrotunda, *petala*, patentia, integra: *unguibus* rectis, linearibus, longitudine vix calycis infidentia. *Glandulae* nectariferae quatuor, quarum una utrinque inter stamen brevius et pistillum, et una utrinque inter stamina longiora et calycem posita. *Siliqua* oblonga, inferne torosa, scabra, bilocularis, bivalvis: *Dissepimento* valvulis duplo longiore, magno, compresso. *Semina* parva, globosa.

8097. *SINAPIS arvensis* siliquis multangulis toroso-turgidis rostro ancipiti longioribus. *Linn. Sp. pl.* 2. p. 933. — *Pollich. fl. pal.* n. 642. *Hall. stirp. helv.* n. 467, *Pistillum* flore luteo. *Bauh. pin.* 95.

Sinapi arvense praecox semine nigro. (foliis incisis vel sinuatis.) *Mapp. fl. Alf.* p. 286.

Sinapi arvense praecox semine nigro foliis integris. *Mapp. l. c.*

Icon. Curtis fl. Lond. t. 321. — *Oeder. fl. Dan.* t. 783.

Germin. Ackerfensf. Wilder Sensf.

Call. Moutarde des Champs.

Nat. Utrinque in agris arvis et vineis interdum nimis copiose.

Flor. Majo, Junio. *Semina* Julio, Augusto matura. ☉

Caulis pedalis, sesquipedalis et altior, erectus, teres, striato-fulcatus, hispidus, ramosa. *Folia* alterna, inferiora petiolata, superiora sessilia, sinuato-dentata, infima subpinnata,

100 TETRADYN. SILIQUOSA.

omnia hispida. *Flores* caulem ramosque tenantes, racemosi. *Corollæ* luteae. *Petala* tentia. *Siliquæ* teretiusculæ, subangulatæ, laevæ aut hirsutæ, rostro duplo breviori lævi subtrigono compressiusculo terminatæ. *Sen* subrotunda, nigricantia, parva.

Variet. Foliis plus minus profunde incis.

Ob nimiam sæpe in agris abundantiam frumentum vera pestis.

Flores apibus melliferi.

Bonum pecoribus pabulum præbet.

1028 *SINAPIS alba* filiquis hispidis, rostro obliquo longissimo. *Linn. Sp. p.* p. 933. — *Hall. stirp. helv. n.* 466.

Sinapi album filiqua hirsuta femine alberusco. *Bauh. hist. 2. p.* 858.

Sinapi apiifolio. *Tabernæmont. p.* 838.

Icon. Blackw. herb. t. 29.

Germ. Weisser Senf. Englischer Senf.

Gall. Moutarde blanc.

Hab. Passim rarius inter segetes. Colitur hinc inde.

Flor. Majo. *Semina* Julio matura. ☉

Facile dignoscitur a sequenti specie, *Calibus* succulentioribus, *Corollis* majoribus, *liquis* a caule magis distantibus, brevioribus hispidis, rostro longiore obliquo, ensiformi fructis.

Pro usu medico, culinari et domestico, semina ut ex sequente specie recipiuntur.

Flores apibus mel præbent.

1029. *SINAPIS nigra* filiquis glabris, semine adpressis. *Linn. syst. veg.* 503.

TETRADYN. SILIQUOSA.

101

Sinapis nigra filiquis glabris apice tetrago-
nis. *Linn. Sp. pl.* 2. p. 933. *Pollich. fl.*
pal. n. 643. *Hall. stirp. helv.* n. 465.

Sinapi filiqua latiuscula glabra, semine rufo
et vulgare. *Bauh. hist.* 2. p. 856. *Icon.*

Sinapi rapifolio. *Tabernæmont.* p. 838. *Icon.*

Sinapi *Regnault bot.* *Blackw.* t. 446.

Sinapi *Schwarzer Senf.* Gelber oder rother
Senf.

Call. La Moutarde noir.

Hab. In agris passim infrequens. Colitur hinc
inde.

Flor. Junio, Julio. *Semina* Augusto matura. ☉

Radix fusiformi-cylindrica, fibrosa, al-
ba. *Caulis* 2-3 pedalis et altior, erectus,
terre glabriusculus, ramosus. *Folia* alterna,
petiolata, infima pinnata, scabriuscula: pinnis
alternis, sessilibus, distantibus, dentatis, exte-
rioribus maxima, semitriloba; caulina superiora mi-
nora, suprema subsessilia, integriuscula. *Flores*
nunc ramosque terminantes. *Corollæ* luteæ,
parvæ. *Siliquæ* cauli adpressæ aut paralellæ,
nunc, toroso-turgidæ, subtetragonæ, com-
pessæ, læves. *Semina* fusco-ferruginea, glo-
bulosa, undique minutim rugosa.

Variat ad ripas sabulosas dumetosas *Caulibus*
terrebus tenuioribus, *Foliorum* foliolis magis di-
versis.

Sinapi *Sinapios Semina.*

Semina acrida, calida, rubefaciencia, vesicantia,
et. Cum panis fermento et aceto vini parata massa,
et *Sinapismi* nomine cuti imponitur.

et colinis pulvis feminum cum aceto aut uvarum
et juxta carnes editur.

102 TETRADYN. SILIQUOSA.

Semina pressione exhibent oleum lapidum, et facile rancescens, et destillatione oleum fragrans, et Flores apibus maxime melliferi.

1030. *SINAPIS incana* filiquis racematis adpressis lævibus, foliis inferioribus ratis scabris, summis lanceolatis, et scabro. *Linn. Sp. pl. 2. p. 934.*

Sinapi foliis radicalibus pinnatis, hirsutis incanis, filiqua cornu suo breviori. *Stirp. helv. n. 463.*

Erysimum foliis subincanis filiquis brevissimis. *Herm. parad. 115. Vaillant. b. parif. 51.*

Icon. Jacq. Hort. Vind. t. 169.

Germ. Graublättriger Senf.

Gall. Moutarde grise.

Hab. In Ditione Basileensi copiose prope fileam in agro extra dem *Spalenthor* versus *Alschweiler*, nec non prope *Mulhusian* *Hofero*.

Flor. Junio, Julio. *Semina* Augusto matura.

Radix fusiformi-cylindrica, tenuis, fibrosa, albida. *Caulis* cubitalis et altior, erectus, brevissime villosus, glaucus, asper, tomentosissimus. *Rami* sparsi, brachiati, diffusi, asperi, glauco-incani. *Folia* radicalia caulinaque infima profunde pinnatifida, lyrata, segmentis versus apicem sensim majoribus, inæqualiter dentatis, obtusis, extimo maximo, subrotundato; caulina superiora subovata, vel oblonga, minora, inæqualiter sinuato-dentata ad basin sæpe auriculata; summa lanceolata acutiuscula, omnia petiolata villosulo-aspera, canescentia. *Flores* caulem ramosque terminantes, racemosi. *Racemi* virgati. *Corolla* petala lutea, obovata, calyce duplo longior.

TETRADYN. SILIQUOSA. 103

Siliquæ cauli adpressæ, oblongæ, compressæ, angustæ, tenues, breves, rectæ, rostro brevi obtuso instructæ, villosulæ, maturæ glabriusculæ, 5-7 spermae.

Nota. Ad rariores Floræ cives pertinet.

Obt. individua in Gallia meridionali et Hispania lecta, nostratibus simillima.

Planta in horto botanico culta, mutationi non succedit.

CCCXXV. R A P H A N U S.

Linn. Gen. pl. ed. Schreb. n. 1098.

Calyx tetraphyllus, erectus: foliolis oblongis, conniventibus, deciduis, basi gibbis. Corolla tetrapetala, cruciformis. Petala obcordata, patentia, unguis calyce paulo longiores. Glandulae nectariferae quatuor: duo inter stamina breviora et pistillum, duo inter stamina longiora et calycem. Siliqua oblonga, cum torulis ventricosa, subarticulata, terminalia subrotunda glabra.

1001. RAPHANUS *Raphanistrum* filiquis teretibus articulatis lævibus unilocularibus. Linn. Sp. pl. 2. p. 935. Pollich. fl. pal. n. 644. Hall. stirp. helv. n. 468.

Raphanistrum flore albo striato filiqua articulata. Mapp. fl. Als. p. 266.

Rapistrum flore albo striato, filiqua articulata. Bauh. pin. 95.

1. Flore albide incarnato s. purpureo. Rapistrum purpureum. Tabernæmont. p. 792. Icon.

γ. Flore luteo aut flavidiusculo firiato.

Raphanistrum segetum, flore luteo, vel
lido. *Mapp. l. c.*

Lampfana flore melino. *Tabernamont. p. 7*
Icon.

Icon. Curtis fl. Lond. t. 267. — Schkahr. b
handb. t. 188.

Germ. Hederich. Akerrettig.

Gall. Radis sauvage. Raifort sauvage.

Hab. In agris et arvis hinc inde nimis abundans
γ. amat agros colliculolos argillaceo-
careos.

Flor. Majo, Junio, Julio, Augusto. Sep-
tembri. *Semina* Julio, Augusto, Septem-
bri. Novembri matura. ☉

Radix fusiformi-cylindrica, tenuis, ter-
fa, albida. *Caulis* pedalis, sesquipedalis et
tior, erectus, teretiusculus, subangulosus, h
saepe rubellus, rariter piloso-hispidus, qu
doque glabriusculus, ramosus. *Rami* alter
Folia alterna, inferiora petiolata, pinnatifi-
sinuato-dentata, saepe basi auriculata; sup-
riora subseffilia, lyrata, inciso-ferrata, omni-
praesertim ad costam mediam et nervos pi-
rariis hispidi. *Flores* caulem ramosq
sterminantes, racemosi, laxiusculi. *Corollae*
tentes, albae, dilute carnea, venis violac
aut nigricantibus pictae, aut lutescentes. S-
quae erectae, teretes, articulatae, attenuat
glabrae, apice rostro compresso instructae.

Planta agris frumentaceis ob nimiam abundantiam
saepe noxia. Juniores evulsione repetita exstirp-
nece est.

Flores melliferi. Semina oleifera.

Bonum pecoribus praebet alimentum.

TETRADYN. SILIQUOSA.

103

1032. *RAPHANUS sativus* filiquis tere-
tibus torosis bilocularibus. *Linn. Sp. pl.*
2. p. 935.

Raphanus minor oblongus. Bauh. pin. 96.

Raphanus major orbicularis f. rotundus.
Bauh. pin. 96.

2. *Raphanus niger. Bauh. pin. 96.*

3. *Raphanus chinensis annuus oleiferus.*

Icon. Blackw. t. 87.

Germ. Rettig.

Gall. Radis.

Hab. In China. Colitur utrinque in hortis sub
multis variationibus.

Flor. Julio, Augusto. Semina Augusto matu-
ra. ☉ ♂

Pharm. RAPHANI Radix recens.

Vires antiscorbuticas diureticas incitantes habet.

Comeduntur radices recentes crudae, cum sale cu-
um. In Var. 3. Semina maxime oleosa sunt.

Flores apibus melliferi.

CLASSIS XVI.
MONADELPHI.

PENTANDRIA.

* *Geranium cicutarium, pimpinellaefolium
moschatum.*

DECANDRIA.

CCCXXVI. GERANIUM. Monogyna. Cal.
5-coeca, rostrata.

* *Flores papilionacei monadelphii: Sparti
Genista. Lupinus. Anthyllis. Ononis.*

POLYANDRIA.

CCCXXVIII. MALVA. Polygyna. Cal. e
rior 3-phyllus. Arilli mono-
spermi, verticillati, plur

CCCXXVII. ALTHÆA. Polygyna. Cal. e
rior 9-fidus. Arilli monos-
permi, verticillati.

CLASSIS XVI.
MONADELPHIA.
DECANDRIA.

CCCXXVI GERANIUM.

Linn. Gen. pl. ed. Schreb. n. 1118.

Calyx monophyllus, quinquepartitus, f. pentaphyllus: *foliis* ovatis, acutis, concavis, persistentibus. *Corolla Petala* quinque-lobata f. ovata, patentia. *Germen* quinquelobatum, rostratum. *Stylus* subulatus, staminibus longior, persistens. *Stigmata* quinque, reflexa, *Capsula* pentacocca, rostrata, loculis interne persistentibus, singulis affixis caudae f. aristae terminali, longae, demum spirali. *Semina* solitaria, raro bina, ovato-oblonga.

† *Staminibus quinque antheriferis.*

1053. GERANIUM *cicutarium* pedunculis multifloris, floribus pentandris, foliis pinnatis incisis obtusis, caule ramoso. *Linn. Syst. veg. p. 615.* — *Pollich. fl. pal. n. 646.*

Geranium chaerophyllum caule erecto-decumbente, foliis oppositis pinnatis, incisis, pedunculis elongatis multifloris, cotyledonibus trilobis. *Cavan. Diff. 4. p. 226. n. 319. Roth. fl. germ. 2. p. 136.*

208 MONADELPHIA. DECANDRIA.

Erodium cicutarium pedunculis multifloris, foliis pinnatis, pinnis pinnatifidis incisos, corollis calyce majoribus, caule hirsuto, strato. *Linn. Sp. pl. ed. Willden. 3. p. 1.*

Erodium pedunculis multifloris, foliis pinnatis, foliolis sessilibus pinnatifidis incis. *Smith. brit. 2. p. 727.*

Geranium caule procumbente, petiolis multifloris, foliis duplicato-pinnatis, pinnis acute incis. *Hall. stirp. helv. n. 94.*

Geranium Cicutæ folio minus et supinum. *Bauh. pin. 319. Dill. Cat. Giff. p. 48. Mapp. fl. Als. 126.*

Geranium arvense. *Tabernæmont. p. 1. Icon.*

Icon. Cavanill. l. c. t. 95. fig. 1. — Cuv. fl. Lond. — Oeder. fl. Dan. t. 986.

Germ. Schierlingblättriger Storchschnabel. Gall. Bec de grue cicutin.

Hab. Utrinque in arvis, agris, vineis, passim fabulosis apricis sterilibus, sepibus paucis frequens.

Flor. Martio, Aprili, Majo, Junio. *Sem.* Majo, Octobri matura. ☉

Radix fusiformis, albida. *Caulis* digitatus, palmaris, pedalis et altior, teres, plus minus villosus, scabriusculus, geniculis incrassatis instructus, erecto-decumbens, ramosus. *Rami* alterni, diffusi. *Folia* radicalia numerosa, longius petiolata, caulina opposita, pinnata, pinnis oppositis alternisque, subsessilibus, profunde incis, acutis, villosiusculis, læte viridibus. *Stipulae* ovato-acuminatæ, marcescentes. *Flores* umbellati. *Pedunculi* communes exserti, alterni, longi, erecti, solitarii, villosi.

MONADELPHIA. DECANDRIA. 109

umbellatum 3-5-6-7 flori. *Radix* floriferi
ti, fructiferi deflexi. *Involucrum* subsep-
tatum, ad radiorum basin, acutum, revo-
m. *Calyx* profunde quinquepartitus, *laciniis*
imbo-ovatis, seta terminatis, patentibus,
cristatis, margine albidis, aut fusco-ru-
bis. *Corolla Petala* quinque, calyce subma-
ior, ovalia, obtusa, integra, unguibus pilo-
sis, violaceo-purpurea, rubella, aut albida,
purpureis picta. *Stamina Filamenta* de-
candria purpurea, alterna breviora, antheris de-
candria. *Antherae* quinque, violaceo-purpureae.
Stigma Arista barbata.

riat. Corollis purpureo-violaceis, rubel-
lis, roseis et albis.

L. Primo vere et autumnio saepe in arvis agrisque
et sepibus inveniuntur plantulae hujus speciei te-
nues, foliis solummodo radicalibus, et scapis ra-
dicalibus umbelliferis, floriferis instructae.

ota. Planta in pharmaciis non confundenda, cum
Fumaria officinali, ut ante aliquot annos
vidi.

Flores apibus melliferi.

In sabulosis arenam figit. Bonum ovibus pabulum.

34. GERANIUM *pimpinellæfolium*

pedunculis multifloris, floribus pentan-
dria, foliis pinnatis, pinnulis subovatis
subsessilibus, incilis, corollis calyce sub-
minoribus, caule ramoso, decumbente.

Erodium pimpinellifolium pedunculis mul-
tifloris, foliis pinnatis ovatis, sessilibus
inciso-dentatis, corollis calyce minori-
bus, aristis glabris, caule decumbente
subpiloso. *Linn. ed. Willden. 3. p. 630.*

Geranium cicutarium caule ramoso prostra-
to, foliis pinnatis, pinnulis subovatis.

110 MONADELPHIA: DECANDRIA.

Gavanill. Diff. p. 226. n. 318. — R. fl. germ. 2. p. 135.

Geranium pimpinellæ folio. Dill. Cat. Giff. app. 69. — Mapp. fl. Alf. p. 124

Geranium IV. et V. Tabernæmont. p. 1 Icon.

Icon. Cavanill. l. c. t. 93. fig. 1.

Germ. Pimpinellenblättriger Storchschnabel

Gall. Bec de grue à sévilles de Pimperne

Hab. et Flor. passim cum præcedente in vis et cultis. ☉

Maxime affine præcedenti, ita ut *Linna* utrinque combinaverit. Differt tamen, *Cotyledonibus* cordatis indivisis, nec trilobis, *Caubus* magis ad terram decumbentibus, *Foliis* pinnis basi latioribus, *Corollis* calyce submissis, aut aequalibus, *Seminum Aristis* glabris.

1035. GERANIUM moschatum pediculis multifloris, floribus pentandris, liis pinnatis incisis, cotyledonibus pinnatifidis. *Linn. Sp. pl. 2. p. 951.*

Geranium procumbens, foliis pinnatis, pinnulis ovatis incisis, cotyledonibus pinnatifidis, floribus umbellatis. *Cavanill. Diff. 4. p. 227. n. 320.*

Erodium moschatum pedunculis multifloris, foliis pinnatis, foliolis subpetiolatis inqualiter incisis. *Aiton. hort. Kew. 2. 414. Linn. ed. Willden. 3. p. 631.*

Geranium petiolis multifloris, caule diffusis, foliis ovatis pinnatifidis. *Hall. stirp. he n. 945.*

Geranium cicutæfolio moschatum. Bauh. p. 319.

Jacqu. hort. Vind. t. 55. Blackw. t. 150.

Cavanill. l. c. tab. 94. fig. 1.

Bisam-Störchschnabel.

Bec à grue musqué.

In regione Basileensi, *Augusta t. Stæhe-*
o. Prope Muttentz t. Joh. *Bauhino*. An
ontanea? forsan ex hortis ubi passim co-
ar emigravit.

Junio, Julio. Semina Augusto, Sep-
tembri matura. ☉

radices ex radice plures, pedales, fesqui-
met longiores, decumbentes, villosi,

Folia opposita, petiolata, pinnata. *Pin-*
nae alternæ, ovatae, inciso-serratae. **Flores**
in raculis longis universalibus, axillaribus,
et. **Corollæ** violaceo-purpureascentes,
intus albidæ, vix calyce longiores. **Sta-**
mentum decem, quinque alterna an-
testituta.

Haec planta, præsertim folia moschum redolent.

. **ACUS MUSCATA** f. **MOSCHATAE** Her-
recens.

Ubi in medicina exoleta, passim ad coediendos
æsertim Pastinacam in culinis adhibetur. Conf.
Harsa Semplici p. 139.

minibus decem fertilibus. Pedunculis bi-
floris a. perennia.

1. **GERANIUM phæum** pedunculis
bifloris, foliisque alternis, calycibus sub-
aristatis, caule erecto, petalis undulatis.
Linn. Sp. pl. 2. p. 953.

Geranium phæum caule herbaceo, foliis al-
ternis hirsutis palmato-quinquelobis den-

724 MONADELPHIA. DECANDRIA.

tatis incis, floribus subspicatis, ca-
bus mucronatis. *Cavanill. Diss. II*
210. n. 292.

Geranium foliis rugosis hirsutis semise-
bis, lobis dentatis, petalis ferratis,
minatis. *Hall. stirp. helv. n. 934.*

Geranium phaeum f. fuscum petalis re-
planis, folio maculato, et non ma-
lato. *Mapp. fl. Alf. p. 125.*

Geranium phaeum f. pullo flore *Clusii.*
Bauh. 3. p. 477. Icon.

β. Petalis reflexis.

Geranium fuscum pedunculis bifloris op-
positifoliis geminis, caule patulo, pe-
integerrimis. *Linn. Mant. 97.*

Geranium phaeum f. fuscum, petalis re-
Morif. hist. 2. p. 515. Mapp. fl. A
125. (folio maculato et non macula-
to.)

Icon. Cavanill. l. c. tab. 89. fig. 2.

Germ. Brauner Storchschnabel.

Gall. Bec a grue brun.

Hab. In Margraviatu superiore retro *Ba-*
wier ad metallifodinam *Hausbaden*,
non in monte prope *Crenzach*. In Alfa-
pratis montosis retro *Barr*, cum vari-
abunde. *t. Mappo.*

Flor. Junio, Julio. *Semina* Augusto matur-

Caulis sesquipedalis, cubitalis et alti-
erectus, villosus, nodoso-articulatus, rami
Folia alterna, molliter villosa, subrugosa, i-
riora longe petiolata, superiora subsessilia,
peltata, quinqueloba, lobis inaequaliter in-
dentatis. *Stipulae* lanceolatae, flaccidae.
Res in pedunculo universali longo, oppositiflo-
bifloro. *Calyx* profunde quinquepartitus,
me

mentis ovatis, aristatis. *Corollae* petala quinque
sepiissime reflexa, subrotunda, acuta ore undula-
ta, fusco-rubentia.

Planta elegans ad rariores Florae cives pertinet.

Nota. *Geranium fuscum*, reflexum, et livi-
dum Aiton. h. Kew. f. patulum Vill. fl.
Delph. 3. p. 371. forsan merae sunt variatio-
nes *Geranii phaei* a diverso solo natali depen-
dentes.

036. *Geranium reflexum* in Hispania passim cum
sore et seminibus maturis lectum, in horto Carls-
ruhani cultum, petala reflexa laciniata, transmuta-
vit in integra. In loco natali saepe in uno eo-
demque individuo Calyx subaristatus, aristatus,
et muticus, Petala plus minusve laciniata et in-
tegra observantur.

1037. *GERANIUM sylvaticum* pedun-
culis bisfloris, foliis subpeltatis quinque-
lobis inciso-ferratis, caule erecto, pe-
talis emarginatis. *Linn. Sp. pl.* 2. p. 955.

Geranium batrachioides canle herbaceo
erecto dichotomo, foliis oppositis quin-
quepartitis, incis: floribus coeruleis
magnis, calycibus inflatis. *Cavanill. Diff.*
4. p. 211. n. 295.

Geranium caule erecto, foliis rugosis septi-
lobis, lobis acute ferratis, petalis emar-
ginatis. *Hall. stirp. helv. n.* 932.

Geranium batrachioides folio *Aconiti*. *Bauh.*
pin. 317.

Icon. *Cavanill. l. c. tab.* 85. *fig.* 2.

Germ. *Wald-Storchschnabel.*

Gall. *Bec-de Grue des bois.*

Heb. In *Brisgoviae monte Feldberg*. In *Marg-*
graviatus superioris monte Belchen infre-
quens. In *Ditione Basileensi circa Langen-*
bruck et passim alibi in umbrosis sylvis al-

214 MONADELPHIA. DECANDRIA.

tiorum Juræ montium circa *Wasserfa*
nec non in monte *Ballon* Alsatiæ.

Flor. Majo, Junio. *Semina* Julio matura.

Caulēs ex radice plures, cubitales et alti
res, erecti, subtetragoni, crassiusculi, inferi
glabri, nudi, superne villosi, dichotomi, fi
liosi. *Folia* radicalia longe petiolata, numero
subrotunda, rugosa, subhirsuta, ultra mediū
septiloba: *lobis* latis inciso-serratis; *cauli*
opposita, petiolata, radicalibus analoga; *rami*
subsessilia, triloba, serrata, caulinis minor
Stipulae binae, lineari-subulatæ. *Flores* axill
res terminalesque, pedunculati. *Pedunculi* su
biflori, terminales 2-3-4, subsessiliati. *Cal
ces* aristati, hirsuti. *Corollæ* speciosæ, rubr
coeruleae. *Petala* quinque, obovata, plus m
nusve, quandoque vix emarginata, patentib
venis quinque ad septem aqueis picta.

Variat in hortis præcipue Corollis albis, venis pu
pureis pictis.

Distinguitur a *G. pratensi*, Caulibus crassior
bus inferne glabris nec villosulis, Foliis minus pro
funde septempartitis, laciniis multo latioribus, no
trilobis subtrifidis, sed inciso-serratis, Corollis ri
bro coeruleis, nec coeruleis, Petalis obovatis, mi
nus latis, sæpissime emarginatis, nec semper integri

Differt a *G. palustri*, Caulibus inferne gl
bris, nec utrinque pilosis, Foliis minus profunde
septemlobis.

Nota. *Geranium batrachyoides*. Cavanil
l. c. differt aliquatenus ab individuis nostratibus
Caulibus demissioribus, Corollarum petalis spe
ciosioribus integris.

Verum hortorum præsertim anglicorum decus, cui
G. pratensi et *palustri*.

Flores apibus melliferi.

1038. GERANIUM palustre pedunculis
bifloris longissimis declinatis, foliis quin-
quelobis incisis, petalis integris. *Linn.*
Sp. pl. 2. p. 954. Pollich. fl. pal. n. 647.

Geranium palustre caule herbaceo villosa:
foliis oppositis villosis quinquelobis in-
cisis, pedunculis longissimis declinatis.
Cavanill. l. c. p. 211. n. 294.

Geranium caule diffuso, foliis rugosis semi-
quinquelobis, lobulis trifidis, floribus
spatis, petalis integerrimis. *Hall. stirp.*
Helv. n. 933.

Geranium Batrachioides maximum minus
laciniatum, folio Aconiti. *Bauh. hist. 3.*
p. 477. — Mapp. fl. Alf. p. 125.

Geranium Batrachioides palustre flore san-
guineo. *Dill. hort. Eltham. 160. t. 134.*
f. 161.

Icon. Oeder. fl. Dan. t. 596. Cavanill. l. c. t.
t. f. 2.

Germ. Sumpffstorchschnabel.

Gall. Bec de-grue des marais.

Hab. Prope Pforzheim in prato sylvatico ad
fluvium *Nagold*. In Ditione Basileensi prope
Langenbruck ad viam circa *Arisdorf* in fos-
sis. In Alsatia monte *Ballon t. Joh. Bau-*
hino. In Palatinatus locis humidiusculis ad
pratorum margines juxta alnos ac dumeta
inter *Neckargemünd, Wemersbach, Mos-*
bach t. Pollichio.

Flor. Junio, Julio. Semina Julio, Augusto
matura. ¶

Caules ex radice plures, cubitales et altio-
res, adscendentes, obliqui, quandoque erecti-

116 MONADELPHIA, DECANDRIA.

usculi, obsolete tetragoni, incrassato-gemmati, subdichotomi, utrinque hirsuti, pilis densis spectantibus. *Rami* divaricati, hirsuti. *Fol.* radicalia numerosa, longe petiolata, subrotunda, villosa, ultra medium leptiloba, lobis trifidis, obtusis. *Petioles* teretiusculi, hirsuti. *Caulina* opposita, petiolata, profunde quinque palmata, segmentis incisivo-dentato-serratis, acutis. *Rameae* sessilia, opposita, caulinis minora, angustiora. *Stipulae* utrinque binae, ovato-acuminatae, bifidae, margine membranaceae, seminariae longae, erectae, glabrae. *Flores* axillares terminalesque, longe pedunculati, biflori, nutantes. *Bractae* subpedicellis, quatuor, angustae, lineari-acutae. *Calycis foliola* trifida, angustula, lineari-lanceolata, aristata. *Corolla* tubulosa, campanulato-patens, coeruleo-violacea. *Petala* quinque, obovata, integra, quandoque submarginata, striis quinque aqueis percul-

O b f. Tota planta, corolla excepta, hirsuta.

Flores apibus mel praebent.

1039. *GERANIUM pratense* pedunculis bifloris, foliis subpeltatis multipartitis pinnato-laciniatis rugosis acutis, talis integris. *Linn. Sp. pl. 2. p. 5*
Pellich. fl. pal. n. 648.

Geranium pratense caule herbaceo erecto foliis oppositis, subpeltatis multipartitis rugosis; lobis pinnatis acutis. *Cavendish. l. c. p. 210. n. 293.*

Geranium caule erecto, foliis rugosis hirsutis multilobis, lobis trifidis, lobulis bipinnatis, floribus umbellatis. *H. stirp. helv. n. 931.*

MONADELPHIA. DEÇANDRIA. 117

Geranium batrachioides, Gratia dei germanorum. *Bauh. pin.* 318. — *Mapp. fl. Als.* p. 125.

Leon. Curtis fl. Lond. t. 220. *Cavanill. l.* 6. t. 87. f. 1.

Ger. Wiesenstorchschnabel.

Gall. Bec-de-grue des-prés.

Hab. Retro Ettlingen, auf dem Kaltenbrunnen, auf der Herrnwiese, im Buhlerthal, Oppenauerthal in pratis montolis riguis dumetosis. In Margraviatu superiore circa Sirnitz et in monte prope Crentzach. In Ditione Basileensi in pratis circa Gempen et Stehelino. In Alsatie montolis pratis udis prope Liechtenberg, in monte versus dem Steinthal copiose, nec non auf dem Hochgebürg Hochfeld t. Mappo. In Palatinatu ad vias humidiusculas dumetosas passim inter Franckenstein, dem Bicard et Hartenburg; circa Heidelberg, Mosbach in pratis et dumetis copiose post Pollichium vidi.

Flor. Junio, Julio, Augusto. Semina Augusto, Septembri maturat. 7

Caules ex radice plures, cubitales et altiores, erecti, angulati, subpubescentes, ramosi. *Folia* radicalia-caulinaque inferiora, longe petiolata, superiora rameaque subseffilia, subrotunda, subpeltata, superne viridia, subtus villosa-canescientia, venosa, profunde ad basin sere multipartita: *laciniis* trilobato-pinnatifidis, acutis. *Stipulae* utrinque binae, acutae, fuscorubellae, marcescentes. *Flores* axillares terminalesque, pedunculati, biflori. *Bractee* ad basim pedicellorum, quatuor, lineari-acutae, fuscae. *Calycis* foliola subquinquecostata, acuta, aristata, villosula. *Corollae* speciosae, amoene

118 MONADELPHIA. DECANDRIA.

coeruleae, quandoque albae. *Petala* subrotundo-ovata, obtusa, integra; striis aqueis cta. *Antherae* oblongae, coeruleae.

Variat praesertim in hortis, Corollis albidis.

Egregium hortorum decus.

Flores melliferi.

1040. GERANIUM *pyrenaicum* ca

erecto ramoso, pedunculis bifloris, foliis radicalibus caulisque petiolatis, lacinjiis ad medium septilobis, lobis bifidis obtusis; rameis quinquelobis, lobis trifidis acutiusculis; supremis profunde tripartitis, laciniis acutis integris, petalis bifidis calyce duplo longioribus.

Geranium caule herbaceo villosa, foliis oppositis subpeltatis orbiculatis, lobis trifidis obtusis, corollis purpureo-violaceis. *Cavan. l. c. p. 203.*

Geranium pyrenaicum. Linn. Mant. 97.

Geranium pedunculis bifloris, foliis multifidis, laciniis obtusis inaequalibus, petalis bifidis. *Gerard. fl. Gallopr. p. 434. fig. 16. n. 2.*

Geranium columbinum perenne, *pyrenaicum*, maximum. *Tournef. inst. 268.*

Germ. Pyrenaeischer Storchschnabel.

Gall. Bec-de-grue des Pyrenées.

Hab. In Principatu superiore in locis graminis dumetosis ad Lacum Bodamicum prope Moersburg ubi mihi obvenit *An. 1801* passim non infrequens.

Flor. Majo. Semina Junio, Julio matura.

Caules ex radice plures, sesquipediales cubitales et altiores; erecti, quandoque ad

MONADELPHIA. DECANDRIA. 119

inferiores, inferne obtuse angulati, superne
 minusculi, villosi. *Folia* radicalia caulinaque
 inferiora longe petiolata, villosa, rotundata,
 medium circiter septiloba, lobis trifidis, ob-
 tusis; superiora rameaque breviter petiolata,
 unguiculata, lobis trifidis acutiusculis; superiora
 pedunculis opposita, subsessilia, multo mi-
 nora, profunde tripartita, lobis trifidis; superio-
 ra sessilia, profunde tripartita, lobis ovato-
 angulatis, acutis, integris. *Stipulae* utrinque
 lineares, bifidae vel trifidae, fuscorubentes. *Flores*
 illae terminalesque, pedunculati. *Pedun-*
culi fructiferi patentissimi, biflori. *Pedicelli*
ciliales, infracti, basi *Bracteis* quatuor, angu-
 lis, linearibus, rubellis, villosis, suffulti. *Ca-*
ulis foliola quinque, lanceolata, subtrico-
 ta, villosula, apice mutica. *Corollae* viola-
 coeruleae. *Petala* calyce duplo longiora,
 illa, supra basin acutam, utrinque villorum
 versalium serie e latere emergentium in-
 vecta, venis saturationibus picta. *Filamenta*
 decem, fertilia. *Antherae* coeruleae.

Nota. Planta omnino ad rariores patriae cives per-
 tinet, hucusque in Germania vix detecta.

Individua ad Lacum Bodanicum lecta, simillima
 uno illis, quæ ex pascuis Pyrenaeorum alpinis nec
 e Gallia meridionali collecta reservo.

Nota. Filamenta decem in omnibus hucusque tam
 spontaneis quam in horto cultis individuis, An-
 theris perfectis instructa fuerunt. Idem obser-
 vavit Ill. Cavanilles. Hinc differentia speci-
 fica à b. Linnaeo sine dubio ex auctoritate
 Cl. Gerardi data, emendanda erat.

¶ In planta fere deflorata, saepe Antherae pluri-
 mae, ut facile sit etiam in aliis speciebus, deper-
 ditae sunt.

¶ *Staminibus decem fertilibus. Pedunculis*
bifloris. β. annua.

1041. GERANIUM *robertianum*

pedunculis bifloris, foliis ternatis pinnatis, incisilis, calycibus pilosis decemangulatis.

Geranium pedunculis bifloris, calycibus pilosis decemangulatis. *Linn. Sp. pl.* 2. 955. — *Pollich. fl. pal.* n. 649.

Geranium *robertianum* foliis ternatis, pinnatis, incisilis calycibus rubris decemangulatis, pilosis. *Cavanill. p.* 215. n. 345.

Geranium foliis duplicato-pinnatis, pinnatis ultimis confluentibus, calycibus strigosis hirsutis. *Hall. stirp. helv.* n. 943.

Geranium *Robertianum* rubens et viride. *Bauh. pin.* 319. *Mapp. fl. Alf. p.* 124.

Geranium. *Tabernæmont. p.* 132. *Icon.*

♂. Corollis albis.

Geranium *Robertianum* flore albo. *Mapp. fl.*

Icon. Curtis fl. Lond. Bulliard herb. t. 201. *Oed. fl. Dan. t.* 694. — *Cavanill. l.c. t.* 86. f. 1. *Blackw herb. t.* 480.

Germ. Rubrechtskraut. Bocks-Storchschnabel. Gall. Bec-de-grue Robertin.

Hab. Circa Carlsruhe et alibi, in nemoribus dumetis, et muris passim abunde.

Flor. Majo, Junio, Julio et sequ. *Semin.* Julio Augusto matura. ☉

Caulis pedalis sesquipedalis et altior, erectus, teres, incrassato-geniculatus, subdichotomus, pilis raris praeditus, saepe fusco-purpureus. *Folia* radicalia caulinaque petiolata rareriter pilosa, tripartita, pinnulis inciso pinnatifidis. *Stipulae* acutiusculæ, villosæ, parvæ, apertae. *Petiolorum* basin. *Flores* axillares terminalesque longe pedunculati. *Pedunculi* teretes, subvillosi.

MONADELPHIA. DECANDRIA. 121

his, biflori. *Pedicelli* ad basin *Bracteis* minutis
 fuscopurpureis, præditi. *Calycina* foliis quin-
 que, sæpe fusco-purpurea, longe pilis, ari-
 tata, alterna tricolata. *Corollæ* petala quinque,
 ovata, obtusa, integrifolia, rubella, lineis
 tribus albidis picta, quandoque albida, calyce
 longiora.

Pærm. GERANII ROBERTIANI Herba.

Pasta foetens, austera, adstringens, vulneraria,
 hæc medicina hodie fere exolevit.

Conices Herba recens cocta, usque et admota pellit. In
 temeraria pecorum, infusum propinant pastores. Linn.
 Saec. p. 241.

42. GERANIUM *lucidum* L. foliis
 bifloris, calycibus transversim elevato-
 rugosis, lobis rotundatis.

Geranium lucidum p. bifloris, ca-
 lycibus pyramidalibus, lobatis, elevato-
 rugosis, foliis quinquelobis, rotundatis.
 Linn. Sp. pl. 2. p. 955. Michx. fl. pal.
 n. 650.

Geranium caule herbaceo foliis oppositis pi-
 losis lucidis rotundato-lobatis, calyci-
 bus pyramidalibus transversim rugosis. Ca-
 van. l. c. p. 214. n. 299.

Geranium foliis subrotundis semiquinquelobis,
 lobis obtusis, calycibus transversim
 rugosis. Hall. stirp. helv. n. 942.

Geranium lucidum saxatile. Bauh. pin. 318.
 Cavanill. l. c. t. 80. f. 2. — Oeder. fl.
 Dan. t. 218.

Ger. Glänzender Storchschnabel.

ll. Bec a grue luisante.

224 MONADELPHIA. DECANDELLA.

Hab. In Palatinatu circa Steinbach in sylvis
rupes umbrosas, t. Pollichio.

Flor. Majo, Junio. *Semina* Julio maturantur.

Caules ex radice plures, pedales, scilicet
pedales, adscendentes, erectiusculi, teres
amoene rubelli, glabri, incrassato-geniculi
ramosi. *Folia* caulina opposita petiolata, tri-
da, subrotunda, 5-7 lobata, lobis subtri-
dentato-crenulatis, rariter pilosis. *Stipulae*
ad basin petiolorum utrinque binæ, parvæ,
triangulæ, acutæ, rubentes. *Flores* alternatim axillares
terminalesque, pedunculati. *Pedunculi* bifi-
dissimi. *Pedicelli* basi *Bractea* triphylla minuta, angustata
rubella suffulti. *Calyces* pyramidato-quinque-
guli, subaristati, transversim sulcis quatuor-
decim longitudinalibus præditi, virides. *Corolla* parva, calice
longior. *Petala* integra, purpureo-dilute colorata.

Planta elegans, omnino ad rariores Floræ clas-
sem pertinet, jam ex longinquo ob caules splendentes
bellos conspicua.

1043. *GERANIUM molle* caulibus erectis
longe pilosis, foliis orbiculatis 5-7-
lobis, trifidis obtusis, pilosis, caulinis
oppositis, rameis superioribus alternis, 1-
pedunculis bifloris, calycibus pilosis mu-
ticis, petalis bifidis calycem superan-
tibus.

Geranium molle pedunculis bifloris foliis
que floralibus alternis, caule ramoso ere-
ctiusculo, calycibus muticis, arillis liliatis
vibis. *Linn. Sp. pl.* 2. p. 955. *Poll.*
pal. n. 651.

Geranium molle caule suberecto: foliis
orbiculatis subseptemlobis trifidis obtu-
satis caulinis oppositis, superioribus alter-
nantibus. *Cavanill. l. c. p.* 203. n. 281.

MONADELPHIA. DECANDRIA. 129

Geranium foliis mollissimis hirsutis reniformibus semiquinquesidis, lobis semitripartitis obtusis. *Hall. stirp. helv. n. 939.*

Geranium columbinum villosum, petalis bifidis purpureis. *Vaill. bot. parisi. 79.*

Icon. Curtis fl. Lond. t. 141. Cavanill. l. c. t. 13. f. 3. Vaillant. l. c. t. 15. f. 3. Oeder. 3. Dan. t. 679.

Ger. Weicher Storchschnabel.

Gall. Bec-de-grue mollet.

Hab. Utrunque passim ad vias parietes, et sepe frequens. Copiose circa *Carlsruhe.*

Flor. Majo, Junio, Julio. *Semina* Julio, Augusto matura. ☉

Caulis ex radice saepe plures, pedales, sesquipedales et altiores, teretes, versus basin rubri, caeterum virides, utrinque longius et tenerrime pilosi, molles. *Folia* caulina opposita, superiora alterna, petiolata, pilosa, mollia, orbiculata, 5-7 loba, lobis trifidis, integerrimis, obtusis; suprema breviter petiolata, alterna, profunde quinquepartita: laciniis lanceolatis, integris bifidisque. *Stipulae* utrinque line. membranaceae, subbifidae, pilosae, marcescentes. *Flores* axillares, terminalesque, pedunculati. *Pedunculi* pilosi, biflori. *Pedicelli* bini. *Bractea* quadriphylla, parva, membranacea, suffulti. *Calyx* quinquephyllus. *Foliola* lanceolata, longe pilosa, mollia, mutica, viridia, corolla breviora. *Corollae* petala coerulea, aut rubello-violacea, bifida. *Antherae* decem, violaceo-coeruleae.

Nota. Facile hanc ab aliis speciebus affinis, Caulis semper longe et tenerrime pilosi non viscidis distinguunt.

1044. *GERANIUM dissectum* caule
erectiusculo villosulo, foliis quinquepart
trifidis, pedunculis bifloris folio brev
ribus, calyce aristato, petalis em
natis calyce subbrevioribus.

Geranium dissectum pedunculis bifloris,
liis quinquepartito-trifidis, petalis en
ginatis longitudine calycis. *Lin. Sp.*
2. p. 956. — *Pollich. fl. pal. n. 652.*

Geranium dissectum caule villosulo erecto
culo, foliis quinquepartito-trifidis;
dunculis bifloris, folio brevioribus.
vanill. l. c. p. 199. n. 276.

Geranium caule erecto, foliis quinquel
lobis trifidis acutis. *Hall. stirp. helv.*
937.

Geranium majus, foliis imis longis ad m
pediculum divis. *Vaill. bot. parij*

Icon. Cavanill. l. c. t. 78. f. 2. — Oeder
Dan. t. 936. Vaill. l. c. t. 15. f. 2.

Ger. m. Zerschlißener Storchschnabel.

Gall. Bec de grue dissequé.

Hab. Utrunque inter stipulas in agris, ad
sepes passim frequens.

Flor. Junio, Julio. *Semina* Augusto matur

Distinguitur a *G. columbino* affini, *Ca*
bus saepe erectiusculis, villosulis, *Foliis*
linis quinquepartitis, *Laciniis* trifidis, r
bifidis, quandoque indivisis, nec quinquel
Pedunculis folio brevioribus, nec longiori
Corollis parvis, calyce subbrevioribus,
bellis, sanguineis, bifidis, nec dilute co
leis, integriusculis.

5. *GERANIUM columbinum* caule

procumbente foliis quinquepartito-multifidis, pedunculis bifloris folio longioribus, calycibus aristatis, petalis oblongis subemarginatis, longitudine calycis.

Geranium columbinum pedunculis bifloris folio longioribus, foliis quinquepartito-multifidis: laciniis acutis, arillis glabris, calycibus aristatis. *Linn. Sp. pl. 2. p. 956. Poll. fl. pal. n. 653.*

Geranium columbinum caule procumbente: foliis oppositis palmatis, laciniis pinnatis: pedunculis solitariis longissimis bifloris. *Cavanill. l. c. p. 200. n. 277.*

Geranium caule procumbente foliis quinquepartitis, lobis trilobis, lobulis trifidis. *Hall. stirp. helv. n. 938.*

Geranium columbinum foliis dissectis, pediculis florum longissimis. *Vaillant. bot. par. p. 79. Mapp. fl. Alf. p. 124.*

Ger. Cavanill. l. c. t. 82. f. 1. Vaillant. l. c. t. 15. f. 4.

erm. Tauben-Storchschnabel.

Hall. Bec-de-Grue Columbin.

Ger. Utrisque ad vias et sepes passim.

Flor. Junio, Julio. Semina Julio, Augusto maturat. ☉

Caules ex radice plures, palmares pedales, longiores, sæpissime procumbentes, teretes, subullosi, articulati, viridi-rubentes, quandoque fusco-rubelli, ramosi. *Folia* caulina opposita, petiolis breviora, orbiculata, proinde septem aut quinquelaciniata: *Laciniis* pinnatis, linearibus, utrinque hispidulis. *Stipulae* subulatæ, subtrifidæ, quandoque bifidæ,

126 MONADELPHIA. DECANTRI.

fulco-rubellæ. Flores axillares, termina pedunculati. Pedunculi solitarii, biflor longiores. Pedicelli basi Bracteis quatuor latis, rubellis, suffulti. Calyx pentagonus aristatus, villosus. Corollæ petala quatuor oblonga, obtusa, subemarginata, coeruleo purpureis picta, calycis fere longitudinem æquant.

In arenosis cum præcedente specie, figit.

1046. GERANIUM *malvæfolium*.

libus erectiusculis, brevissime mollibus, foliis petiolatis, orbiculatis ultra medium septilobis, trifidis, villosulis, mollibus, pedunculis floribus, calyce mutico piloso, emarginatis calycem vix superantibus filamentis decem, alternis casis.

Geranium malvæfolium. Scopol. fl. 2. p. 37. n. 847. Roth. fl. germ. 2.

Geranium rotundifolium. Pollich. fl. 654.

Geranium pusillum caule herbaceo r. foliis subrotundo lobatis; lobis trifidis floribus minimis pentandris coarctatis. *Cavan. l. c. p. 202.*

Geranium pusillum. Willdenow. Lin. pl. 3. p. 713.

Geranium foliis hirsutis semiseptilobis semitrilobis obtusis. Hall. stirp. 1. 940.

Geranium columbinum majus flore r. coeruleo. Vaillant. bot. parif. p. 124. Mapp. fl. Als. p. 124.

β. Caulibus procumbentibus.

MONADELPHIA. DECANDRIA. 137

Frax. 4. *Vaillant l. c. t. 15. f. 1. optima.*
Cavanill. l. c. t. 83. f. 1.

Ger. Malvenblättriger Storchschnabel.

Gall. Bec-de-grue mauvin.

Hel. Utinque ad vias sepes aggeres passim.
 Copiose circa Carlsruhe in dumetis im
felsenengarten et alibi.

Flor. Majo, Junio. *Semina* Junio, Julio ma-
 tura. ☉

Radix subsufiformis, crassa, albida. *Cau-*
lis ex radice copiosi, pedales, cu-
 linae et altiores, erecti, teretes, brevissime
 articuli, molles, incrassati, articuli, basi ru-
 bellis, ceterum saepe virides, ramosi. *Rami* al-
 terni, erecti. *Folia* villosula, mollia, radica-
 la numerosa, longe petiolata, orbiculata, ad
 medium 7-9 lobata: lobis trifidis obtusis: lo-
 bus medio tricenato, obtusissimo; *caulina*
mentis opposita, petiolata, radicalibus limi-
 ta; *superiora* alterna, profundius quinque-
 lobis, trifida: *laciniis* angustioribus acutiuscu-
 lis. *Stipulae* binae, acuminatae, bifidae, ru-
 bellae, margine pilosulae. *Flores* axillares,
 terminalesque pedunculati. *Pedunculi* unciales,
 semunciales, biflori. *Pedicelli* incrassati semi-
 unciam longi, *Bracteis* quatuor setaceis, suf-
 fultis. *Calycina* foliola quinque, ovata, acuta
 pilosa. *Corollae* petala quinque, coe-
 rulea, emarginata, vix calycem superantia.
Filamenta decem, alterna castrata, *Antherae*
 purpureae, violaceo-coeruleae.

Variatio β. Caulibus procumbentibus palmari-
 bus pedali- sesquipedalibus vix altiori-
 bus magis rubellis promiscue cum α. crescit.

Nota. Individua hujus bene multa sedulo examinata,
 etiam in horto Botanico culta, semper Filamentis
 decem, et alternis quidem castratis instructa.

Geranium pusillum Linn. nullibi a vanillide observatum, hucusque frustra Gallia, Helvetia, Hispania et Germania quam forsan planta dubia ex Catalogo plantarum movenda? Sic mihi certe videtur.

1047. *GERANIUM rotundifolium* c

le prostrato divaricato, pubescente, cido, foliis oppositis longe petiolo inferioribus orbiculatis linuato quing lobis crenatis, superioribus subcune trilobis, pubescenti-viscidis; pedun lis bifloris, calyce villoso aristato, p lis calyce sublongioribus, integris.

Geranium rotundifolium pedunculis biflo petalis integris obtusissimis longitud calycis, caule prostrato, foliis reni mibus incis. *Linn Sp. pl.* 2. p. 957.

Geranium rotundifolium caule prostra foliis oppositis, inferioribus subrotu semiquinquepartitis, superioribus f cuneatis; petalis integerrimis, cal paulo longioribus. *Cavanill. l. c. p. 2 n. 300.*

Geranium viscidum caule decumbente, liis semiquinquelobis tridentatis et qu quelobatis. *Hall. stirp. helv. n. 941.*

Geranium folio malvae rotundo. Bauh. 2 318. — Mapp. fl. Alf. p. 123. an?

Icon. Cavanill. l. c. t. 93. f. 2.

Germ. Rundblättriger Storchschnabel.

Gall. Bec-de-grue à févilles ronds.

Hab. Circa Carlsruhe et alibi in cultis, ru ratis, ad sepes et vias passim.

Flor. Majo, Junio. *Semina* Junio, Julio, t tura. ☉

Cau

Caules ex radice plures, palmares, pedatet longiores, prostrati, incrassato-geniculateretes, villosi-viscidi, inferne fusco-rubri, ramosi: *Rami* divaricati, viscidii. *Folia* ovata, longe petiolata, orbiculata, quinque-loba, crenata, pubescenti-mollia, subtus praemixta viscida, sinibus puncto fusco saepe notata: superiora minora, subcuneato-rotundata, basi. *Stipulae* acutae, breves, rubellae. *Flores* axillares terminalesque, pedunculati, bibracteati. *Pedicelli* *Bractea* tetraphylla, linearis, acuta, amoene rubra, suffulti. *Calyx* lobis quinque, ovato-acuta, pilosa, aristata. *Corollae* petala quinque, incarnato-purpurea, unguibus albido-viridibus, liriata, inaequalia, calyce paulo longiora. *Antherae* decem, breviter saepe.

Nota. Minime confundendum cum *G. malvae-folio*.

†††† *Pedunculis unifloris.*

1048. *GERANIUM sanguineum* pedunculis unifloris; foliis quinquepartitis trifidis orbiculatis. *Linn. Sp. pl. 2p. 958.* — *Pollich. fl. pal. n. 655.*

Geranium sanguineum caule articulado rubro hispido; foliis orbiculatis, profunde 5-7 partitis, laciniis trifidis: pedunculis solitariis longissimis unifloris articulatis. *Cavanill. l. c. p. 195. n. 269.*

Geranium pedunculis unifloris, foliis septilobis, petalis bifidis. *Hall. stirp. helv. n. 930.*

Geranium sanguineum maximo flore. *Bauh. pin. 318.* — *Mapp. fl. Alf. p. 125.*

130 MONADELPHIA. DECANDRIA.

Geranium XIII. *Tabernæmont. p. 126.*
p. 1158. Icon.

Icon. Oeder. fl. Dan. t. 1107. — Bulli
herb. t. 12. — Cavanill. l. c. t. 76. f.

Germ. Bluthrother Storchschnabel.

Gall. Bec-de Grue sanguin.

Hab. Utrunque in sylvaticis graminosis pas-
 In collibus apricis graminosis calcareo
 gillaceis ad montem *Thurnberg*, in vi-
 retro *Durlach*, *Grözingen* prope *W*
garten, *Bruchsal*, *Wissloch* et alibi
 sim frequens.

Flor. Majo, Junio, Julio. *Semina* Julio.
 gusto matura. ☞

Caules ex radice plures, pedales, fesc-
 pedales et longiores, teretes, geniculato-
 crassati, hirsuti, fusco rubentes, ramosi. *Fo*
 opposita, petiolata, orbiculata, profunde f-
 tem aut quinquepartita, laciniis trifidis, fi-
 rate viridia, villosa. *Stipulae* basi latæ, acu-
 reflexæ. *Flores* axillares, longe peduncul-
Pedunculi erecti, medio subinflexi, unifi-
Corolla speciosa, magna, patens. *Petala* f-
 quinea, venis saturationibus prædita, lata, fi-
 rotundo-obcordata.

Inter omnia Gerania germanica Corollis maxi-
 conspicuum. Verum hortorum decus.

Flores apibus mel concedunt.

MONADELPHIA. POLYANDRIA

CCCXXVII. A L T H A E A.

Linn. Gen. plant. ed. Schreb. n. 1132.

Calyx duplex: exterior monophyllus 6-
 fidus: laciniis angustissimis, minus persisten-

interior monophyllus semiquinquefidus. *Corolla* *alabastrina* fere quinquepartita, obcordata, *præcorollata*, plana, tubo flaminum basi affixa. *Germea* orbiculatum. *Stylus* cylindraceus, brevis. *Stigmata* numerosa, setacea, longitudine Styli. *Capsule* plurimæ, monospermæ. *Semina* compressa reniformia.

1049. *ALTHAEA officinalis* foliis simplicibus tomentolis. *Linn. Sp. pl.* 2. p. 966. *Pollich. fl. pal. n.* 656. *Cavan. Diff.* 2. p. 93.

Althæa tomentosa, herbacea, caule erecto, foliis cordato-lanceolatis, obsolete trifidis. *Hall. stirp. helv. n.* 1074.

Althæa Dioscoridis et Plinii. *Bauh. pin.* 315. *Mapp. fl. Als.* p. 19.

Althæa *Tabern: mont.* p. 1152. *Icon.*

Icon. Bulliard. herb. t. 373. — *Regnault. bot. Oeder. fl. Dan. t.* 530. — *Blackw. t.* 90. — *Schkuhr. bot. handb. t.* 192.

Germ. Eibisch. Heilwurz.

Call. Guimaure ordinaire.

Herb. Ad viarum fossarumque margines hinc inde infrequens. Prope *Durlach* cum *Teucrio Scordio* rarius. In Palatinatu ad vias ac in plateis desertis circa *Dürkheim*; ad fossas prope *Lomersheim* cum *Scordio* et *Samolo valerandi* ad vias humidiusculas inter *Oggersheim* et *dem Holzhof*; locis humidis inter *Durkheim* et *Lamsheim t. Pollichio*. In Alsatia prope *Marlenheim* et alibi passim.

Her. Julio, Augusto. *Semina* Augusto, Septembri matura. ♀

222 MOMADELPHIA. POLYANDRIA.

Radix magna, ramosa; ramis longis, ascendentibus, teretibus, simplicibus, grassis, digiti, extus cinereo-albida, transverse subgosa, intus alba, transverse striata. *Caulis* una radice plures, cubitales, tripedales et tiore, erecti, teretes, albido-tomentosi, molli. *Folia* alterna, petiolata, cordato-ovata inferiora subtriloba, inciso-crenata, tomentosa, mollissima, quinquenervia. *Flores* in ramis terminalibus subcompositis. *Corollae* exalbido-incarnatae, quandoque albae.

Pharm. ALTHÆÆ *Radix*; *Herba*; *Flores*

Planta tota, cum primis radicibus mucilaginosa, emulsiens, demulcens, involvens.

Flores apibus mel præbent.

Nota. Digna quæ ad fossas aquaticas propagetur.

1050. ALTHÆA *hirsuta* foliis trifidis, hispidis supra glabris. *Linn. Sp.* 2. p. 966. *Pollich. fl. pal. n.* 657.

Althæa hirsuta foliis cordatis, pilosis, hispidis supra glabris, radicalibus numero subrotundo-lobatis, caulinis trilobis quinquelobis, pedunculis solitariis u floribus. *Cavanill. l. c. p.* 95.

Althæa foliis radicalibus reniformibus, caulinis palmatis, trilobis et quinquelobis calycibus hispidis. *Hall. stirp. helv.* n. 1073.

Alcea hirsuta. *Bauh. pin.* 317. *Mapp. Alf. p.* 11.

Icon. Jacqu. fl. Austr. t. 170. *Cavanill. l. c.* 29. f. 1.

German. Haariger Eibisch.

Gall. Guimauve velue.

Hab. In Palatinatu ad vias gramineas inter *Oppenheim* et *Weinheim*; retro Alzey ad ambulacra graminea juxta agros, versus *Niederulm* et *Mainz* post *Pollichium* vidi. In Alsatia prope Argentinam in dumetis in *Illkircher Wald* versus agros hodie raro

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix fibrosa, albida. **Caulis** palmaris, sesquipedalis, ascendens, quandoque se decumbens, teres, inferne sæpe ramosus, iis patulis hispidus. **Folia** alterna, petiolata, seniora, reniformi-rotunda, subseptem vel pinqueloba, obtuse ferrato-dentata, superiora palmata, profunde quinqueloba, serrata, rima tripartita, omnia hispidiuscula. **Stipulae** lanceolatae, acutae, pilosae. **Pedunculi** cillares, solitarii, erecti, uniflori, longitudine vicum, pilosuli. **Calyx** exterior 7-8 partitus: **seminibus** lanceolatis, acutis, patentibus, ciliatis; **interior** quinquesfidus, longior, erectus, hirtus. **Corolla** quinquepartita, parva, purpureo-bida aut coerulea, laciniis obtusis, subcrenatis, calyce sublongioribus.

Ad rariores hodie floræ cives pertinet.

051. *ALTHÆA rosea* caule erecto, foliis rugosis, cordatis 5-7 angulatis. *Cavan. Diff.* 2. p. 91.

Alcea rosea foliis sinuato-angulosis. *Linn. Sp. pl.* 2. p. 966.

Malva rosea folio subrotundo. *Bauh. pin.* 315.

Malva arborea. *Tabernæmont. p.* 1148. 1149. *Icon.*

334 MONADELPHIA. POLYANDRIA.

*Icon. Cavanill. l. c. t. 28. f. 1. Regnault bo
Germ. Stockrosen-Eibisch: Pappelrose. Hd
rose.*

Gall. La Rose Tremiere ou Doutremer.

Hab. In Oriente. Colitur in hortis passim.

*Flor. Septembri, Octobri. Semina Octob
Novembri maturat. 7.*

*Caules ex radice sæpe plures, 5-9 pe
les, erecti. Folia alterna, petiolata, 5-7 fol
rugosa. Flores terminales, copiosi, racemati
Calyx duplex, exterior sæpe sexpartitus. 7
rollæ speciosæ, magnæ, profunde quinquep
titæ, saturate ex nigro rubræ, sæpius plen
venosæ.*

Multifariæ variationes in hortis coluntur.

Pharm. MALVÆ ARBOREÆ Flores.

In usum medicum cumprimis feliguntur quæ C
rollis atropurpureis instructæ sunt. Externe præcip
emolliendi causa adhibentur.

Egregium hortorum tempore inflorescentiæ dec
Flores apibus mel largiuntur.

CCCXXVIII. M A L V A.

Linn. Gen. plant. ed. Schreb. n. 1134.

*Calyx duplex, exterior triphyllus, ang
stior, foliolis cordatis, acutis, persistentibus
interior monophyllus, semiquinquefidus, m
jus, latius, persistens. Corolla profunde ad h
fin quinquepartita: segmenta obcordata, pr
morfa, plana, tubo staminum basi affixa. St
lus cylindraceus brevis. Stigmata plurima, i
tosa. Capsula subrotunda, composita. Semin
subsolitaria, reniformia.*

1052. MALVA rotundifolia caule prostrato, foliis cordato-orbiculatis obsolete quinquelobis, pedunculis fructiferis declinatis. *Linn. Sp. pl.* 2. p. 969. — *Pollich. fl. pal. n.* 658. — *Hall. stirp. helv. n.* 1070.

Malva vulgaris, flore minore, folio rotundo. *Bauh. hist.* 2. p. 549. — *Mapp. fl. Alf. p.* 187.

Malva sylvestris minor. *Tabern. p.* 1150. *Icon.*

A. Malva sylvestris rotundo variegato folio. *Mapp. l. c.*

Icon. Curtis fl. Lond. t. 157. — *Bulliard. herb. t.* 161. — *Cavanill. l. c. t.* 26. *f.* 3. — *Schkuhr. bot. t.* 192.

Germ. Gänsepappeln. Käsepappeln.

Gall. Petite Mauve.

Hab. Utrunque in ruderatis, cultis, ad vias, plateas, muros passim frequens.

Flor. Junio, Julio. *Semina* Julio, Augusto maturata. ☞

Caules palmares, pedales et longiores, decumbentes, teretes, villulosculi, ramoli. *Rami* alterni. *Folia* alterna, petiolata, cordata, subrotunda, obsolete quinqueloba, dentato-ferrata, subpubescentia, medio sæpe fusco-purpureascentia. *Stipulae* ovatae, acutae. *Flores* axillares, 3-5 fasciculati, pedunculati. *Corollae* parvae, albidæ, venis purpureis pictæ.

Pharm. MALVÆ VULGARIS f. MINORIS
Herba. Flores. Semina.

Cum reliquis malvaceis emollientes vires exercet.
Veteribus culinaris fuit.

1053. *MALVA sylvestris* caule
herbaceo, foliis septemlobatis
pedunculis petiolisque pilosis. *Linn.*
pl. 2. *p.* 969. *Pollich. fl. pal. n.* 659.

Malva caule erecto, foliis lobatis, lobis si-
ratis quinis et septenis. *Hall. stirp. he-*
n. 1069.

Malva vulgaris flore majore, folio sinuato
Bauh. hist. 2. *p.* 949. — *Mapp. fl. A-*
p. 187.

Malva sylvestris major. *Tabernæmont.*
1150. *Icon.*

Icon. Curtis fl. Lond. Bulliard. herb. t. 2:
Cavanill. l. c. t. 26. *f.* 2.

Germ. Waldmalve. Pferdeappel.

Gall. Mauve sauvage.

Hab. Utrinque in ruderatis, ad vias, sep-
muros; amat pagos, urbes, ruderata.

Flor. Junio, Julio. *Semina* Augusto matura

Differt a præcedente specie, *Caulibus* alt-
ribus 3-4 pedalibus, adscendentibus, quan-
que erectis, *Foliis* majoribus profundius 5
lobis. *Pedunculis* longioribus, *Corollis* mu-
majoribus albido-purpurascens, venis sa-
ratoribus pictis.

Variat *Corollis* niveis.

Pharm. MALVÆ MAJORIS f. SYLVESTR
Herba.

Vires præcedentis.

Flores apibus accepti.

1054. *MALVA mauritiana* caule erec-
herbaceo foliis quinquelobatis obtus
pedunculis petiolisque glabriusculis. *L*
Sp. pl. 2. *p.* 970.

ROSADELPHIA. POLYANDRIA. 137

Malva caule erecto subglabro foliis cordatis crenatis quinquelobatis, floribus numerosis magnis rubescentibus, stipulis connatis. *Cavanill. Diff. 2. p. 77.*

Malva hederaceo folio. *Bauh. pin. 315.*

an. Cavanill. l. c. t. 25. f. 2.

an. Mauritanische Malve.

an. Mauve mauritiane.

Malva Passim ad pagos et urbes cum præcedente specie. Circa *Carlsruhe am Holzmagazin* et alibi non infrequens.

Flor. Julio, Augusto. **Semina** Augusto, Septembri matura. ☉

Affinis præcedenti a qua differt, *Corollis* submajoribus saturatius rubris, venis purpurascens pictis. *Fructibus* pulcherrime reticulatis.

1955. **MALVA** *crispa* caule erecto, foliis angulatis crispis, floribus axillaribus glomeratis. *Linn. Sp. pl. 2. p. 970.*

Malva foliis crispis. *Bauh. pin. 315.*

Malva *crispa*. *Tabernæmont. p. 1151. Icon.*

Icon. Cavanill. Diff. 2. t. 23. f. 1.

Ger. Krausblättrige Malve.

Gall. Mauve frisée.

Malva In Syria. Circa *Carlsruhe am Holzmagazin*, ad viam über die Schieswiese, et alibi, forsan ex hortorum rejectamentis.

Flor. Julio, Augusto. **Semina** Augusto, Septembri matura. ☉

Folia elegantissime crispata. *Corollae* minores in præcedente specie.

Vires præcedentium.

1056. **MALVA** *Alcea* caule erecto, fr
multipartitis scabriusculis. *Linn. Sp.*
2, p. 971. — *Pollich. fl. pal. n.* 660.

Malva caule erecto, foliis scabris trilob
lobis lateralibus bipartitis, medio
partito et quinquepartito. *Hall. f.*
helv. n. 1071.

Alcea vulgaris major flore ex rubro rō
Bauh. pin. 316. *Mapp. fl. Alf. p.* 10

Alcea. *Tabernæmont. p.* 1155. *Icon.*

Icon. Cavanill. Diff. 2. t. 17. f. 2. — *Blæ*
t. 304.

Germ. Rosenpappel. Siegmarskraut.

Gall. Mauve Alceé.

Hab. Utrunque passim in sepibus et ad agro
margines.

Flor. Junio, Julio. *Semina* Augusto, 1
tembri matura. ♀

Caulis cubitalis, tripedalis et altior,
ctus, teres, piloso - scabriusculus, ramē
Rami alterni. *Folia* alterna, petiolata, pro
de quinquepartita, segmentis tri-quinque
titis, villosis. *Stipulæ* parvæ, subulatæ. *Fl*
terminales, paniculati, densi. *Calyx* dup
exterior triphyllus, interior semiquinquesi
major. *Corollæ* speciosæ, incarnato-purpu
Capsulæ glabræ, subreniformes, nigræ.

Pharm. ALCEÆ Radix; Herba.

Planta ob flores amoenos passim in hortis pri
tim anglicis colitur.

Flores apibus mel concedunt.

Rarius hodie in medicinam recipitur. Vires
thææ.

1057. *MALVA moschata* foliis radicalibus reniformibus incisfis, caulinis quinquepartitis pinnato-multifidis. *Linn. Sp. pl.* 2. p. 971. — *Pollich. fl. pal. n.* 661.

Malva foliis radicalibus reniformibus incisfis, caulinis quinquepartitis pinnatis, pinnis dentatis. *Hall. stirp. helv. n.* 1072.

Alcea folio rotundo laciniato. *Bauh. pin.* 316. *Mapp. fl. Als. p.* 10.

Alcea tenuifolia crispa. *Bauh. hist.* 2. p. 1067. — *Mapp. l. c. p.* 11.

Leon. Curtis fl. Lond. t. 228. — *Oeder. fl. Dan.* 1905. *Cavanill. Diff.* 2. t. 18. f. 1.

Germ. Moschuspappel.

Gall. Mauve musquée.

Hab. In Marggraviatu superiori ad aridos aggeres circa *Haslach* infrequens; ad agrorum margines versus *Hünigen*; circa *Höllstein*, in agro Basileensi ad *Langenbrück*. In Alsatia prope Argentinam ad *Ostwinkel* hodie rarius, ad *Marienkirch* frequens. In Sundgovia versus Rhenum, et ad *St. Cosmum t. Joh. Bauhino*. In Palatinatu Circa *Lautern* in aggeribus; inter *Frankenstein* et *dem Bicard* ad vias. Circa *Waldmohr*, *Kibelberg*, *Schönenberg*, *Sann*, cum Anthemide tinctoria post *Pollichium* vidi. In Sponhemiae aggeribus colliculosis, præcipitiis cum Jafone montana frequens.

Flor. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ♀

Radix ramosa, sublignosa, albida, prostrata. *Caules* ex una radice plures, cubitales, tripedales, erecti, teretes, ramosi, punctis asperis hirsutis, purpureis adspersi. *Folia* alterna,

340 MONADELPHIA. POLYANDRIA.

petiolata, inferiora cordato-reniformia subpartita: *laciniis* pinnatifidis, multifido-laciniatis, acutis, subhirsutis; superiora brevius tiolata, in tenuiores lacinias divisa. *Stipbinae*, erectae, lanceolatae, margine undulhirsutae. *Flores* terminales axillaresque, pediculati, fasciculati, solitariique. *Pedunculi* ciales, teretes, pilosi. *Calyx* duplex: *exter* triphyllus, foliolis lineari-lanceolatis; *inter* quinquemfidus, ovato-acutus, saepe lacinia punctato-hirsutus: margine serrato glandifus. *Corolla* speciosa, profunde ad basin quaequepartita, *laciniis* cuneiformi-triangulari emarginato-bifidis, carneis, venis saturatis, anastomosantibus notatis, basi albidis, z gine utrinque ciliatis. *Filamentorum* tubus alitus, pilosus. *Antherae* reniformes, carne dein purpureae, demum coerulescentes.

Flores Moschum redolent. Non confundenda praecedente specie.

Flores apibus melliferi.

CLASSIS XVII.
DIADELPHIA.

HEXANDRIA.

- CCCLXIX. FUMARIA. *Cal.* 2-phyllus. *Cor.*
ringens, basi gibbosa necta-
risera. *Filamenta* antheris 3.

OCTANDRIA.

- CCCLXXX. POLYGALA. *Cal.* 2-laciniae alæ-
formes. *Cor.* vexillum cylin-
dricum. *Stamina* connexa.
Caps. obcordata, 2-ocularis.

DECANDRIA.

* *Stamina omnia connexa.*

- CCCLXXI. SPARTIUM. *Filamenta* adhæren-
tia germi. *Stigma* adnatum.
villosum.
- CCCLXXII. GENISTA. *Pistillum* deprimens
carinam. *Stigma* involutum.
- CCCLXXIII. ULEX. *Cal.* diphyllus. *Legumen*
vix calyce longius.
- CCCLXXV. ANTHYLLIS. *Cal.* turgidus inclu-
dens *Legumen*.
- CCCLXXIV. ONONIS. *Legumen* rhombeum,
sessile. *Vexillum* striatum.

**** *Stigma pubescens*** (nec priorum notæ.)

CCCXLIV. COLUTEA. *Legumen* inflatum
supra basin dehiscens.

CCCXXXVI. PHASEOLUS. *Carina Stylus*
spirales.

CCCXXXVIII. OROBUS. *Stylus* linearis, te-
tiusculus, supra villosus.

CCCXXXVII. PISUM. *Stylus* supra carinam
villosusque.

CCCXXXIX. LATHYRUS. *Stylus* supra planum
villosusque.

CCCXL. VICIA. *Stylus* sub stigmate barbatus

***** *Legumen subbiloculare*** (nec priorum.)

CCCXLIX. ASTRAGALUS. *Legum.* 2-lo-
lare, rotundatum.

****** *Legumina submonosperma*** (nec priorum)

CCCL. TRIFOLIUM. *Legum.* vix calyce lon-
gius, 1-f. 2-spermum. *Fl-*
res capitati.

******* *Legumen subarticulatum.***

CCCXLVIII. HEDYSARUM. *Legum.* articul-
subrotundis, compressis. *Ca-*
rina obtusissima.

CCCXLV. CORONILLA. *Legum.* istimis inte-
ceptum, rectum.

CCCXLVI. ORNITHOPUS. *Legum.* artici-
latum, arcuatum.

VII. HIPPOCREPIS. *Legum.* compresso-membranaceum: altera futura emarginaturis ad medium excisa.

I. MEDICAGO. *Legum.* spirale, membranaceo-compressum. Pifillum carinam deflectens.

* *Legumen uniloculare polyspermum*
(nec priorum.)

III. ROBINIA. *Vexillum* reflexo-patens, subrotundum.

II. ERVUM. *Calyx* 5-partitus, subaequalis, longitudine fere corollae.

III. CYTISUS. *Legum.* pedicellatum. *Calyx* bilabiatus.

LOTUS. *Legum.* teres, farctum seminibus cylindricis.



CLASSIS XVII.
D I A D E L P H I A
H E X A N D R I A.

CCCXXIX. F U M A R I A.

Linn. Gen. pl. ed. Schreb. n. 1154.

Calyx diphyllus: *foliis* oppositis, æ-
libus, lateralibus, erectis, acutis, parvis,
ciduis. *Corolla* oblonga, tubulata, ringens,
lato prominente, faucem claudente. *Lab-*
superius planum, obtusum, emarginatum,
flexum. (*Vexillum*) *Nectarium*, est basis
superioris postice prominens. *Labium* inferius
simile superiori, versus basin carinatum. (*C-*
rina) *Nectarium* est basis carinata. *Faux* tri-
gonæ, obtusæ, perpendiculariter bifida. (*A-*
Filamenta duo æqualia, basi latiora. *A-*
theræ in singulo filamento tres, terminal
Stigma orbiculatum, compressum. *Silicula* r-
tunda aut oblonga, mono aut polysperma, m-
nolocularis.

1058. FUMARIA *cava* radice cava, cau-
simplici, bracteis florum integris. *Hoffm-*
fl. Germ. p. 248.

Fumaria bulbosa var. *æ cava.* *Linn. Sp. 1*
2. p. 983.

Fumaria radice bulbosa, caule simplici l-
folio, bracteis ovato-lanceolatis. *Ha-*
stirp. helv. n. 348.

Fum

DIADELPHIA. HEXANDRIA. 145

Fumaria bulbosa radice cava major. Bauh. pin. 143. Mapp. fl. Alf. p. 114.

Fumaria bulbosa alba et purpurea, I. II. III. IV. Tabernæmont. p. 92. 93.

Ger. Oeder. fl. Dan. t. 605. — Blackw. t. 534

Ger. Hohler knolliger Erdrauch.

Call. Fume terre bulbeuse.

Hab. Utrunque passim in nemoribus, ad sepes circa pagos et urbes. Circa Carlsruhe prope Mühlberg an der Ziegelhütte et prope Grünwinkel frequens.

Flor. Martio, Aprili, Majo. Semina Junio maturat. 7

Radix bulbosa, subrotunda, fusco-ferruginea, junior solida, adulta cava. Caulis palmatus, pedalis, et altior, erectus, glaber, simplicifolius. Folia alterna, petiolata, tripinnatifida, incisa, glabra. Racemus terminalis. Bractee florum solitariae, ovato-lanceolatae, integerrimae, glabrae. Corollae calcaratae, purpurascens, incarnatae, vel albidæ.

Pharm. ARISTOLOCHIAE CAVÆ Radix.

Panta florens pulchra. Flores apibus mel præbent.

1059. *FUMARIA solida* radice solida bulbosa, caule simplici, bracteis digitatis. *Hoffm. fl. germ. 248.*

Fumaria bulbosa var. β. et γ. solida. Linn. Sp. pl. 2. p. 933. — Pollich. fl. pal. n. 662.

Fumaria radice non cava, major et minor, Bauh. pin. 144. — Mapp. fl. Alf. p. 114. 115.

Fumaria bulbosa minor. Tabernæmont. p. 94. Icon.

146 DIADELPHIA. HEXANDRIA.

Icon. Oeder. fl. Dan. t. 1224. — Curtis M t. 291.

Germ. Dichter Erdrauch.

Gall. Fume terre bulbeuse solide.

Hab. Utrinque præsertim in montosis ad passepes, pomaria.

Flor. Martio, Aprili. *Semina* Junio matur.

A præcedente specie distinguitur. *Radix* folida, *Bracteis* cuneiformibus, digitatis.

Pharm. ARISTOLOCHIAE FABACIAE Radix.

Olim vires vulnerariae radicibus hujus et præcedentis speciei tribuebantur. Hodie vix in usu sunt.

Flores melliferi.

1060. *FUMARIA officinalis* pericarmonospermis racemosis, caule diffusilinn, *Sp. pl. 2. p. 984. Pollich. pal. n. 4*

Fumaria foliis multifidis, lobis subrotundilanceolatis, fructibus monospermis. *Hist. stirp. helv. n. 346.*

Fumaria officinarum et Dioscoridis. *Bapin. 143. Mapp. fl. Alf. p. 114.*

Fumaria prima. *Tabernæmont. p. 85. Ic*

Icon. Curtis fl. Lond. t. 112. Oeder. fl. D t. 940. — Bulliard. t. 189. — Blackw. t. 2

Germ. Gemeiner Erdrauch. Taubenkropf.

Gall. Fumeterre.

Hab. Utrinque in cultis, ruderatis, agris, neis, passim abunde.

Flor. Majo, Junio. *Semina* Julio, Augusto matura. ☉

Radix fibrosa, fusco-flava. *Caules* palmares, pedales, cubitales; diffusi, angulosi,

Herba glabri, succulenti, ramosi, subflexuosi. Folia alterna, petiolata, bipinnata, glauca, pubescentia, pinnis bitrifidis. Flores caulem ramumque terminantes, pedunculati, spicato-racemosi. Pedicelli clavati. Corollae parvae, callosae, ex viridi alboque purpurascens, quantitate albidæ. Pericarpium Silicula uniloculari, subcordata, monosperma. Semen subrotundum.

Fig. FUMARIÆ Herba.

Herba succulenta, amariuscule, detergens, resoluens, aperiens, corroborans.

Nota. Caveant Pharmacopoei et medici ne confundant Fumariæ herbam cum Geranio Cicutario aut Robertiano, quod paucis ab hinc tempore in pharmaceutica quadam officina factum esse vidi.

1661. *FUMARIA parviflora* spicis laxis, filiculis monospermis globosis, acuminatis, caule diffuso, foliolorum laciniis, linearibus canaliculatis. *Smith. fl. brit.* 2. p. 750.

Fumaria filiculis globosis monospermis, foliis tenuissime divisis, laciniis ultimis angustis setaceis. *Lamark. Encycl.* 2. p. 561.

Fumaria foliis tenuissimis, floribus albis, circa Monspelim nascens. *Vaill. bot. parisi.* 56.

Fumaria minor tenuifolia, caulibus procumbentibus et caducis. *Bauh. pin.* 194. — *Morif. hist.* 2. Sect. 3. p. 261. t. 12. f. 11.

1672. *Vaillant. l. c. t.* 10. f. 5.

Curm. Kleinblütiger Erdrauch.

Call. Fume terre à fleurs petites.

148 DIADELPHIA. HEXANDRIA.

Hab. Circa *Baden* in cultis et arvis, frequenter ibi ad arcem in horto aulico.

Flor. Majo, Junio. *Semina* Julio maturantur.

Differt a præcedente specie, *Foliis* tenuioribus, *foliolorum* laciniis angustioribus, linearibus, canaliculatis; *Spicis* laxis, *Comminoribus* albidis.

Obs. Planta in horto botanico culta, constans. Metinet locum inter *F. officinalem* et *spicam*. Differt a *F. spicata*, *Caulibus* dinon erectis, *Spica* non densa, ovali, sed ovali-cylindrica. *Corollis* non rubris, sed

Individua hujus in Hispania et gallia meridiana lecta, nostratibus simillima sunt.

Nota. *Fumariam spicam*, quæ in Hispania cultis cum hacce specie abunde crescit et locos delectat, frustra hucusque in terris Germanicis quæsi.

1062. *FUMARIA claviculata* caulescens scandente, filiquis oblongis pressis dispermis, foliis inferioribus ternatis, rameis simpliciter ternatis hofis. *Linn. Sp. pl. ed Willden. 3. p.*

Fumaria claviculata filiquis linearibus laevibus cirrhiferis. *Linn. Sp. pl. 2. p. 9*

Fumaria claviculis donata. *Bauh. pin. — Moris. hist. 2. p. 266. Sect. 3. fig. 3.*

Fumaria alba latifolia. *Raj. angl. 3. p. Icon. Oeder. fl. Dan. t. 340.*

Germ. Gabelntragender Erdrauch.

Gall. Fume Terre claviculée.

Hab. Circa *Carlsruhe am Holzmagazin*, dubio ex Horto botanico migravit.

STADELPHIA. HEXANDRIA. 149

v. Majo, Junio. Semina Julio matura. ☉
Caulis copiosi, ramosi. Rami apicibus
inf. Folia latiuscula, subcirrhosa, lacte
is. Corollæ albæ,

STADELPHIA. OCTANDRIA.

CCCCXX. POLYGALA.

Gen. plant. ed. Schreb. n. 1154.

Herb. pentaphyllus, parvus: foliolis ova-
tis: duobus infra corollam, unico supra
eam, duobus mediis subovatis, planis, maxi-
coloratis, (Alae) persistens. Corolla sub-
spicea. Vexillum fere cylindricum, tu-
um, breve; ore reflexo, parvo bifidum.
Limbo concava, compressa, versus apicem
reticulata. Stamina octo, connata, carina in-
te. Capsula obcordata, compressa, margine
ext. bilocularis, bivalvis, Dissipimento val-
utinario, margine utrinque dehiscens. Se-
mina solitaria, ovata, umbilico glanduloso,

603. POLYGALA vulgaris floribus cri-
statis racemosis, caulibus herbaceis sim-
plicibus procumbentibus. *Linn. Sp. pl.*
2. p. 986. — Pollich. fl. pal. n. 664.

Polygala caule erecto, foliis linearibus acu-
tis. Hall. stirp. helv. n. 344.

Polygala major. Bauh. pin. 215. — Mapp.
fl. Als. p. 243.

12. Oeder. fl. Dan. t. 316. — Bulliard. herb.
t. 177

Germ. Gemeine Kreuzblume.

Gall. Laitier commun.

Hab. Utrunque in sylvis gramineis, pratis tam siccis quam humidis, pascuis montosis sylvaticis, collibus gramineis, juniperitis, ericetis ubique frequens.

Flor. Majo, Junio. *Semina* Julio matura. 7

Radix repens, lignosa, fibrosa. *Caules* digitales, palmares, pedales, adscendentes, quandoque procumbentes, angulosi, glabri, inferne ramosi. *Folia* alterna, inferiora latiora, superiora sensim longiora, angustiora, sparsa, sessilia, linearia, acuta, glabra. *Spica* racemosa, laxa, terminalis. *Bractea* 3, lanceolatae, breves, violaceae, aut coeruleae, ad singulorum pedicellorum basin. *Corollae* coeruleae, violaceae, incarnatae, purpureae, quandoque niveae. *Corpuscula* amoena, duo, tripartita, penicilliformia, carinae versus apicem affixa.

Pro diverso solo natali mire ludit.

Pharm. POLYGALÆ VULGARIS Radix; Herba.

Herba amariuscula vaccarum lac auget, quare Polygalæ nomen.

Flores apibus mel concedunt.

Infusum herbæ loco Theae chinensis tam ob gratum saporem quam ob salubritatem commendatur. Lübeckische Anzeig. 1758. 4. St.

1064. *POLYGALA amara* floribus cristatis racemosis, caulibus erectiusculis, foliis radicalibus obovatis, majoribus. *Linn. Sp. pl.* 2. p. 987.

Polygala procumbens, foliis imis subrotundis, supremis linearibus. *Hall. stirp. helv.* n. 343.

Polygala buxi minoris folio. Vaillant. bot. parif. p. 161.

Polygala vulgaris, foliis circa radicem rotundioribus, flore coeruleo, sapore admodum amaro. Bauh. pin. 215.

Icon. Vaillant. l. c. t. 32. f. 2. — Oeder. fl. Dan. t. 1169.

Germ. Bittere Kreuzblume.

Gall. Laitier amère.

Hab. In pratis montosis udis aquosis passim. *Auf dem Kaltenbrunnen, prope Herrnalb, Frauenalb, auf der Herrnwiefe.* In Margraviatu superiore in sylva *Wolfsaker* prope *Mültheim* et circa *Zienhen*. In Principatu superiore ad *Lacum Bodamicum* in pascuis udis passim abunde.

Flor. Junio, Julio. *Semina* Julio, Augusto matura. ♀

Præcedenti affinis, a qua distinguitur. *Caulibus* ex una radice copiosioribus, demissioribus, digitalibus, adscendentibus, aut procumbentibus, *Foliis* radicalibus et caulinis inferioribus latioribus, subrotundo-obovatis, firmiter, *Floribus* minoribus, cæruleis, albidis, vel incarnatis.

Pharm. POLYGALÆ AMARÆ Radix; Herba.

Radix et Herba valde amara, lingue diu inhaerens, incidens, resolvens, roborans, in pectoris morbis, plethisi præsertim pituitosa, magno sæpius cum succella exhibetur, et a quibusdam radici *Polygalæ Senegæ* substituitur.

1065. *POLYGALA Chamaebuxus* floribus imberbibus sparsis: carina apice subrotundo, caule fruticoso, foliis lanceolatis. *Linn. Sp. pl. 2. p. 989.*

252 **DIADELPHIA. OCTANDRIA.**

Polygaloides procumbens, foliis duris, ovatis, nervo aristato. *Hall stirp. helv. n. 34*

Polygala fruticosa, Buxifolio, flore flavicente et ex purpura rubente. *Tourneinst. — Mapp. fl. Als. p. 244.*

Chamæbuxus flore Coluteæ. *Bauh. pin. 47*

Icon. Jacqu. fl. Austr. t. 233.

Ger m. Buchsblättrige Polygala,

Gall. Laitier à fêvilles de Buis.

Hab. In Principatu Fürstenbergenſi in rupib calcareis prope Fürſtenberg abunde. In E tionis Baſileenſis monte *Hauenſtein* prope *Langenbruk* frequens, nec non prope *Baſſal.* In Alſatia prope *Hochfelden* de *Hochgebürg.*

Flor. Majo, Junio. *Semina* Julio, Auguſto matura. *Fruticulus.*

Radix repens, multiceps, longa, gracilis *Caulis* frutescens, adſcendens, sæpe procumbens, digitalis, palmaris. *Folia* alterna, ſubpetiolata, elliptica, coriacea, dura, ſempe virentia, integra, buxeis analoga, ariſta bre terminata. *Flores* axillares, pedunculati. *Fundunculi* ſolitarii, uniſlori, aut biſlori, ſquamliſbracteati. *Corolla* ſpecioſa, ochroleuca. *Apice* macula crocea ſaut purpurea pictæ. *Crinæ* criſtæ nullæ.

Variat. Calycibus aliſque elegantiffime ex rubro puraſcentibus.

Nota. Ad rariores Floræ cives hic elegantiffim in flore fruticulus pertinet.

Nullibi elegantius et copioſius florentem vi quam in Principatu Fürſtenbergenſi in rupib calcareis prope Fürſtenberg ubi certo inſig decus.

Difficilius in horto cicuratur.

DIADELPHIA. DECANDRIA, 153

DECANDRIA

CCCXXXI SPARTIUM

Linn. Gen. pl. ed. Schreb. n. 1166.

Calyx monophyllus, summo margine brevissimus, inferne versus apicem quinque densissimus, notatus, coloratus, parvus. *Corolla* papilionacea, pentapetala. *Vexillum* obcordatum, minimum, reflexum. *Alae* ovatae, oblongae, vix breviores, filamentis annexae. *Carina* bipetala, lanceolata, oblonga, alis longior, margine carinali villis connexo, filamentis inserta. *Stamina Filamenta* decem, connata. *Stigma* villosum. *Legumen* cylindricum, longum, obtusum, uniloculare, bivalve. *Semina* plura, globoso-reniformia.

1066. *SPARTIUM scoparium* foliis ternatis folitariisque, ramis inermibus angulatis. *Linn. Sp. pl. 2. p. 996.* — *Pollich. fl. pal. n. 665.*

Spartium foliis inferioribus ternatis, hirsutis. superioribus simplicibus. *Hall. stirp. helv. n. 354.*

Cytiso Genista scoparia, vulgaris, flore luteo. *Mapp. fl. Alf. p. 90.*

Genista angulosa et scoparia. *Bauh. pin. 395.*

Genista. *Tabernæmont. p. 1509. Icon.*

Leon. Curtis fl. Lond. Oeder. fl. Dan. t. 313.

Blackw. t. 244. — *Schkuhr. bot. handb. t.*

195. — *Pfleger arb. et frut. Bad.*

Ger m. Pfriemen. Gienst.

154 DIADELPHIA. DECANDRIA.

Gall. Génét commune.

Hab. Utrunque in sabulosis sylvis sterilibus facioribus, aridis. Abunde in sylvis arenosis circa *Carlsruhe* et alibi.

Flor. Majo, Junio. *Semina* Julio maturant.

Frutex 3-4-8 pedalis et altior, erectus sempervirens. *Rami* alterni angulati, copiose graciles, virgati. *Corollae* speciosae, sulphureae. *Vexillum* magnum, obtusum, patens, basiflexum, striis fulvis notatum. *Legumina* marginata pilosa, nigricantia, compressa, elasticè dilata cum strepitu.

Pharm. SPARTII f. GENISTAE SCOPARI.
Summitates; Folia; Flores; Semina.

Planta masticata nauseosa, amara, adstringens. Flores inaperti Capparidum instar conditi in quibusdam regionibus comeduntur, sed minus grati sunt; spermatophyta mel praebent.

Semina nauseosa purgant, et vomitum excitant.

Caules teneriores a pauperculis straminis loco adhibentur. Caeterum ad multifarios usus domesticos est requisita planta. Ad coria paranda apta. Cortex maceratus fila crassa ad telas viliores dat apta. *Schreber*

In naturae oeconomia maximi momenti, defensionis arborum cotyledones in montosis arenosis praefertim sylvae nigrae contra aeris injurias, arenamque figunt etc.

Lignum ex annosioribus luteo-rubellum, durum. Planta combusta, multum Salis alcalini relinquit.

Nota. In hortis anglicanis Carlsruhensibus hospitatur *Spartium junceum* et *radiatum*.

CCCXXXII. GENISTA.

Linn. Gen. pl. ed. Schreb. n. 1167.

Calyx monophyllus, tubulatus, parvum bilabiatus: *labio superiori* bidentato, profunde

in diviso; *inferiori* tridentato, subæquali. *Corolla* papilionacea. *Vexillum* oblongum, a *carina* remotum, totum reflexum. *Alæ* oblongæ, laxæ, breves. *Carina* recta, emarginata, *veulo* longior. *Stamina* Filamenta decem, *ovata*. *Stigma* acutum, involutum. *Legumen* ~~filamentum~~ turgidum, uniloculare, bivalve. *Semina* folitaria, subreniformia.

† *Inermes*.

1067. GENISTA *sagittalis* ramis ancipitibus, articulatis, foliis ovato-lanceolatis. *Linn. Sp. pl.* 2. p, 998. — *Pollich. pal.* n. 666.

Genista foliis cauli adpressis ellipticis, fine patenti ovato-lanceolato. *Hall. stirp. helv.* n. 353.

Genistella herbacea sive *Chamaespartium*. *Bauh. hist.* 1. p. 393. *Mapp. fl. Alf.* 122.

Chamægenista sagittalis. *Bauh. pin.* 395.

Chamaespartium primum. *Tabernæmont. p.* 1512. *Icon.*

Icon. Jacqu. fl. Austr. t. 209.

Germ. Pfeilförmiger Ginster.

Gall. Génét herbacée.

Hab. Circa *Carlsruhe*, *Mühlburg* et alibi in pascuis sterilibus graminosis et sylvis sabulosis ericetosis passim frequens.

Flor. Junio. *Semina* Augusto matura. *Suffrutex.*

Caules ex radice plures, adscendentes, palmatis, pedales, inferne sublignosi, superne herbacei, a foliis decurrentibus alati. *Folia*

156 DIADELPHIA. DECANDRIA.

alterna, subsessilia, lanceolata, integra, pilosa.
Flores terminales, racemoso-spicati. *Corolla*
 lutea. *Legumina* pilosa, compressa, nigra,
 4-6 sperma. *Semina* suborbiculata, compressa,
 glaberrima, fusco-flavicantia.

In pascuis arenosis sterilibus sæpe arenam
 Planta junior ovibus et capris pabulum bonum
 pet. Flores apibus mel concedunt.

1068. *GENISTA tinctoria* foliis lance-
 latis glabris, ramis striatis, teretibus er-
 ctis. *Linn. Sp. pl.* 2. p. 998. — *Pollic-
 pal.* n. 667.

Genista inermis foliis glabris, confertis, elli-
 pticis, lanceolatis, floribus sessilibus, fructu
 catis. *Hall, stirp. helv.* n. 350.

Genista tinctoria germanica, *Bauh. pin.* 39.
 — *Mapp. fl. Alf.* p. 121.

Genistella tinctoria. *Tabernæmont.* p. 151.
Icon,

Icon. *Oeder. fl. Dan.* t. 526. *Pfleger. arb.*
frut. Bad.

Germ. Färbe-Ginster,

Gall. Genêt des Teintres.

Hab. Utrunque in pratis pascuisque siccis
 ceo-arenosis et calcareis passim non in-
 quens.

Flor. Junio, Julio. *Semina* Augusto matur-
Suffrutex.

Caulis basi ramosus, lignosus, pedalis, ce-
 bitalis, erectiusculus, quandoque adscender-
Rami sparsi, angulati, glabriusculi, viride-
Folia alterna, sessilia, conferta, lanceola-
 acuta, integra, glabriuscula. *Stipulae* binæ
 foliorum basin, subulatæ, parvæ. *Flores* te-

simples, racemosi, denfi. *Corollæ* luteæ. *Le-*
gumen glabrum polyspermum.

Pharm. Genistæ Herba; Flores; Semina.

Vires Spartii scorparii habet.

Tactoribus ad fila colore luteo et viridi imbuenda
inferri; cum aqua calcis cocta, creta confusa et ite-
rum cum alumine cocta, reddit colorem saturate fla-
vum pictoribus sub nomine (Schuttgelb) notum.
Böhner techn. Gesch. der Pflanzen. 2. p. 221.

Frutex in hortis anglicis elegans.

Flores apibus mel præbent.

1069. *GENISTA pilosa* foliis lanceolatis
obtusis caule tuberculato decumbente.
Linn. Sp. pl. p. 999. — Pollich. pal.
n. 668.

Genista inermis procumbens, foliis duris
subhirsutis, spicis floriferis brevibus. Hall.
stirp. helv. n. 351.

Genista ramosa, foliis Hyperici. Bauh. pin.
395. — Mapp. fl. Als. p. 122.

Icon. Jacqu. fl. Austr. t. 208. — Pfleger arb.
et frut. Bad. — Schkuhr. bot. t. 195.

Germ. Haariger Ginster.

Gall. Genêt rampant.

Hab. In sylvaticis montosis arenosis et granita-
ceis utrinque passim abunde. Prope *Frau-*
enalb, Herrnalb, Reichenbach, Kalten-
brunn, auf dem Tobel, im Bühler-, Oppen-
euer- Durbacher- Thal. In Margg. sup.
prope *Badenweiler, Oberweiler, Schweig-*
hof et alibi in montosis apricis abunde.
In Palatinatu circa *Lautern, Heidelberg,*
Schriesheim et alibi in montosis. In Alta-
tiz Vogelis passim frequens.

Flor. Junio, Julio. *Semina* Augusto maturat.

Fruticulus palmaris, pedalis, cubitus procumbens. *Rami* angulosi, tuberculati. *Folia* alterna, terna, simplicia, lanceolata, obtusa, integra, superne glabra, inferne sericea. *Stemmi* terminales saepe longissimi. *Corollae* luteae. *Legumina* scabriuscula, piloso-sericea.

In montosis arenosis sabulum figit. Rupes graecae decompositas in superficie, saepe pulcherrime deaurat.

Folia capris bonum pabulum.

Flores apibus mel concedunt.

†† *Spinosae.*

1070. GENISTA germanica spinis compositis, ramis floriferis inermibus, foliis lanceolatis. *Linn Sp. pl.* 2. p. 999. *Pollich. fl. pal.* n. 669.

Genista foliis ovato-lanceolatis, hirsutis spinis in ramis senescentibus ramosis. *Hall. stirp. helv.* n. 352.

Genista Spartium minus, germanicum. *Mart. fl. Als.* p. 121.

Genista minor spinosa, germanica. *Bauh. pin.* 395.

Genistella aculeata. *Tabern.* p. 1510. *Icon.*

Icon. Pfleger arb. et frut. Bad.

Germ. Teutscher Ginster. Stechginster.

Gall. Genêt germanique.

Hab. Utrunque in sylvis aridis arenosis, collibus apricis dumetosis asperis. Prope *Durlach* retro dem *Thurnberg*, prope *Grözingen* et alibi, minus frequens.

Flor. Majo, Junio. *Semina* Julio, Augusto matura. **B**

DIADELPHIA. DECANDRIA. 159

Frutex sesquipedalis, cubitalis. *Rami* ves-
si, spinosi, pungentes. *Spinæ* compolitæ.
in ramis junioribus, alterna, ovato-lan-
cea, hirsuta. *Flores* terminales, racemosi.
corollae flavæ. *Legumina* brevia, scabriuscula,
baccata, nigra, 2-3 sperma.

Ex cineribus Genistarum multum salis alcalini ob-
tinetur.

Nota. In hortis Carlsruhanis adhuc hospitatur *GENISTA* sibirica. Linn.

CCCXXXIII. U L E X.

Linn. Gen. pl. ed Schreb. n. 1169.

Calyx diphyllus, persistens: *foliolis* ova-
to-oblongis, concavis, rectis, aequalibus, ca-
pulis paulo brevioribus. *Corolla* papilionacea,
emipetala. *Vexillum* obcordatum, emargina-
tum, erectum, maximum. *Alae* oblongue, ob-
tusæ, vexillo breviores. *Carina* dipetala, recta,
obtusæ, margine inferiore connivens. *Stamina*
Filamenta decem, connata. *Legumen* oblon-
gum, turgidum, calyce vix longius, rectum,
uniloculare, bivalve. *Semina* pauca, subrotun-
da, emarginata.

1071. *ULEX europæus* foliis villosis
acutis, spinis sparsis. *Linn. Sp. pl. 2. p.*
969.

Ulex dentibus calycinis obsoletis conniven-
tibus, bracteis ovatis laxis, ramulis cre-
ctis. *Smith. fl. Brit. p. 756.*

Ulex folio sub singulis spinis subulato, pla-
no, acuto. *Linn. hort. Cliff. 356.*

160 DIADELPHIA. DECANDRIA.

*Genista spartium majus brevioribus et
gioribus aculeis. Tournef. inst. M.
fl. Alf. p. 121.*

*Genistellæ spinosæ affinis, Nepa quibus
Bauh. hist. 1. p. 400.*

*Scorpius primus. Clus. hist. 1. p. 106.
bernæmont. p. 1515. Icon.*

*Icon. Oeder. fl. Dan. t. 608. — Schluhr.
Handb. t. 196.*

Germ. Europäischer Hecksame.

Gall. Ulex européen.

*Hab. In Alsatia montibus circa Besfor
Mappo.*

*Flor. Per totam æstatem. Semina Septem
Octobri maturat. 5*

*Frutex 2-4 pedalis et altior, erectus,
molissimus, quammaxime spinosus. Rami in
spinam et folium, juniores fulcati, angu-
lirfuti, virides, spina terminati. Spinæ lon-
validæ, spinulis foliisque minoribus auc-
apice albidæ, pellucidæ. Folia alterna, sp-
similia, infra easdem. Flores axillares, nec
ex spinis prodeuntes, breviter peduncu-
Bractæ binae, calyci oppositæ, ovatae.
lyx diphyllus, foliolis integriusculis, vill-
luteolis. Corollæ luteæ, villosulæ. Legum-
oblongum, turgidum, nigricans, villosum,
spermum.*

*Nota. Hunc Fruticem hucusque frustra in terris
densibus quæsvi.*

*Ad sepes densissimos impenetrabiles circa ho-
formandos commendatur.*

DIADI

CCCXXXIV. ONONIS.

Linn. Gen. pl. ed. Schreb. n. 1173.

Calyx quinquepartitus, longitudine fere corollae; *laciniis* linearibus, acuminatis. *Corolla* papilionacea. *Vexillum* cordatum, striatum. *Alae* ovatae, vexillo dimidio breviores. *Carina* acuminata, alis fere longior. *Stamina* Filamenta decem, connata. *Stigma* obtusum. *Legumen* rhombeum, turgidum, subvillosum, uniloculare, bivalve. *Semina* pauca, reniformia.

1072. *ONONIS spinosa* floribus axillari-
bus solitariis, foliis ternatis; superioribus
solitariis, ramis spinosis.

Ononis spinosa floribus racemosis solitariis
foliis ternatis solitariisque, caule spinoso.

Linn. Syst. veg. p. 651. Poll. fl. pal. n. 670.

Anonis caule decumbente, ramis senescen-
tibus spinosis. Hall. stirp. helv. n. 356.

Anonis spinosa flore purpureo. Bauh. pin.
389. — Mapp. fl. Alf. p. 24.

Ononis prima. Tabernæmont. p. 913. Icon.

Leon. Regnault. bot. — Bulliard. herb. t. 105. —

Oeder. fl. Dan. t. 783.

Germ. Stachlichte Hauhechel. Ochsenbrech.

Gall. l'Arrete Boeuf. — Bougrande.

Herb. Utrinque in arvis et agris neglectis, ad
pascua sabulosa arida passim frequens.

Flor. Junio, Julio, Augusto. Semina Augusto,
Septembri maturat. 7

Radix profunde descendens, cylindramosa, tenax, extus fusca, intus alba. Caex una radice plurimi, pedales, sesquipedales, et altiores, adscendentes, sæpissime erecti, teretes, villosi, rubelli, ramosi, spinosi. *Folia* tiolata, alterna, inferiora ternata, superiora simplicia, ovalia, obtusa, serrulata, glabriuscula. *Stipulae* rotundato-ovatae, bifidae, margine integræ, glabriusculæ. *Flores* axillares ternalesque, solitarii, breviter pedunculati. *Cilium* piloso-glandulosum. *Corollæ* purpureæ, innatae, albidæ. *Vexillum* striis longitudinalibus purpurascentibus pictum. *Legumina* ventriplaniuscula, villosa - scabriuscula, calyce viora, 4-6 sperma.

Obs. Individua hujus spontanea et in horto botanico hospitantia erecta, rarius adscendentia, juvenilia et adultiora spinis instructa. Flores nunquam gemini, sed semper solitarii, axillares, alterni.

Pharm. ONONIDIS s. RESTÆ BOVIS Her

Planta peculiaris odoris, diuretica. Decoctio radices in Ischuria ex calculo vesicæ ubi emulsivaria atque mucilaginosae incassum adhibita fuere, in cum levamine adhibuit Bergius. vid. ej. med. p. 600. Planta agris et pratis noxia, eradicata. Combusta multum salis alcalini dat.

1073. *ONONIS mitis* floribus omnifloris, alternis, axillaribus, ternis simplicibusque subtus cum filis et calyce pilosis, caulibus ramis inermibus, villosis.

Ononis spinosa var. *mitis*. floribus simplicibus solitariis lateralibus, ramis inermibus. *Linn. Sp. pl. 2. p. 1006. Pol. pal. n. 670. (sub O. spinosa.)*

Ononis spinis carens purpurea. *Bauh. 389. Mapp. fl. Als. p. 25.*

arm. Stachellose Hauhechel.

all. L'Arrete bocuf sans epines.

ab. Utrunque in arvis, agris et pascuis sterilibus, pratisque passim. In collibus calcareo-argillaceis retro Durlach prope Grözingen, Stein et alibi frequens.

Var. Junio, Julio, Augusto. Semina Augusto, Septembri matura. ♀

Differt à præcedente specie, Caulibus ramisque inermibus, fusco-viridibus, Foliis Stipulae majusculis, subtus longius pilosis, Capitulis longius pilosis, glanduliferis, magis laetidis. Corollae forma, magnitudo, et color e præcedentis speciei.

Nota. Haec sine dubio sub Ononide arvensi Murray, Linnaei filii, Rothii, Hoffmanni, aliorumque passim botanicorum militat. à qua recedit, Floribus semper solitariis, alternis, nunquam geminis, Corollis non duplo majoribus quam in specie spinosa, sed ejusdem magnitudinis.

¶ Species spinosa et haec mitis ex seminibus jam ultra 16 annos in horto cultae, semper constantissimae.

Odor fortior, magis hircosus, quam in præcedente specie.

Var. ONONIDIS Radix; Herba.

Planta ob odorem fortiorem præcedente magis

074. ONONIS altissima floribus longius racemoso-spicatis, geminatis, foliis inferioribus ternatis, superioribus simplicibus serratis glabriusculis, stipulis amplexicaulibus latis, ferrulatis, caule erecto ramoso, piloso.

Ononis altissima caulibus erectis villosis
 mosis, ramis apice spicatis, stipulis
 ferratis amplexicaulibus. *Lamarck. cycl.*

Ononis hircina. Jacqu. Hort. Vind. 1. p.

Ononis mitior. Clus. hist. p. 99. Icon.

Icon. Jacqu. Hort. Vind. t. 93.

*Germ. Gröſte Hauhechel. Boks-Hauhechel.
 Gall. l'Arrete Boeuf puante.*

*Hab. In Brisgovia collibus apricis am Kaij
 ſtuhl paſſim non infrequens.*

*Flor. Junio, Julio, Auguſto. Semina Au-
 ſto, Septembri matura. 7.*

Differt a præcedente specie, *Caulibus* ſi-
 cioribus, 3 - 4 pedalibus, magis erectis,
 ridibus, *Ramis* patenti-erectis, ſemper viri-
 bus. *Foliis* duplo longioribus et latioribus
 utrinque glabris, læte viridibus. *Stipulis* a-
 plexicaulibus, multo latioribus et longioribus
 patentibus, ferrulatis, margine rarius pilosis
Floribus in *Spica racemosa* caulem ramosc
 terminante, ſemper binis, nunquam ſolitas
Corollæ ut in præcedentibus speciebus, palli
 purpureæ, non majores. *Legumina* infla
 pilosa, 2-3 ſperma.

Nota. Planta ſpecie diverſiſſima a præcedente, m-
 me confundenda cum *Ononide arve*
Hoffmanni, Rothii, aliorum botanicor-
 Porro non pro *O. foetente Allionii*
Pedem. t. 41. f. 1. habenda, quæ habitu
Stipulis parvis a noſtra planta longe recedit

Obſ. Ad rariores omnino Floræ cives pertinet.

Planta ex ſeminibus in horto ultra tredecim
 annos culta, mutati non ſuccubuit.

Odor plantæ maxime hircinus.

In hortis Carlsruhanis adhuc hoſpitatur *On-*
fruticosa Linn. pulcherrima.

CCCXXXV. ANTHYLLIS.

Linn. Gen. plant. ed. Schreb. n. 1174.

Calyx monophyllus, ovato-oblongus, ventriculus, villosus: ore quinquedentato, inæquæ, persistens. *Corolla* papilionacea. *Vexillum* longius, lateribus reflexis, ungue longitudine calycis. *Alae* duae, oblongæ, vexillo breviores. *Carina* compressa, longitudine alarum, bifida. *Stamina Filamenta* decem, connata. *Legumen* subrotundum, tectum, intra calycem, minimum, bivalve. *Semen* 1 - 2.

1075. *ANTHYLLIS vulneraria* herbaea, foliis pinnatis inæqualibus, capitulo duplicato. *Linn. Sp. pl. 2. p. 1012.* — *Pollich. fl. pal. n. 671.*

Vulneraria foliis radicalibus ovatis, caulinis pinnatis, umbellis ovatis. *Hall. stirp. helv. n. 398.*

Vulneraria rustica. *Bauh. hist. 2. p. 362.*

Lagopodium flore luteo. *Tabernæm. p. 909. Icon.*

Leon. Oeder. fl. Dan. t. 988. — *Schkuhr. bot. handb. t. 196.*

Germ. Wundkraut. Wundklec.

Gall. Vulnèraire des paysans.

Rob. Utrinque in pratis montosis apricis, ad viarum marginalia graminea, in collibus graminosis aridis passim copiose.

Flor. Junio, Julio. Semina Augusto matura. ♀

Caulis palmaris, pedalis, adscendens, tereti, pubescens. *Folia* caulina pinnata, foliola

166 DIADELPHIA. DECANDRIA.

maxime inæqualia, lanceolata; *impari* maxim
Flores terminales, denſi, in duo capitula ſu
rotunda, multiflora, digeſſi. *Bracteae* digitat
3-5-6 fidae, ſub ſingulo capitulo.

Ad plantas vulnerarias olim numerabatur,

Flores apibus mel largiuntur.

CCCXXXVI, PHASEOLUS.

Linn. Gen. plant. ed, Schreb. n. 1180.

Calyx monophyllus, bilabiatus: *labio* ſup
riore emarginato, inferiore tridentato. *Corol*
papilionacea. *Vexillum* cordatum, obtuſu
emarginatum, reclinatum, lateribus reflex
Alae ovatae, longitudine vexilli, unguibus le
gis infidentes. *Carina* cum contentis genita
bus ſpiratis! *Filamenta* diadelpa, intra c
rinam ſpiralia. *Stylus* ſpiraliter inflexus, ſuper
pubeſcens. *Legumen* longum, rectum, cori
ceum, obtuſum, cum acumine. *Semina* reſ
formia, oblonga, compreſſa.

1076. *PHASEOLUS vulgaris* volubi
floribus racemoſis geminis, bracteis c
lyce minoribus, leguminibus pendul
Linn. Sp. pl. 2. p. 1016.

Phaseolus vulgaris. Lob. Ic. 59.

Smilax hortensis ſ. *Phaseolus major. Bau
pin. 339.*

β. *Phaseolus coccineus* volubilis, floribus r
cemofis, geminis, bracteis calyce br
vioribus, leguminibus pendulis. *Linn.*

DIADELPHIA. DECANDRIA. 167

Phaseolus caule volubili, floribus racemosis, bracteis calycinis adpressis. Lamarck. Encyl. 3. p. 68.

Icon. β. Schkuhr. bot. Handb. t. 199.

Germ. Gemeine Bohne. β. Feuerbohne.

Gall Phaseole vulgaire. 2. Phaseole écarlate.

Hab. In India orientali. Colitur in hortis, agris, et vineis passim.

Flor. Majo, Junio. Semina Augusto, Septembri, Octobri matura. ☉

Multifarizæ prostant variationes, inter quas Phaseolus coccineus potius magis peculiarem speciem culinaris constantem, constituit.

Ufus fructuum culinaris notissimus.

1077. PHASEOLUS nanus caule erectiusculo lævi, bracteis calyce majoribus, leguminibus pendulis, compressis, rugosis. Linn. Sp. pl. 2. p. 1017.

Phaseolus vulgaris italicus humilis seu minor, albus cum orbita nigricante. Bauh. hist. 2. p. 358.

Germ. Zwergbohne.

Gall. Phaseole nain.

Hab. In India. Colitur passim cum præcedente.

Flor. et Semina cum præcedente matura. ☉

Ufus culinaris.

CCCXXXVII. P I S U M.

Linn. Gen. pl. ed. Schreb. n. 1184.

Calyx monophyllus, quinquefidus, acutus, persistens: laciniis duabus superioribus brevioribus. Corolla papilionacea. Vexillum latissi-

168 DIADELPHIA. DECANDRIA.

mum, obcordatum, reflexum, emarginatum cum acumine. *Alae* duæ, subrotundæ, convergentes, vexillo breviores. *Carina* compressa, femilunata, alis brevior. *Stamina* diadelphica. *Legumen* magnum, longum, teretiusculum (compressum deorsum) apice sursum acuminatum, uniloculare, bivalve. *Semina* plura globosa.

1078. *PISUM sativum* petiolis teretibus stipulis inferne rotundatis, crenatis, pedunculis multifloris. *Linn. Sp. pl.* 1026.

Pisum stipulis crenatis. Linn. Hort. Cliff.

β. *Pisum hortense majus. Bauh. pin.* 342.

γ. *Pisum sine cortice durior. Bauh. pin.* 342.

δ. *Pisum umbellatum* stipulis quadrifidis atque trifidis, pedunculis multifloris terminalibus. *Mill. dict. n.* 3.

ε. *Pisum quadratum*, majus quadratum. *Bauh. pin.* 342.

Icon. Regnault. bot.

Germ. Gemeine Erbsen.

Gall. Erve commune.

Hab. . . . Coluntur in hortis, agris et vineis.

Flor. Majo, Junio. *Semina* Septembri, Octobri matura. ☉

Ufus culinaris notus.

CCCXXXVIII. O R O B U S.

Linn. Gen. pl. ed. Schreb. n. 1185.

Calyx monophyllus, tubulatus, ore obliquo, quinquedentato, brevissimo. *Corolla* 1

ptoracea. *Vexillum* obcordatum, apice late-
 dusque reflexum, longius. *Alae* duæ, oblon-
 gæ, affurgentes, conniventes, longitudine fere
 æquæ. *Carina* inferne bifida, acuminata, af-
 furgens: marginibus conniventibus, parallelis,
 compressis, fundo vertricofo. *Filamenta* dia-
 delpha. *Stylus* filiformis, erectus. *Stigma* li-
 gnatum, latere inferiore a medietate ad apicem
 stipubescens. *Legumen* teres, longum, apice
 acuminato adscendente, uniloculare, bivalve.
Semina plurima, subrotunda.

1079. *OROBUS vernus* foliis pinnatis
 ovatis, stipulis semilagittatis, integerrimis,
 caule simplici. *Linn. Sp. pl.* 2. p.
 1028. — *Pollich. fl. pal. n.* 672.

Orobus caule simplicissimo, foliis senis ova-
 tis, acutis. *Hall. stirp. helv. n.* 416.

Orobus sylvaticus purpureus vernus. *Bauh.*
pin 351. — *Mapp. fl. Als. p.* 227.

1. *Orobus* latifolius parvo flore, prorsus albo.
Bauh. pin. 251. — *Mapp. fl. Als. p.* 228,

Leon. Blackw. herb. t. 208.

Germ. Frühlings-Walderbsen.

Gall. Orobe printannier.

Hab. Utrunque in sylvis umbrosis, præsertim
 montosis ad sylvarum marginatiam passim non
 infrequens. *β.* In Allatiæ montosis auf dem
Hochgebürg prope Monasterium *Peris* ver-
 sus dem schwarzen See t. *Mappo.*

Flor. Martio, Aprili, Majo. *Semina* Junio
 matura. *z*

Caules palmares, pedales, ex radice sæpe
 pures, erecti, angulosi, simplices, glabri.

170 DIADELPHIA, DECANDRIA.

Folia alterna, petiolata, pinnata; *foliolis* sessilibus, ovatis, acutis, integris, glazete viridibus. *Stipulae* cordato-sagittatae tegræ, latæ. *Flores* in pedunculo communi go laterali, erecto, racemosi. *Racemus* uteralis. *Corollæ* violaceo-purpureæ, quæque incarnatæ, rariùs albidæ, amœnæ.

Planta elegans, bonum præbet pecoribus pap
Flores apibus mel concedunt.

1080 *OROBUS tuberosus* foliis p
tis lanceolatis, stipulis semisagittati
tegerimis, caule simplici. *Linn. S.*
2. p. 1028. — *Pollich. fl. pal. n. 6.*

Orobis caule simplici, foliis fenis ellip
radice tuberosa. *Hall. stirp. helv. n.*

Orobis foliolis imis ovatis, superioribus
ceolatis, caule alato, simplici. *Cri*
austr. p. 377.

Orobis sylvaticus, foliis oblongis, gla
Mapp. fl. Alf. p. 227.

Astragalus sylvaticus. *Thal. herc. 80. t.*

ß. *Orobis tenuifolius* caule inferne quac
gulo, superne alato, foliis bi seu
jugis, linearibus, stipulis semisagit
Roth. fl. Germ. 2. p. 170.

Orobis sylvestris, angustifolius, Asphc
radice. *Bauh. pin. 351. — Mapp. fl.*
p. 227.

Orobis quorundam radice bulbosa, 1
albo. *Bauh. hist. 2. p. 326.*

Orobis pannonicus III. *Clus. hist. 2. p.*

Icon. Curtis fl. Lond. — Oeder. fl. Da
781. — *Schkuhr. bot. Handb. t. 200.*

DIADELPHIA, DECANDRIA. 171

Germ. *Knollige Walderbste*.

Hall. *Orobe tubercul.*

Lab. In sylvis montosis lapidosis, nec non in pratorum marginalibus sylvaticis passim cum β .

Fler. Majo, Junio. Semina Julio, Augusto matura. γ

Differt a præcedente affini specie, *Radice* terosa, *Caulibus* manifeste præsertim superne pappibus, alatis, alis membracis. *Foliorum* foliis senis, superne quandoque quatuor, lanceolatis, arista terminatis.

Variat. Caulibus ad terram prostratis, Stipulis sæpe ad basin denticulato-hamatis.

Variatio β . mere differt Foliolis angustioribus.

Corollæ albae et violaceo-purpureae.

In Scotia radix ob dulcem saporem pro cibo adhibetur et pro medicamento quo rustici et famem et sitim pectoris morbos depellunt. Hall. l. c.

Flores apibus mel præbent.

1081. *OROBUS niger* caule ramoso, foliis sexjugis ovato-oblongis. Linn. *Sp. pl.* 2. p. 1028. — Pollich. *fl. pal.* n. 674.

Orob. caule ramoso, foliolis ovatis duobus. Hall. *stirp. helv.* n. 418.

Orob. sylvaticus *Viciæ* foliis. Bauh. *pin.* 252. — Mapp. *fl. Alf.* p. 227.

Icon. Oeder. *fl. Dan.* t. 1170. — Schkuhr. *bot. Handb.* t. 200.

Germ. *Schwarze Walderbste*.

Hall. *Orobe noiratre*.

Hab. In sylvis montosis elatioribus passim. In Margg. superiore retro Oberweiler, Badenweiler et Seringen. In ditione Basileensi

572 DIABELPHIA. DECANDRIA.

in monte Muteto et *Mönchenstein* t. *Cass. Bauhino*. In Palatinatus sylvis montol circa *Steinbach* abunde; in sylvis circ *Lauterecken* t. *Pollichio*. In Allatiæ Vog lis passim, inter *Hafslach* et *Niedek* fr quens.

Flor. Majo, Junio. *Semina* Augusto matura.

Caulis sesquipedalis, cubitatis, et altio erectus, glaber, ramosus. *Rami* alterni, p tenti erectiusculi, subdivisi. *Folia* alterna, pu nata; foliolis 10-12-14, oppositis, ovato-o longis, acutis, arista terminatis. *Stipulae* la ceolatae, parvae, integrae. *Flores* in longis p dunculis axillaribus racemosi. *Racemi* 6-9 flori subsecundi. *Corollae* purpureae, demum dec loratae. *Legumina* teretia, longa, glabra, pe dula, nigra.

Planta ficcata, tota nigra evadit.

Radix dulcis mucilaginoso.

Bonum et acceptum herba pecoribus pabulu praebet.

Flores mel apibus concedunt.

CCCXXXIX. LATHYRUS.

Linn. Gen. pl. ed. Schreb. n. 1186.

Calyx monophyllus, semiquinquesidu campanulatus: *laciniis* lanceolatis, acutis, f perioribus duabus brevioribus; infima longior *Corolla* papilionacea. *Vexillum* obcordatur maximum, lateribus et apice reflexum. *Al* oblongae, lunulatae, breves, obtusae. *Caris* semiorbiculata, magnitudine alarum, iisque l tior, in medio introrsum dehiscens. *Filamen* diadelpha. *Stigma* a medietate Styli ad apicer

DIADELPHIA. DECANDRIA. 173

stipe villosus. *Legumen* longum, cylindraceum, vel compressum, acuminatum, unilobare, bivalve. *Semina* plura, cylindracea, pabosa, vel parum angulata.

† *Pedunculis unifloris.*

1082. *LATHYRUS Aphaca* pedunculis unifloris, cirrhis aphyllis, stipulis sagittato-cordatis. *Linn. Sp. pl.* 2. p. 1029. — *Pollich. fl. pal.* n. 675.

Lathyrus aphyllus, stipulis sagittatis latissimis. *Hall. stirp. helv.* n. 442.

Vicia lutea foliis convolvuli minoris. *Bauh. pin.* 345.

Aphaca. *Lobel. Icon.* 70. — *Mapp. fl. Als.* p. 26.

Icon. Curtis fl. Lond.

Germ. Gelbe Ackerplatterbse.

Gall. Gesse des blés.

Hab. In agris ad viarum marginalia nec non in vineis passim, Circa *Carlsruhe* inter segetes non infrequens.

Flor. Junio, Julio. *Semina* Augusto, Septembri matura. ☉

Caulis palmaris, pedalis, et altior, decumbens, debilis, tetragonus, glaber, subramosus, aphyllus. *Stipulae* oppositæ, maximæ, foliaceæ, cordato-sagittatæ, obtusiusculæ, utrinque ad basin denticulo notatæ, ceterum integre, subtus nervosæ, glaucae. *Cirrhi* solitarii, simplices, stipulis longiores. *Flores* longe pedunculati. *Pedunculi* axillares, solitarii, uniflori, stipulis longiores, tetragoni, *Bractea* minus subcalyce instructi. *Corolla* luteo-sulphurea,

174 **DIADELPHIA. DECANDRIA.**

Legumen sesquiunciale, compressum, latiusculum, glabrum. *Semina* 7-8, subrotunda, nitida, maculis minutis fuscis picta.

1083. LATHYRUS *Nissolia* pedunculis unifloris, foliis simplicibus, stipulis subulatis. *Linn. Sp. pl.* 2. p. 1029. — *Pölich. fl. pal. n.* 676.

Lathyrus pedunculis unifloris, foliis gramineis, stipulis minimis. *Hall. stirp. heb.* n. 441.

Nissolia vulgaris *Mapp. fl. Als.* p. 210. — *Lindern Tournef. Als. p.* 73. cum Icone

Lathyrus sylvestris minor. *Bauh. pin.* 344.

Germ. Nissolische Blatterbse.

Gall. Gesse Nissole.

Hab. Utrunque inter segetes passim. *Prop. Goldschier et Kehl* et alibi passim in Rheni vicinia. In Marggr. superiore inter segetes Weilenfes, ubi jam observavit *C. Bauhinus*. In Palatinatu circa *Lautern* in agris passim, prope *der langen Meil.* In Alsatiæ agris passim, prope *Erstein*, *Schweighausen* et alibi.

Flor. Junio, Julio. *Semina* Augusto matura. C

Caulis pedalis et altior, erectus, angulosus, glaber, simplex. *Folia* alterna, amplexicaulia, simplicia, lineari-lanceolata, integerrima, nervosa, glabra, erecta. *Stipulae* setaceae, minimæ. *Cirrhi* nulli. *Flores* axillares pedunculati. *Pedunculi* solitarii, uniflori, per raro biflori, longi. *Corollæ* purpureæ. *Legumina* longa, angusta, pendula, glabra, polysperma. *Semina* subrotunda, compressa.

††. *Pedunculis bifloris.*

084. **LATHYRUS** *hirsutus* pedunculis bifloris, cirrhis diphyllis, foliolis lineari-lanceolatis, leguminibus hirsutis, seminibus scabris. *Linn. Sp. pl. 2. p. 1032.* — *Pollich. fl. pal. n. 677.*

Lathyrus foliis geminis ellipticis, floribus solitariis et geminis, siliquis hirsutis. *Hall. stirp. helv. n. 437.*

Lathyrus angustifolius siliqua hirsuta. *Bauh. pin. 344.* — *Mapp. fl. Als. p. 165.*

erm. Haarige Platterbse.

Hall. Gessé velue.

Tab. Inter segetes passim rarius. In ditione Basileensi inter segetes versus *Mutenz*; in agris *auf dem Birsfeld*, inter segetes retro *Wartenberg* circa pontem *Wiesæ*. In Palatinatu in præfectura *Lutrensi* inter *Otterbach* et *Sambach* in agris t. *Pollichio*. In *Alsatia* in agris prope *Besfort t. Mappo*.

Flor. Junio, Julio. *Semina* Augusto matura. ☉

Caulis sesquipedalis et altior, erectus, annuato-membranaceus, ramosus. *Folia* alterna, petiolata, foliolis duobus lanceolatis, acuminatis, integris, trinerviis, subtus scabriusculis. *Stipulae* sagittatae, lanceolatae, integriusculae, aliquandoque denticulo instructae. *Flores* axillares, pedunculati. *Pedunculi* fulcati, scabri, pice biflori, rarius triflori, nutantes. *Bractae* curvae, subulatae, ad pedicellorum basin. *Corollae* pallide purpureae. *Legumina* longa, compressa, hirsuta, polysperma. *Semina* rotunda, rugosa, sublutea.

176 DIADFLPHIA. DECANDRIA.

Nota. Ad rariores floræ cives pertinet.

Obs. *Lathyrus odoratus* Linn. passim ob suavitatem in hortis colitur.

†††. *Pedunculis multifloris.*

1085. *LATHYRUS tuberosus* pedum multifloris, cirrhis diphyllis: foliolis libus, internodiis nudis. *Linn. Sp. p.* 1033. — *Pollich. fl. pl. n.* 678.

Lathyrus scapis multifloris, foliis ov capreolis trifidis. *Hall. stirp. helv. n.*

Lathyrus arvensis repens tuberosus. *B. pin.* 344. — *Mapp. fl. Alf. p.* 164.

Chamæbalanus. *Tabernæmont. p.* 891. *L*

Germ. Knollige Platterbse. Erdnüsse.

Gall. Gesse tubereuse.

Hab. In agris præsertim argilloso-lapideis calcareis montosis passim. Circa *Pforz Dillstein, Weissenstein, Hugenfeld* et in Marggr. superiori perinde infrequè In ditone Balileenfi inter *Spitelmata Holze* copiose; versus *Mutenz*, in monteteti agris; *auf den Galgenfeldern.* Brisgovia prope *Villingen* in agris careo-lapidosis. In Alsatiae agris perin prope *Niederhausbergen* et retro *Lingheim.* In Palatinatu circa *Lautern*; in *Mannweiler* et *Rockenhausen* in agris abunde, nec non circa Worms copiose *Pollichium* vidi.

Flor. Julio, Augusto. *Semina* Septembri, Octobri matura. *¶*

Radix tuberosa, nigricans. *Caules* cubiles, tripedales, adscendentes, erecti, quodque procumbentes, angulosi, nudi, glaberrimi
rami

Ns alterna, petiolata. Foliola bina, opposita, ovata, acuminata, aristata, glabra, laeteviridia. Cirrhi trifidi. Stipulae sagittatae, lanceolatae, integrae. Flores axillares, pedunculati. Pediculi 4-5-6 flori. Corollae amoenae, speciosae rubro-purpureae, aut roseae, suaviter odoratae. Legumina compressa, glabra, poly-

tabra suis gratissima. quibusdam in regionibus edula. Herba bonum pabulum pecoribus praebet.

Flores apibus mel largiuntur.

Aqua florum destillata, suaviter odorata. Gled. 2. p. 180.

Magnum saepe in agris montosis lapidoso-calcareo-fuligineis vitium, suis post cerasium messim optime succedendum.

Est Plantam florentem oculos mire delectantem, nullibi copiosius, quam in agris calcareo-lapidosis inter Villingen et Donaueschingen vis.

1086. *LATHYRUS pratensis* pedunculis multifloris, cirrhis diphyllis simplicissimis: foliolis lanceolatis. *Linn. Sp. pl. 2. p. 1033. — Pollich. fl. pal. n. 679.*

Lathyrus scapis multifloris, foliis lanceolatis, capreolis simplicibus. Hall. stirp. helv. n. 436.

Lathyrus sylvestris luteus foliis Viciae. Bauh. pin. 344. — Mapp. fl. Alf. p. 264.

Vicia. Tabernæmont. p. 890. Icon bona.

Icon. Curtis fl. Lond. t. 170. — Oeder. fl. Dan. t. 527.

Germ. Gelbe Wiesenplatterbse.

Gal. Gesse de prés.

578 DIADELPHIA. DECANDRIA.

Hab. Utrunque in pratis humidiusculis et cioribus, nec non in agris ac sepibus sive non infrequens.

Flor. Junio, Julio, Augusto. *Semina* Augusto Septembri matura. 7

Radix repens, albida. *Caulis* sesquipedalis, tripedalis et altior, debilis, ascensu saepe Cirrhis sustentatus, tunc erectus, et doque decumbens, angulosus, subpilosus. *Folia* alterna, petiolata, bina, opposita, lanceolata, trinervia, integra, subtulosa. *Petioli* trigoni, stipularum longitudo. *Cirrhi* simplices. *Stipulae* semisagittatae, coelatae, integrae, basi utrinque saepe ciculo hamato instructae. *Flores* axillares, bipedunculati. *Pedunculi* solitarii, angulosi, apice 8-10 flori. *Flores* erecti, race secundi. *Corollae* luteae. *Legumina* sesquialia, compressa, glabra, nigra, polysperma. *Semina* 7 — 12, subrotunda, glabra, perispermata fusca.

Egregium pecoribus in pratis pabulum praebet.

Nota. Planta omnino praesertim in pratis et vicinis, aliisque humidiusculis magis colenda — propaganda!

Flores apibus mel concedunt.

1087. *LATHYRUS sylvestris* pedalis multifloris, cirrhis diphyllis: foliis ensiformibus, internodiis membranaceis.
Linn Sp. pl. 2. p. 1033. — *Pollich fl.* n. 680.

Lathyrus caule alato, foliis geminis ensiformibus. *Hall. stirp. helv.* n. 434.

Lathyrus sylvestris major. *Bauh. pin.* *Mapp. fl. Alf.* p. 164.

fl. Dan. t. 327.

erm. Grofse Wald-Platterbfe.

hall. Gelfe de Bois.

hab. Utrunque in dumetis et fylvis paffim.

Amat vinearum marginalia dumetofa.

Flor. Junio, Julio, Augufto. Semina Augufto, Septembri matura. 7

Caulis fufertatus a dumetis, 5 - 6 - 7 pedibus, anceps, alatus, glabriufculus, ramofus. Folia alterna, petiolata. Foliola bina, opposita, lineari-lanceolata, integerrima, nervofa, fimplici. Cirrhi ramofi. Stipulae femihafatae, acuminae, integrae. Flores pedunculati. Pedunculi millares, angulofi, praelongi, folitarii, 5-7 flori. Corollae purpureae, rofeae, rarius albae, fpeciofae. Legumina longa, angulofa, glabra, polyfperma. Semina rotunda, fimplici.

Flores apibus mel concedunt.

Boaum pecoribus pabulum planta junior submini-

1088. *LATHYRUS latifolius* pedunculis multifloris, cirrhis diphyllis: foliolis lanceolatis, internodiis membranaceis. *Linn. Sp. pl. p. 1033.*

Lathyrus caulibus alatis, foliis ovato-lanceolatis, rigidis, fcapis multifloris. Hall. fimplici. helv. n. 433.

Lathyrus major latifolius, flore purpureo fpeciofior. Bauh. hift. 2. p. 302.

erm. Breitblaettrige Platterbfe.

hall. Gelfe à fêvilles larges.

hab. In ditione Bafileenfi in Monte Muteto et prope Binningen t. c. Bauhino.

186 **DIADELPHIA. DECANDRIA.**

Flor. Junio, Julio. *Semina* Augusto,
tembri matura. 7

Differt a praecedenti specie, *Caulibus*
floribus, *Foliis* ovatis, latioribus, superio-
rum lanceolatis, angustioribus, rigidioribus
Florum racemis 9-12 floris, *Corollis* m-
bus speciosioribus.

Egregium in hortis passim decus.

Flores apibus mel praebent.

Nota. LATHYRUM heterophyllum ped-
lis multifloris, cirrhis diphyllis tetraphyl-
liis foliolis lanceolatis, internodiis membra-
tis Lin n. frustra hucusque in terris Baden-
Palatinis, Alsaticis et confinibus quaesivi.
a C. Bauhino olim in ditione Basileensi
Klüber allegata, frustra a Cl. Lachenalic
que ibi hodie quaesita est.

1089. **LATHYRUS palustris** pedun-
culis multifloris cirrhis polyphyllis, fl-
liis lanceolatis. *Linn. Sp. pl.* 2. p. 103
Pollich. fl. pal. n. 681.

Lathyrus foliis senis, floribus racemosis
Hall. stirp. helv. n. 431.

Lathyrus palustris flore orobi nemo-
syni. *Rupp. fl. Jen.* 210. t. 1.

Lathyrus peregrinus, foliis viciae,
subcaeruleo pallideque purpurascens
Bauh. pin. 344.

Clymenum parisiense flore coeruleo. *Desf. inst.* — *Mapp. fl. Als.* p. 78.

Cicerula sylvestris. *Tabernaemont. p.*
Icon.

Icon. *Oeder. fl. Dan.* t. 399.

Ger m. *Sumpf-Platterbse.*

Gall. *Gesse des marais.*

DIADELPHIA. DECANDRIA. 181

l. In pratis pascuisque paludosis Rheno vicinis passim. Circa *Carlsruhe* prope *Eckenslein* et *Linckenheim* in pratis udis infrequens. In *Alsatia* *auf der Krafft* retro *Blobsheim* t. *Mappo*. In Palatinatu in pratis *gegen dem Holzhof* cum *Gratiola officinali* t. *Pollichio*. In Principatu superiore ad Lacum *Podamicum* in pascuis udis.

rr. Julio, Augustio. *Semina* Septembri maturantur. 3

Caulis sesquipedalis, bipedalis, erectus, ubique decumbens, anceps, debilis. glaberrimus, simpliciusculus. **Folia** alterna, petiolata, ovata. **Foliola** quatuor, sæpe sena, opposita, ovato-lanceolata, superiora angustiora, interna, venosa, glabra, apice aristata. **Cirrhi** nulli. **Stipulae** subulatæ, sæpe sagittatæ, eximie. **Flores** axillares. **Pedunculi** longi, solitarii, 6 flori. **Racemi** unilaterales. **Corollae** ex albo, violaceo-caeruleæ. **Legumina** compressa, glabra.

CCCXL. V I C I A.

Linn. Gen. pl. ed. Schreb. n. 1187.

Calyx monophyllus, tubulatus, erectus, aequinquefidus, acutus: *dentibus* superioribus brevioribus. **Corolla** papilionacea. **Vexillum** orbiculatum, ungue lato oblongo, apice emarginatum, cum acumine, lateribus reflexum, linea longitudinali compressa, elevata. **Alae** duæ, oblongæ, erectæ, semicordatæ, ungue oblongo, filis breviores. **Carina** ungue oblongo, bidentata, ventriculo compresso, semiorbiculato, brevior. **Filamenta** in plurimis diadelphea

Stigma obtusum, sub apice transverse barb-
tum. *Legumen* longum, coriaceum, uniloc-
ulare, bivalve, acumine terminatum. *Semi-*
plura, subrotunda, in quibusdam ovalia, com-
pressa.

† *Pedunculis elongatis.*

1090. *Vicia pisiformis* pedunculis mu-
tifloris, petiolis polyphyllis, foliolis ovi-
tis: infimis sessilibus. *Linn. Sp. pl.* 2, 1
1034. — *Pollich. fl. pal.* n. 682.

Vicia foliolis ovatis maximis, filiquis rac-
mosis pendulis. *Hall. stirp. helv.* n. 428.

Pisum sylvestre perenne. *Bauh. pin.* 343.

Pisum sylvestre. *Clus. hist.* 2. p. 129.

Icon. Jacqu. fl. Austr. t. 364.

Germ. Erbsenartige Wicke.

Gall. Vesce pisiforme.

Hab. In Palatinatu circa *Steinbach* in sylvarum
montosarum densis dumetis quibus in mon-
tem *Donnersberg* ascenditur, t. *Pollichio*
In Sponhemia prope *Winningen* ad Mo-
sellam in dumetis sylvaticis montosis den-
sis, nec non prope *Winterburg* vidi. In
Alsatia prope *Colmar*.

Flor. Junio, Julio. *Semina* Augusto matura. 2

Radix repens, prælonga, fusca. *Caulis*
tripedales, et altiores erecti. debiles, suban-
gulati, striati, glabri, ramosi. *Folia* alterna,
sessilia, apice cirrhosa. *Foliola* quatuor - sex ad
octo, subsessilia, opposita aut alternantia, infi-
ma cauli approximata, subrotundo-ovata, ob-
tusa, subpedicellata, integrifuscula, omnia gla-

hiscula, glauco-viridia, arista minuta terminata. **Stipulae** parvae, subrotundæ, acutæ, basi densimembratæ. **Flores** axillares, in pedunculis solitæ, **fulcatis**, erectis aut patentiusculis, glabris multifloris. **Corollæ** parvulæ, viridi-ochroleucæ, **friatæ**. **Vexillum** vix emarginatum. **Alae** angustæ, hamatæ. **Carina** fissa, rostro obtuso terminata. **Legumina** oblonga, compressa, glabra, pendula, unciam longa. **Semina** 3-4-6, subglobosa, glabra, fusca, demum nigricantia.

Ad rariores Floræ cives pertinet.

Planta egregium pecoribus pabulum præbet, facile in desertis locis nec non in cultis propaganda.

Flores apibus mel concedunt.

1091. **VICIA dumetorum** pedunculis multifloris, foliolis reflexis, ovatis mucronatis, stipulis subdentatis. *Linn. Sp. pl. 2. p. 1035.*

Vicia foliis ovatis, oblongis, stipulis ferratis, filiquis racemosis pendulis. *Hall. stirp. helv. n. 427.*

Vicia sylvatica maxima piso similis. *Bauh. hist. 2. p. 315.*

Cracca sylvatica. *Rivin. t. 50.*

Germ. Strauch-Wicke.

Ital. Vesce des Buiffons.

Tab. In Ditione Basileensi, in sylva *Hünigensi*, *Weilensi*, monte Muteto inter *Muttenz* et *Prattelen auf der Meyenfels* frequens. In Principatu Fürstenbergenli prope *Donaueschingen* et alibi passim in sylvis.

Flor. Junio, Julio. *Semina* Augusto matura. 7

Differt a præcedente specie, *Caulibus* crassioribus, altioribus, sæpe 6-7 pedalibus, alato-

184 DIADELPHIA. DECANDRIA.

membranaceis. *Foliis* longioribus, cirrho longiori, tripartito, diviso, terminatis. *Folia* semper sex, subalternis, ovalibus, (nec subterundo-ovatis) angustioribus, *Stipulis* in basi anguli folii foliaceis, maximis sagittatis, integris a foliolis remotis, amplexicaulibus, foliolis angulo duplo latioribus, mucronato-aristatis. *Corollis* longioribus in singulo racemo novem- et tredecim, purpureo-violaceis, (non ochroleucis). *Leguminibus* longioribus 8-9-spermis.

Nota. Planta in horto Botanico cum præcedenti culta, constantissima permanfit.

Ad rariores omnino floræ cives pertinet.

1092. *VICIA sylvatica* pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. *Linn. Sp. pl.* 2. p. 1035.

Vicia pedunculis multifloris, foliolis ellipticis, stipulis lunatis dentatis. *Smith. fl. brit.* 2. p. 768.

Vicia foliis ovatis, stipulis argute dentatis filiquis, racemosis pendulis. *Hall. stirp. helv.* n. 426. omnino.

Vicia multiflora maxima perennis tetro odore, floribus albis: lineis coeruleis notatis. *Pluk. Alm.* 387. t. 71. f. 1.

Icon. Oeder. fl. Dan. t. 277.

Germ. Waldwicke.

Gall. Vesce des Bois.

Hab. In ditione Basileensi in sylvis montosis circa *Meltingen* t. *Zeyhero*. In Principatu Fürstenbergensi prope *Unadingen* in præcipitiis calcareo-gypsaceis dumetosis cum *Spiræa Aruncus* nuper vidi.

Flor. Junio, Julio. *Semina* Augusto maturat. 1

Caulis cubitales et altiores, adscendentes, debiles, tetragoni, sulcato-friati, glabri, rarius ramosi. *Folia* alterna, petiolata, cirsioideum ramoso diviso terminata. *Foliola* in singulis foliis 12-14, alterna, subpedicellata, ovali-elliptica, omnia obtusa, integra, glabra, arista in apice terminata, semiunciam longa, tres lineas lata, laete viridia. *Stipulae* amplexicaules, parvae, argute sinuato-dentato-sciaceae. *Floris* axillares, penduli, in pedunculo parvis, friato, longo, racemosi. *Racemi* 9-13-
lori, unilaterales. *Corollae* albae, lineis punctisque coeruleis notatae, semiunciam longae. *Legumina* glabra, 7-8 lineas longa.

Nota. Foliola exacte ovali-elliptica, obtusa quandoque retusa, arista terminata, alterna 3-4 lineas a se invicem remota. Pedunculi axillares, solitarii, folio saepissime duplo longiores, in nostris saepe individuis.

Magnus omnino adhuc botanicorum dissensus in hac planta.

1093. *Vicia Gerardi* pedunculis multifloris, petiolis polyphyllis, foliolis ovali ellipticis, villosis, stipulis lanceolatis acutis integris appendiculatis.

Vicia pedunculis multifloris, petiolis polyphyllis, foliolis villosis, stipulis acutis integris appendiculatis. *Gerard. fl. gallopr. p.* 497. *fig.* 19. (*optima.*)

Vicia Cassubica Linn. *Sp. pl.* 2. *p.* 1035. — *Linn. ed. Willden.* 3. *p.* 1096.

Vicia multiflora pedunculis subduodecim floris racemosis, foliis inferioribus ovatis, supremis lanceolatis, retusis, inferne pubescentibus, stipulis integris, fliquis monospermis. *Pollich. fl. pal. n.* 683, *con. Jacqu. fl. Austr.* 3. *t.* 229.

185 DIADELPHIA. DECANDRIA.

Germ. Gerardische Wicke.

Gall. Vesce de Mr. Gerard.

Hab. In Palatinatu prope *Hartenburg* in collibus virgultosis gramineis post *Pollichium*.

Flor. Junio, Julio. *Semina* Augusto, Septembri matura. 7

Caules ex radice copiosi, sesquipedi tripedales, erecti, tetragoni, striati, villosuli, maxime ramosi. *Rami* alterni, patuli, erecti, cauli conformes. *Folia* alterna, sessilipinnata, cirrho ramoso terminata, praelobata. *Foliola* alterna oppositave, ovali-elliptica, tusa integra, villosula, subincana, quando late viridia, firmula, arista terminata, 26 superiora inferioribus sensim breviora et aristiflora. *Stipulae* lanceolato-acutae, integerrimae semisagittatae. *Flores* axillares, in pedunculo erecto solitario, folio subjacenti, breviori, locis racemosi, penduli, unilaterales, 11-16, imbricati. *Calyces* breves. *Corollae* violaceae. *Legumina* compressa, glabra, semiunciam longa, mono-rarius disperma. *Semina* rotunda, glabra, fusca,

Obs. Individua passim in Gallia meridionali et in Italia lecta, nostratibus simillima, magis tamen villosa-sericea. Individua in Franconia prope Langen in sylvis dumetosis montis Zatzberg cum *Perill.* Praefidi de Schreber lecta, modo recedunt a palatinis.

1094. *Vicia Cracca* pedunculis multifloris, folio longioribus, floribus locis racemosis imbricatis, pendulis, foliis lanceolatis obtusis acutisque, pubescentibus, stipulis semisagittatis, linearilanceolatis, integris, dentatisve.

Vicia Cracca pedunculis multifloris, foliis imbricatis, foliolis lanceolatis

DIADELPHIA. DECANDRIA. 187

bescentibus, stipulis integris. *Linn. Sp. pl.* 2. p. 1035. *Pollich. fl. pal.* n. 684.

Vicia foliis linearibus sericeis, racemis multifloris reflexis, stipulis integerrimis. *Hall. stirp. helv.* n.

Vicia multiflora. *Bauh. pin.* 345. — *Mapp. fl. Alf.* p. 125.

Vicia multiflora perennis nemorensis sive dumetorum. *Bauh. hist.* 2. p. 314.

Aracus. Tabernæmont. p. 891. *Icon.*

¶ *Vicia tenuifolia* pedunculis multifloris, floribus imbricatis foliolis linearibus trineruiis acuminatis, stipulis linearibus integris, caule erecto flexuoso. *Roth. fl. germ.* 2. p. 183. *Linn. Sp. pl. ed. Willden.* 3. p. 1099.

Cracca minor. Tabernæmont. p. 892. *Icon.*

γ. *Vicia polyphylla* foliis lineari-lanceolatis mucronatis villosis pedunculis axillaribus folio longioribus, floribus confertis secundis racemosis. *Desfont. fl. Atlant.* 2. p. 162. — *Linn. Sp. pl. ed. Willden.* 3. p. 1098.

Vicia orientalis multiflora incana angustissimo folio. *Tournef. inst. Cor.* 27.

Icon. a. Curtis fl. Lond. t. 310. — *Oeder. fl. Dan.* t. 804.

Germ. Vogelwicke.

Hall. Vesce multiflore.

Tab. α. et β. Utrunque inter segetes, ad viarum et fossarum marginalia dumetosa, in pratis, arvis, sylvis, collibus præsertim vinearum passim frequens. γ. in campis aridis fabulosis apricis sterilioribus.

188 DIADELPHIA. DECANDRIA.

Flor. Junio, Julio. *Semina* Augusto, Septembri matura. 7

Radix repens. *Caulis* cubitalis, tripedalis et longior, debilis pro ratione loci decumbens pro ratione loci in dumetis et frugibus erectus scandens, pubescens, villosus, quandoque glabriusculus, fragilis, simpliciusculus, saepe ramosus. *Folia* alterna, pinnata, cirrho tripedito terminata. *Foliola* alterna, opposita vel 24-26, oblongo-lanceolata, obtusa, quandoque acuta, semper aristata terminata, integra utrinque plus minusve pubescentia late viridi quandoque albide incana. *Stipulae* binae, semisagittatae, lanceolatae, acutae, integrae, saepe dentatae. *Flores* in pedunculis folio longioribus axillaribus, fulcatis, solitariis, racemosis. *Racemi* multiflori, suberecti, apice incurvi, demum reflexi, floribus 10-45 violaceis, violaceo rubellis, rarius albidis, quandoque pallidis incarnatis subimbricatis, brevissime pedicellatis, unilateralibus instructi. *Calyx* tubulatus coloratus, quinquedentatus, dentibus tribus inferioribus longioribus longe pilosis, medio longissimo, duobus superioribus minimis. *Corolla* *Vexillum* emarginatum, reflexum, coeruleo-violaceum, venis saturatioribus obsolete striatum. *Alae* conniventes, ungue bifidae. *Carina* rostrata obtusa, ungue bifida, albido-coerulea, apice macula saturate violacea utrinque notata. *Legmina* pendula, septem ad decem lineas longae compressa, glabra, fusco-flavicantia. *Semina* 4-5-6-8 ad 10, subrotunda, glabra, nigricantia

Planta omnino elegans, pro diverso solo nata altiore et demissiore. humidior et siccior argillaceo calcareo, siliceo-arenoso, sterili, variabilis, humilis et altior, glabriuscula, villosa, viridis, sericea, incarnata. *Foliola* latiuscula, ovali-lanceolata obtusa et angustiora lineari-lanceolata, acuta. *Stipulae*

DIADELPHIA. DECANDRIA. 189

modemque sæpe individuo integræ sæpe et dentatæ, lanceolatae, quandoque lineari-acutæ. Flores in nemo pauciores 10-13, et copiosiores 13-45. Corollæ majusculæ, longiores, et subbreviares, coeruleæ, violaceo-coeruleæ, incarnatæ et albidæ. Legumina septem lineas ad unciam longa.

Obs. Vicia tenuifoliam et polyphyllam esse hujus varietates, earum in horto cultura ne docuit, docuerunt idem variationes intermedie.

In Gallia meridionali nec non in sterilissimis campis Castiliæ novæ, legi Individua sericeo-albidescens, vix pedalia, fragilissima, rigida, Foliis lineari lanceolatis, acutis, utrinque sericeis, albido-micans instructa. Semina ex his collecta, in horto culta, exhibuerunt plantas multo majores, villosas, viridescens, minus fragiles, et altero jam anno exacte varietatem *a.* repræsentabant.

Egregium pecoribus in pratis et pascuis pabum præbet, hinc planta in pratis magis propaganda et colenda. Eleganter condecorat pratorum et sylvarum marginalia dumetosa.

Flores apibus melliferi.

1095. *Vicia villosa* radice annua, pedunculis folio longioribus longe pilosis floribus imbricatis, foliolis ovali-lanceolatis obtusis, utrinque villosis, stipulis femi sagittatis basi dentatis.

Vicia villosa pedunculis multifloris, floribus imbricatis, foliolis ovalibus, villosis, stipulis nervosis basi dentatis. *Roth. fl. germ. 2. p. 182.*

Term. Einjährige Vogelwicke.

Ball. Vesce multiflore annuelle.

Hab. Passim in agris locisque limosis cum præcedente. In agris inundatis prope *Dachslund, Knielingen, Linckenheim* et alibi.

192 DIADELPHIA. DECANDRIA.

1098. *Vicia lathyroides* leguminifera
 sessilibus solitariis, erectis glabris, fo-
 lis fenis: inferioribus obcordatis. *L.*
Sp. pl. 2. p. 1037. — *Pollich. fl. pal. n. 6*

Vicia montana minima, flosculo purpureo
Linn. Amoen. acad. 1. p. 167.

Vicia praecox verna, minima solonien-
 sino hexaedro. *Morif. hort. blaef.* 3
Hist. 2. p. 63.

Vicia minima praecox Parisienfium. *Ton-*
nes. inf. p. 397.

Icon. Oeder. fl. Dan. t. 58.

Germ. Platterbsenartige Wicke. Kleinste V-
cke. Kleine Frühwicke.

Gall. Vesce Gessier.

Hab. Circa *Carlsruhe* prope *Rintheim* ju-
 dumeta in graminosis passim; prope *Mü-*
burg am Lutherischen Wäldlein ad
 meta non infrequens. In Margraviatu
 superiore inter pontem *Wiesæ* et *Klein-*
ningen ad dextram in prato sicco. In
 latinitate inter *Hartenburg* et *Dürkheim*
 gramineis locis ad radicem montis ill-
 cujus summitati monasterium *limburches-*
 inaedificatum est. t. *Pollichio.* Amat lo-
 aprica graminea dumetosa.

Flor. Martio, Aprili. *Semina* Majo matura.

Radix fibrosa, tenuis, albida. *Caulis* di-
 talis, palmaris, adscendens, aut decum-
 debilis, angulosus, simpliciusculus, ad ba-
 quandoque ramosus. *Folia* alterna, pinnat-
 2-saepe 3 juga, inferiora breviora, latior-
 cuneiformi-obovata, retusa, emarginata,
 lobula, integra, superiora lanceolata, obtu-
 emarginata, longiora, et angustiora. *Cirrh-*
 foliu

~~Inter~~terminans, simplex, brevis, quandoque
~~Stipulae~~ *Stipulae* parvae, semisagittatae. Flo-
~~solitarius~~ *solitarius*, subseffiles. *Corolla* par-
~~purpureo-coerulea~~ *purpureo-coerulea*. *Legumen* erectum, lan-
~~compressum~~ *compressum*, unciale, 5-7 spermum.
~~rotunda~~ *rotunda*, fusco-viridia, nigro minutim
~~punctata~~ *punctata*, parva.

Minimas Viciarum species pertinet, quae facile
~~subducit~~ *subducit*.

1000. *VICIA lutea* leguminibus seffilibus
reflexis, pilosis solitariis pentaspermis,
corollae vexillis glabris. *Linn. Sp. pl. 2.*
p. 1037. — Pollich. fl. pal. n. 687.

Vicia flore luteo pallido, filiquis hirsutis cras-
sis brevibus perpendentibus. *Morif. hist.*
2. p. 62. Sect. 2. t. 21. f. 5.

Vicia leguminibus solitariis deorsum fixis
hirsutis. *Sauv. Monsp. p. 235.*

Vicia sylvestris lutea filiqua hirsuta. *Bauh.*
pin. p. 345.

Germ. Gelbe Wicke.

Call. Vesce jaune.

Lab. In dumetis ad vias et agros inter *Darm-*
satt et *Frankfurt* non infrequens, post
Pollichium ibi, nec non prope *Frankfurt*
passim vidi.

Flor. Junio, Julio. *Semina* Augusto maturat. ☉

Caulis sesquipedalis, cubitalis, debilis, de-

foliens, sæpe erectus, angulosus, glabriuscu-

Folia alterna, pinnata, cirrhis bifidis ter-

Foliola in singulo folio opposita, alter-

ne, 10-12, subpedicellata, lineari lanceo-

la, obtusa, arista albida terminata, integerr-

ima subvillosa. *Stipulae* cordatae, denticulatae,

194 DIADELPHIA. DECANDRIA.

supremæ integriusculæ, *Flores* axillares, sessiles, solitarii, quandoque bini, erecti. *rolla* speciosa, pallide flava. *Vexillum* fundum, obtusum, apice emarginatum. conniventes, carina breviores. *Carina* ubifida, obtusa, ad apicem macula nigra prædita. *Stamina Filamenta* decem, co *Stigma* plumosum. *Legumina* lanceolata, preffa, hirsuta, deorium pendula, polyspe *Semina* rotunda, fusca, nigro punctata.

Nota. Ad rariores omnino in Germania cives net. copiose in calidioribus Galliae Hispani Italiæ agris, et olivetis obveniens.

1100. *Vicia sepium* leguminibus cellatis, subquaternis erectis, foliolis integerrimis: exterioribus decrel tibus. *Linn. Sp. pl.* 2. p. 1038. — *lich. fl. pal.* n. 688.

Vicia foliis ovatis oblongis, scapis brev quadrifloris. *Hall. stirp. helv.* n. 429

Vicia sepium folio rotundiore acuto, ser nigro. *Bauh pin.* 345. — *Mupp. fl.* p. 325. et *Vicia maxima dumetorum* (nec *C. Bauhini*.)

Cracca major. Tabernamont. p. 892. 1

Icon. Oeder. fl. Dan. t. 699.

Germ. Heckenwicke.

Gall. Vescé des haies.

Hab. Utrunque ad sepes, viarum margin: dumetosa, prata, nec non in sylaticis minolis passim abunde.

Flor. Aprili, Majo, Junio. *Semina* Julio, gusio matura. 7

repens. *Caulis* sesquipedalis, cubi-
 latus, pro ratione loci decumbens aut erectus,
 trigonus, sulcatus, simpliciusculus. *Folia* al-
 ternata, pinnata, cirrho trifido terminata. *Foliola*
 bipedunculata, opposita, subalternave, 8-12,
 ovata, obtusa, ariliata, integerrima, læte viri-
 da, glabra, margine villulosiuscula, inferiora
 superioribus majora, superiora sensim decre-
 scentia. *Stipulae* subrotundæ, denticulatæ, par-
 vae, utrinque macula fusca notatae. *Flores* axil-
 lares, breviter pedunculati 2-4-6 flori. *Ra-
 cemi* nutantes. *Corollae* albido-violaceo pur-
 pureae, aut incarnatae. *Filamenta* decem, coa-
 ctæ. *Legumina* tereti-compressiuscula, pen-
 data, maturitate erecta, glabra, unciam circi-
 ter longa. *Semina* 4-5-6, suborbiculata, fusca,
 maculis nigris picta, glabra.

Egregium pecoribus planta pabulum subministrat.

Flores apibus mel concedunt.

1101. *Vicia Faba* leguminibus subseffili-
 bus ternis torulosis, foliolis ovatis inte-
 gerrimis, petiolis absque cirrhis, stipu-
 lis sagittatis basi dentatis. *Willden. Linn.
 Sp. pl. 2. p. 1111.*

Vicia Faba caule erecto, petiolis absque
 cirrhis. *Linn. Sp. pl. 3. p. 1039.*

Faba. Bauh. pin. 338.

Faba minor f. *equina.*

Icon. Regnault. bot.

Germ. Saubohne. Ackerbohne.

Call. Fève des marais.

*Lat. In Persiæ confiniis non procul a mari
 Calpio. Colitur passim frequens. ..*

196 DIADELPHIA. DECANDRIA.

Flor. Junio, Julio. *Semina* Augusto, Septem-
bri matura. ☉

Corollæ speciosæ, suaveolentes, albi
Alæ in medio macula atra holosericea. *Semina*
maxima reniformia, compressa, alba, in var-
tione β . minora.

Pharm. *FABÆ Stipites; Flores; Semina.*

Stipites aut caules, incineratione magnam copi-
salis alcalini exhibent.

Flores recentes gratum odorem spirant, qui ve-
ficcatione perit.

Farina Seminum additur Farinis nomine qui
tuor resolventium notis.

Legumina cum feminibus immatura, tenella in-
linis nota, delicatiora, quam ex majori variatione.

Flores apibus mel præbent.

CCCXLI. ERVUM.

Linn. Gen. pl. ed. Schreb. n. 1189.

Calyx quinquepartitus longitudine corollæ
laciniis linearibus, acuminatis, subaequalibus
Corolla papilionacea. *Vexillum* planum, sul-
rotundum, majus. *Alæ* obtusæ, vexillo dimidi
breviores. *Carina* alis brevior, acuminata. *Sti-*
mina Filamenta diadelpha. Stigma obtusum
imberbe! (qua nota a Viciis differt.) *Legume*
oblongum, obtusum, teres, feminibus pro-
berantibus nodosum. *Semina* 2-4 subrotunda

1102. *ERVUM tetraspermum* pedun-
culis subbifloris, feminibus globosis qua-
ternis. *Linn. Sp. pl. 2. p. 1039.* — *Poi-*
lich. fl. pal. n. 689.

DIADELPHIA. DECANDRIA. 197

cia foliis linearibus filiquis gemellis glabris. *Hall. stirp. helv. n. 423.*

cia minima, cum filiquis glabris. *Mapp. fl. Alf. p. 325.*

. *Curtis. fl. Lond. — Schkuhr. bot. t. 202.*

1. *Vierfaamige Feldwicke.*

! *Ers tetrasperme.*

Utrunque in arvis, cultis, agris, vineis assim frequens.

1. Junio, Julio. *Semina* Julio, Augusto natura. ☉

Aules sesquipedales, cubitales, adscens, tetragoni, glabri. *Folia* alterna, pinnatifido ramoso terminata. *Foliola* 6-8-pinnifida, rarius 10, opposita, subalternave, lanceolata-linearia, integra, apice aristata, glaberrima. *Stipulae* sagittatae, integrae, acutae. *Flores* res, pedunculati. *Pedunculi* solitarii, longe foliorum, biflori, rarius uniflori, subtetraspermi. *Corollae* parvae, albae, venis purpureis pictae. *Legumina* teretiuscula, glaberrima subtetrasperma. *Semina* subrotunda, fusca, dura, nigro punctulata.

Semina avibus accepta. Planta haec cum sequente ovibus pabulum in arvis subministrat.

3. **ERVUM** *hirsutum* pedunculis multifloris, seminibus globosis binis. *Linn. Sp. pl. 2. p. 1039. — Pollich. fl. pal. n. 690.*

icia foliis linearibus, filiquis racemosis dispersis hirsutis. *Hall. stirp. helv. n. 690.*

icia segetum cum filiquis plurimis hirsutis. *Bauh. pin. 345. — Mapp. fl. Alf. p. 325.*

198 DIADELPHIA. DECANDRIA.

Icon. Oeder. fl. Dan. t. 639. — Curtis. fl. Ló
— *Schkuhr. bot. t. 202.*

Germ. Kleine rauhe Feldwicke.

Gall. Ers velue.

Hab. Utrinque in cultis, agris, hortis, vi
passim frequens.

Flor. Junio, Julio. *Semina* Julio, Augi
matura. ☉

Differt a praecedenti specie, *Foliolis* 14
lineari-lanceolatis, truncatis, arista terminatis
Pedunculis 3-4-8 floris, *Leguminibus* hirsutis
dispermis.

1104. *ERYUM Lens* pedunculis subbi
ris, seminibus compressis convexis. *L.*
Sp. pl. 2. p. 1039.

Lens. *Hall. stirp. helv. n. 421. — Rivin. t.*
t. 35.

Icon. Schkuhr. bot. handb. t. 202.

Germ. Linse.

Gall. Lentille.

Hab. Utrinque inter segetes, nec non in vir
quasi spontanea. Colitur passim aut sola
cum cerealibus praesertim *Hordeo*, sat

Flor. Junio, Julio. *Semina* Augusto matura.

Pharm. LENTILIUM Semen.

Variationes semine majori et minori. Plu
notissima. Semina omni tempore pro cibo d
habita, aegre concoquuntur.

CCCXLII. C Y T I S U S.

Linn. Gen. plant. ed. Schreb. n. 1101.

Calyx monophyllus, campanulatus, bre
vissi obtusus: ore bilabiato: labio superi

hirsuto, acuminato; inferiore tridentato. *Corolla* papilionacea. *Vexillum* ovatum, assurgens, lateribus reflexis. *Alæ* longitudine vexilli, rectæ, obtusæ. *Carina* ventricosa, acuminata. *Stamina* Filamenta diadelphea. *Stigma* obtusum. *Legumen* oblongum, obtusum, basi attenuatum, rigidum. *Semina* nonnulla, reniformia, compressa.

1105. *CYTISUS nigricans* racemis simplicibus erectis, foliolis ovato-oblongis. *Linn. Sp. pl.* 2. p. 1041.

Cytisus foliis ovatis nitidis, floribus spicatis, cernuis, calycibus et siliquis sericeis. *Hall. stirp. helv.* n. 361.

Cytisus glaber nigricans. *Bauh. pin.* 390.

Cytisus quartus. *Clus. hist.* 1. p. 95.

Icon. Jacqu. fl. Austr. t. 387.

Germ. Schwärzlichte Bohnenstaude. Schwärzlichter Cytisus.

Gall. Cylise noir.

Hab. In Principatu superiore versus Lacum Bodamicum prope Ueberlingen in collibus cum *Arbuto uva ursi* frequens.

Flor. Julio et Augusto. *Semina* Septembri, Octobri matura. 5

Frutex 2-3 pedalis et altior. *Rami* alterni, erecti, virgati, villosi. *Folia* alterna, petiolata, ternata. *Foliola* ovato-oblonga, integra, inferne villosa, nitida. *Racemi* terminales, prælongi, speciosi, erecti. *Corollæ* luteæ. *Legumina* sericea.

Siccitate tota nigrescit.

200 DIADELPHIA. DECANDRIA.

Ob racemos virgatos praelongos erectos, flo-
que copiosos luteo - aureos, insigne hortorum c-
mentum.

Flores apibus mel præbent.

Ad rariores omnino Floræ cives pertinet.

Nota. In hortis anglicis Carlsruhanis adhuc h-
tantur sequentes species: *Cytisus lat-*
num, sessilifolius, hirsutus, cag-
tus, austriacus, supinus. Linn. et
pureus. Scopoli fl. Carn.

CCCXLVIII. R O B I N I A.

Linn. Gen. plant. ed. Schreb. n. 1195.

Calyx monophyllus, campanulatus,
drifidus; *denticulis tribus inferioribus* ten-
ribus; supremo duplo latiori vix emargi-
Corolla papilionacea. *Vexillum* subrotund-
majus, patens, obtusum. *Alae* oblongæ, ov-
liberae, appendice brevissimo, obtuso. *Ca-*
fere semiorbiculata, compressa, obtusa, lo-
titudine alarum. *Stamina Filamenta* diadel
Stigma villosum antice ad apicem styli. *L-*
men compressum, gibbum, longum. *Semina*
niformia.

1106. ROBINIA *Pseud-Acacia* r-
mis pedicellis unifloris, foliis impari-
natis, stipulis spinosis. *Linn. Sp. pl. :*
1043.

Robinia aculeis geminatis. Linn. hort. C
354. *Duham. arb. 2. t. 42.*

Icon. Wangenheim amer. t. 7. f. 19.

Germ. Gemeine Robinie. Unächter Acac-
baum.

Gall. *faux Acacia épineux.*

Hab. In Virginia. Hospitatur utrinque passim in hortis, ambulacris, ad vineas, nec non in sylvis passim.

Flor. Majo, Junio. *Semina* Augusto matura. 5

Arbor pulchra, florens suaveolens, citissime crescit, præsertim in montosis apricis. Lignum durum exquillum. Folia bonum pabulum pecoribus. Flores apibus melliferi.

Nota. In hortis Carlsruhanis et Schwezingensibus adhuc hospitantur, *ROBINIA* Pseud-Acacia var. inermis. *Robinia hispida*. Linn. (*Acacia Rose* Gall.) *Caragana*. frutescens. pygmaea. *Altagana*, ferox, *Halodendron*, *Chamlagu*, inermis, et viscosa. *Ventenat. descr. pl. nov. 4. t. 4*

CCCXLIV. COLUTEA.

Linn. Gen. pl. ed Schreb. n. 1196.

Calyx monophyllus, campanulatus, quinquefidus, erectus, subæqualis. *Corolla* papilionacea. *Filamenta* diadelpha. *Stigma* linea barata, a medio styli ad ejus apicem, a parte superiore extensum. *Legumen* maximum, latissimum, inflatum, membranaceo-diaphanum! sutura superiore erecta, inferiori gibba, uniloculare, ad basin sutura superiore dehiscens.

107. *COLUTEA arborescens* arborea, foliolis obcordatis. *Linn. Sp. pl. 2. p. 1045.*

Colutea foliis ovatis emarginatis, floribus racemosis. *Hall. stirp. helv. n. 400.*

Colutea vesicaria. Bauh. pin. 396.

202 **DIADELPHIA. DECANDRFA.**

Icon. Pfleger arb. et frut. Badenf. Regni bot.

Germ. Gelber Blasenstrauch. Falsche Se Gall. Baquenaudier. Faux Séné.

Hab: In Brisgovia am Kaiserstuhl inter Salem et Burgheim abunde cum Generosa Itz vidi. In Marggr. superiori, in monte Schwärze retro Oberweiler copiose. In cipatu superiore prope Salem et alibi pa

Flor. Majo, Junio. *Semina* Augusto maturi

Frutex elegans, 4-8 pedalis, erectus. *lia* alterna, impari-pinnata. *Foliola* obscurae laete viridia, integra. *Stipulae* lanceolatae. *res* racemosi. *Racemi* folio sub breviores, flori, nutantes. *Corollae* luteae. *Legumina* flata, velicam referentia, glabra.

Pharm. COLUTEAE Folia.

Folia amariuscula, subnauseosa, alvum mo-
Semina ad drachmam ad duas sumpta, vomitum
movent. Prevot. med. paup. p. 511.

Nota. In hortis Carlsruhanis et Schwetzingen
aliisque passim anglicis adhuc hospitantur.
LUTEA cruenta, fruticosa, foliolis o-
datis. glaucis, crassiusculis, corollis cru-
seu C. orientalis (quae in Germania au-
vix unquam spontanea venit.) et COLU-
Pocockii Ait. hort. Kew. 3. p. 55
halepica. Lamarck Encycl. 1. p. 34

CCCXLV. C O R O N I L L A.

Linn. Gen. pl. ed. Schreb. n. 1198.

Umbellula simplex. *Calyx* monophyl-
brevissimus, compressus, bifidus, erectus;
tibus tribus inferioribus minoribus: supe-
bus duobus coalitis. *Corolla* papilionacea.

stylum cordatum, undique reflexum, alis vix longius. *Alae* ovatae, sursum conniventes, deorsum deliscentes, obtusae. *Carina* compressa, acuminata, adscendens, alis saepius brevior. *Filamenta* diadelpha. *Legumen* longissimum, teres, inter singula semina isilumis contractum, uniloculare, bivalve, articulatim discedens. *Semina* plura.

Nota. C. Emerus Ungues corollae calyce longiores. Legumen tenue subulatum.

1108. **CORONILLA** *Emerus* fruticosa, pedunculis subtrifloris, corollarum unguibus calyce triplo longioribus caule angulato. *Linn. Sp. pl.* 2. p. 1046.

Coronilla frutescens, foliis retusis, septenis, siliquis longissimis, vix articulatis. *Hall. stirp. helv. n.* 389.

Colutea siliquosa f. *Scorpioides* major. *Bauh. pin.* 397.

Emerus *Caesalp.* 117. — *Mapp. fl. Alf. p.* 97.

Emerus minor. *Tournef.* — *Mapp. fl. Alf.* 98.

Colutea siliquosa minor. *Bauh. pin.* 397.

Leon. Miller. Dict. Icon. t. 132. f. 1. major. fig. 2. minor. *Pfleger arb. et frut. Badens.*

German. Scorpions-Peltschen.

Gall. Coronille pauciflore.

Tab. In Brisgovia in dumetis am *Kaiserstuhl* ad rudera arcis *Sponeck*, cum *Ruta graveolente*. In Marggr. superiore supra *Kembs*. t. *Vulpio*, nec non in monte prope *Krenzach*. In Ditione Basileensi in monte Muteto ad radicem Jurae passim frequens. In

Principatu superiore prope *Salem* et in sylvis et dumetis, frequens. In Alsatia rudera antiqua am- *Dannbacher* et prope *dem Blinschweiler Schloss*, copiose t. *Mg* 8. In Alsatia auf dem *Halgensteiner Falkenstein* Hochgebürg, non prope *Blinschweiler*, nec non in montosis siccis agris prope *Litzelburg* t. *Mapp* Flor. Aprili, Majo, Semina Julio, matur.

Frutex 3-6 pedalis, erectus. *Folia* magna, pinnata. *Foliola* septem obovato-cuneata, obtusa, submarginata, integra. *Fl* subtriflori. *Corollæ* luteæ. *Ungues* petalo calyce triplo fere longiores. *Vexillum* a reliis petalis remotum, obcordatum, basi lineat. *Carina* ampla, rostrata. *Legumina* præcylindrico-subulata, tenuia, subarticulata, lysperma.

Pharm. COLUTEÆ SCORPIOIDIS *Folia*

Folia amariuscule, laxantia, hodie in medicina obleta.

Frutex elegans. Hortorum ornamentum. *Folia* apibus mel præbent.

1109. *CORONILLA coronata* fruticosa
foliis novenis obovatis: foliolis inter
caule approximatis, stipulis oppositis
bipartitis. *Linn. Sp. pl. p.* 1047.

Coronilla caule erecto, foliis undenis cuneatis, floribus umbellatis, filiquis articatis pendulis. *Hall. stirp. helv. n.* 388.

Coronilla montana. *Scop. fl. Carn. ed.* 1912. t. 44.

Colutea scorpioides minor coronata. *Berol. pin.* 397.

Colutea scorpioides secunda. *Clus. hist.* 1. p. 98.

Jacqu. fl. Austr. t. 95.

Gekrönte Peltfchen.

Coronille couronné.

In ditione Basileensi, in monte *Sissach* apud, in montibus *Farnsburg* et *Sonnenberg* prope *Meispirach* inter montem *Mutetum* et *Schauenburg* frequens.

Junio, Julio. *Semina* Augusto, maturat. *Suffrutex.*

Radix lignosa, longa, ramosa, petrarum fissi-
petens, difficile hinc eradicanda, luteo-
ca. *Caules* ex radice plures, pedales, sesqui-
pedales, adscendentes, erecti, teretiusculi, fir-
mif, versus radicem sublignosi, superne her-
mif, glabri. *Folia* alterna, pinnata cum im-
mif *Foliola* sublellilia, opposita, alternave
ad undecim, ovalia, obtusa, integra,
terminata, glauca, glabra, septem ad oc-
tas longa, semiunciam lata, superiora
minora. *Stipula* lata bilida, quandoque di-
stilla. *Flores* in longis pedunculis axillaribus
natis, erectis, dense coronato-umbellati,
buli. *Corolla* luteæ. *Legumina* pendula, cy-
lica, articulata, subulato-acuminata, inæ-
qua, unciam ad sesquiunciam fere longa,
pendula, 3-4 sperma. *Semina* oblonga,

Ad rariores Florae cives omnino pertinet.

Flores suaveolentes, apibus mel subministrant.

III. *CORONILLA minima* suffruticosa,
procumbens, foliolis novenis ovatis, sti-
pula oppositifolia emarginata, legumini-
bus angulatis nodosis. *Linn. Mant.* 444,

206 DIADELPHIA. DECANDRIA.

Coronilla caulibus lignosis procumbentibus
foliis ovatis, floribus umbellatis, filis
alatis. *Hall. stirp. helv. n. 390.*

Polygonum Cortusi. *Bauh. hist. 2. p. 351*

Lotus enneaphyllos. *Dalechamp. hist. 1.*

Icon. Jacqu. fl. Austr. t. 271.

Germ. Kleinste Peltchen.

Gall. Coronille mineur.

Hab. In ditione Basileensi, in montis *V
berg* cacumine *Paswang* dicto. *t. Laci
lio* nec non *auf der Kallensflur* prope *Re
ptingen* *t. Zeyhero.*

Flor. Junio, Julio. *Semina* Augusto matura
Suffrutes.

Radix ramosa, lignosa, extus fusca,
albida. *Caules* ex radice plures, digitales,
mares, inferne lignosi, superne herbacei
mosi, prostrati. *Folia* alterna, pinnata cui
pari, opposita alternave, subpedicellata, tri
na, quandoque novena, obovata, obtusa, tri
doque emarginata, aristia terminata, inter
crassiuscula, avenia, glauca, glabra, parva
ad quantuor lineas longa *Stipula* mem
bracea, pallida, decidua, connivens, super
fida. *Flores* in pedunculis nudis, solitariis
minalibus, umbellati, 5-6- saepe 7 rarius
Corollae luteae. *Legumina* subtetragona,
culata, fusca, pendula, unciam circiter longi
4-6 sperma. *Semina* oblonga.

Obs. Individua nostratia omni ex parte conveniunt
specimine a Cl. Jacquino depicto et descri

Specimina in Arragoniae montosis rupestribus
me lecta, nostratibus minora, Foliolis tantum
tem ad novem minoribus instructa sunt. Idem
servavi circa individua in Pyrenaeis nec non
vetis alpinis lecta.

III. *CORONILLA Securidaca* herbacea, leguminibus falcato-gladiatis, foliolis plurimis. *Linn. Sp. pl. 2. p. 1048.*

Securidaca lutea major. Bauh. pin. 398.

Ger. Beilkrautartige Peltfchen.

Gall. Coronille Securidaca.

Hab. Prope Carlsruhe am Holzmagazin, vix spontanea, sed ex horto botanico dissecta. In Margraviatu superiore in monte Krenschach nec non in monte Muteto t. C. Bauhino, hodie tamen exul.

Flor. Junio, Julio. Semina Augusto matura. ☉

Caulis sesquipedalis et altior, erectus, angulosus, sulcatus, glaber, ramosus. *Folia* alternata, pinnata cum impari. *Foliola* opposita, ovata breviter pedicellata, oblonga, obtusa, apice breviter submarginata, subaristata, integra, glabra. *læte viridia*, 13-15, unciam longiusculam lata, impari obovato-cuneiformi, reliquis magis emarginato. *Flores* axillares, pedunculis praelongis, solitariis, falcato-angulatis, glabris, erectis, umbellati, 7-9 flori. *Corolla* luteæ. *Legumina* compressa, ensiformia, articulata, glabra, fusca, erecta, tres ad quatuor uncias longa, apice, aristâ incurva præfixa, polysperma.

Hab. Plantam abundanter præsertim in Castiliarum regionibus inter segetes vidi.

bonum præbet pecoribus pabulum, hinc omnino ære solo propaganda et colenda.

III2. *CORONILLA varia* herbacea, leguminibus erectis teretibus, torosis, numerosis, foliolis plurimis glabris. *Linn. Sp. pl. 2. p. 1048. — Pollich. fl. pl. n. 691.*

208 DIADELPHIA. DECANDRIA.

*Coronilla caule brachiato, angulato, fr
vigenis aristatis, floribus umbellatis. Fl
stirp. helv. n. 387.*

*Coronilla herbacea, flore vario. Mapp
Alf. p. 82.*

*Securidaca dumetorum major, flore va
siliquis articulatis. Bauh. pin. 349.*

Icon. Schkuhr. bot. Handb. t. 205.

Germ. Bunte Peltchen. Bunte Kronwicken

Gall. Coronille bigarré.

*Hab. Utrunque in agris, arvis, vineis ad vias
marginalia dumetosa, et prata passim :
infrequens.*

Flor. Junio, Julio. Semina Augusto maturi

*Radix repens, ramosa. Caules ex rad
plures, cubitales, tripedales, decumbentes,
cati, glabri, herbacei, ramoli. Rami alte
Folia alterna, impari-pinnata. Foliola 15-
19, opposita, alternave pedicellata, obov
obtusa, integra, arista terminata, glabra,
ferne glauco-viridia. Stipulae binae, line
acutæ, parvæ. Flores axillares, in pedunc
sulcatis nudis, solitariis, folio duplo longi
bus, coronato-umbellati, 17-19-21, ere
pedicellati, denum deflexi. Corollae amo
rosæ, pallide purpureae, incarnatæ, quar
que fere albidæ. Legumina linearia, subul
pentagona, geniculato-articulata, glabra,
quiunciam ad binas uncias longa, polysper
Semina cylindrica, utrinque obtusa, glabra*

*Planta florens pulchella. Bonum ovibus publi
præbet.*

Flores apibus mel præbent.

CCCXLVI. ORNITHOPUS.

Linn. Gen. pl. ed. Schreb. n. 1199.

Umbella simplex. Calyx monophyllus, tubulosus: ore quinquedentato, subæquali. Corolla papilionacea. Vexillum obcordatum, imbricatum. Alae ovatae, rectæ, magnitudine vix æquales. Carina compressa, minima. Filamenta diadelphica. Legumen subulatum, teres, arcuatum, articulatum, isthmis interceptum, articulis discedens. Semina subrotunda.

3. ORNITHOPUS *perpusillus* foliis

pinnatis, leguminibus subarcuatis. Linn.

Sp. pl. 2. p. 1049. — Pollich. fl. pal. n. 692.

Ornithopodium caule prostrato, foliis pinnatis, filiquis umbellatis, articulatis. Hall. stirp. helv. n. 393.

Ornithopodium minus. Bauh. pin. 350. — Mapp. fl. Als. p. 226.

Ornithopodium majus. Bauh. pin. 350. Mapp. fl. Als. p. 226.

Ornithopus intermedius caulibus repentibus longissimis, foliis pinnatis, bractea pinnata, leguminibus incurvatis majore. Roth. fl. germ. 2. p. 215.

Leon. Oeder. fl. Dan. t. 730. — Schkuhr. bot. Handb. t. 206.

Germ. Kleiner Vogelfus.

Gall. Pied - d'oiseau delicat.

Ital. In aridis fabulosis sterilibus circa Carlsruhe, Dachsland inter et Mühlburg, et alibi passim frequens. In Marggr. superiori circa Weil et Haltingen. In Palatinatu prope Schwetzingen in sylvis fabulosis; circa Lautern in arvis et ericetis arenosis ubique

210 **DIADELPHIA. DECANDRIA.**

t. Pollichio. In Alfatia prope *Hagen* sterilibus. β . promiscue iisdem in locis *Carlsruhe*, et praesertim in ericetosis *am Hardthof* abunde.

Flor. Junio, Julio. *Semina* Julio, Autumnata. ☉

Caules ex radice plures, unciales, ciales, digitales, decumbentes, subangulololi, in variatione β . palmares, pedales et giores. *Folia* alterna, impari pinnata. *Fo* opposita, alternave, 11-25, sessilia, ovalia tusa, arista vix manifesta terminata, inter utrinque villosa-sericea. *Stipulae* exiguae bulatae. *Flores* 2-3-4-5, umbellati, termin sessiles, subtus *Bractea* foliacea, longiore, nata, suffulti. *Corollae* minutae, albido-lut venis purpureis praesertim ad vexillum p *Legumina* arcuata, articulata, 7-8 sperma

O b f. Individua passim in Gallia meridionali et Hil lecta, magis splendida, tota villosa-sericea, rum nostratibus similia.

In horto botanico culta, major evadit. Va β . minime pro peculiari specie declaranda. Bonum ovibus pabulum praebet.

CCCXLVII. HIPPOCREPIS

Linn. Gen. pl. ed. Schreb. n. 1200.

Umbella simplex. *Calyx* monophyllus, c quedentatus: *denticulis* superioribus conjur minusque divis. *Corolla* papilionacea. *V lum* cordatum, ungui longitudine calycis dens. *Alae* ovato-oblongae, obtusae. *Ca lunulata*, compressa. *Filamenta* diadelphagumen (*Soleae* equinae forma) compresso-m

lanceum, longum, incurvatum futura al-
tius ad superiorem incisum, multoties sinu-
ato subrotundis; hinc *articulis* confians obtuse
sinuatis, plurimis, futura superiore connexis.
Semen solitaria in singulo articulo, oblonga,
incurva.

III. *HIPPOCREPIS comosa* legumi-
nibus pedunculatis confertis, margine ex-
teriore repandis. *Linn Sp. pl.* 2. p. 1050. —
Pollich. fl. pal. n. 693.

Ferum equinum filiquis umbellatis undula-
tis. *Hall. stirp. helv.* n. 391.

Ferum equinum germanicum filiquis in sum-
mitate. *Bauh. pin.* 349. — *Mapp. fl. Als.*
p. 104.

Icon. Rivin. tetr. irr. 97.

German. Teutsche Hufeisen-Hülse.

Call. Fer à cheval vivace.

Mat. Utrinque passim in pratis montosis, col-
libus apricis gramineis asperis siccis, nec non
ad sylvarum marginalia non infrequens. Re-
tro *Durlach* am *Thurnberg* in collibus gra-
mineis calcareo-argillaceis frequens.

Flor. Majo, Junio. *Semina* Julio, Augusto
matura. ☞.

Radix repens, ramosa, fibrosa, dura. *Cau-*
lis palmares, pedales, decumbentes, adscen-
dentes, sulcati, glabri. *Folia* alterna, impari-
pinata. *Foliola* 11-13-15, subpedicellata, op-
posita, subalternaque, lineari-ovalia, obtusa,
ciliolata, integra, inferne villosula, glauco-vi-
ridia. *Stipulae* lanceolatae, acuminatae, integræ.
Flores in pedunculis axillaribus, solitariis, sul-

212 **DIADELPHIA. DECANDRIA.**

ratis, glabris, apice umbelliferi, 5-6-8 f
Pedicelli uniflori, nutantes. *Corollae* luteæ.
gumina compressa, arcuata, utroque la
 sinuata.

Planta puchra saepe colliculos praesertim calca
 graminosos apricos deaurat. Bonum ovibus pab-
 praebet. Flores apibus melliferi.

CCCXLVIII. HEDYSARUM.

Linn. Gen. plant. ed. Schreb. n. 1202

Calyx monophyllus, semiquinquefidus
 ciniis subulatis, erectis. *Corolla* papilionæ
 firiata. *Vexillum* reflexo-compressum, ov
 oblongum, emarginatum, longum. *Alae* ob
 gae, reliquis angustiores, rectae. *Carina* re
 compressa, extrorsum latior, transverse obt
 a basi ad gibbum bifida. *Filamenta* diadelp
Legumen articulis subrotundis, compressis, n
 nospermis, bivalvibus (in *Onobrychi* uno
 ticulo constans echinatum). *Semen* reniform
 solitarium.

1115. HEDYSARUM *Onobrychis* fol
 pinnatis, leguminibus monospermis ac
 leatis, corollarum alis calyce brevior
 bus, caule elongato. *Linn. Sp. pl. 2.*
 1059. — *Poll. fl. pal. n. 694.*

Onobrychis caule erecto ramoso, florib
 spicatis. *Hall. stirp. helv. n. 396.*

Onobrychis folio viciae, fructu echinato n
 jor. *Bauh. pin. 350.* — *Mapp. fl. Alf.*
 212.

Icon. Jacqu. fl. Austr. t. 352.

Grn. Esparsette. Esper. vulgo ewiger Klee. Türkischer Klee.

Call. Esparcette. Saint Foin.

Det. Utrunque passim in collibus praesertim calcareo-argillaceis et pratis montolis non infrequens.

Flr. Junio, Julio. *Semina* Augusto matura. *?*

Radix ramosa, longa. *Caules* ex radice pures, cubitales, tripedales adscendentes, erecti, teretes, striati, ramoli. *Folia* alterna, imparipinnata. *Foliola* 9-13, lineari-oblonga, vel obtusa, integra, subtus glauco viridia, villosiuscula. *Stipulae* lanceolatae, acuminatae. *Flores* in pedunculis praelongis, foliolis nudis, erectis, in spicam cylindricam speciosam longam terminati. *Corollae* purpureae, amoenae. *Alae* brevissimae, oblongae. *Legumina* semi subrotunda, compressa, acute cristato-echinata, monosperma. *Fructus* reniforme, glabrum, cinereum, versus apicem nigricans.

Pianta elegans aprica, colles dumetosos graminum, lapidosos, aridos amans. Passim in pratis artificibus feritur. Bonum praebet pecoribus pabulum.

Flores apibus mel suppeditant.

Semina pullis gallinaceis accepta.

CCCXLIX. ASTRAGALUS.

Linn. Gen. pl. ed. Schreb. n. 1208.

Calyx monophyllus, tubulatus, quinque-
latus acutus. *Corolla* papilionacea. *Vexillum* longius reliquis, lateribus reflexum, emarginatum, obtusum, rectum. *Alae* oblongae, vel breviores. *Carina* longitudine alarum,

214 **DIADELPHIA.** DECANDRIA.

emarginata. Filamenta diadelphea. *Legumen* b
loculare: *loculis* ad alterum latus magis flex
Semina reniformia.

1116. ASTRAGALUS *glycyphyllos* ca
lescens, prostratus, leguminibus sub
quetris arcuatis, foliolis ovalibus,
dunculo longioribus. *Linn. Sp. pl.* 2.
1067. *Pollich. fl. pal.* n. 695.

Astragalus caule prostrato, foliis ovatis, se
pis brevibus racemosis, siliquis sub
quetris incurvis. *Hall. stirp. helv.* n. 41

Astragalus luteus perennis, procumben
vulgaris f. sylvestris. *Morif. hist.* 2. p. 10
— *Mapp. fl. Als.* p. 34.

Glycyrrhiza sylvestris, floribus luteo - pal
centibus. *Bauh. pin.* 352.

Germ. Wildes Süßholz.

Gall. Astragale reglisier.

Hab. Utrinque passim in collibus dumeto
apricis, nec non in sylvaticis gramin
asperis, ad sylvarum marginalia frequer

Flor. Junio, Julio. *Semina* Augusto, Se
tembri matura. ¶

Caules ex radice plures, tripedales et lo
giores, procumbentes teretes, articulati, gl
bri, crassiusculi, ramosi. *Rami* alterni. *Foli*
alterna magna, impari-pinnata. *Foliola* su
sessilia, opposita alternave, 11-13 ovata, o
tusa, integra, glabra, inferne glauco viridi
aristata, reticulato-venosa. *Stipulae* ovatae, ac
minatae, integrae, aut subdenticulatae, magna
Flores axillares in pedunculis solitariis fulca
tis, spica racemosa terminati. *Corollae* pallid

Legumina triangularia, inferne sulcata, pene hamulo subulato terminata, glabra.

Pharm. GLYCYRRHIZA sylvestris Herba; Semen.

Hodie in medicina exoleta.

Infusum foliorum ad Ischuriam et urinæ sabulum a quibusdam commendatur. Planta in Germaniæ pharmacopœia loco Astragali exscapi toto coelo diversifatur.

Folia dulcia pecoribus bonum et acceptum palatum.

Flores melliferi.

III7. *ASTRAGALUS Cicer* caulescens, prostratus, leguminibus subglobosis inflatis, mucronatis pilosis. *Linn. Sp. pl.* 2. p. 1067.

Astragalus caulescens prostratus, pedunculis folio longioribus spicatis, leguminibus didymo inflatis pilosis bilocularibus. Pall. Astrag. p. 57. n. 42.

Astragalus caule ramofo diffuso, spicis erectis, filiquis inflatis rotundis, unguiculatis. Hall. stirp. helv. n. 409.

Astragalus luteus perennis filiqua gemella rotunda velicam referente. Morif. hist. 2. p. 107. — *Mapp. fl. Als. p. 34.*

Cicer sylvestre foliis oblongis hispidis, majus. Bauh. pin. 347.

Icon. Jacqu. fl. Austr. t. 251.

German. Kichernartiger Astragalus.

Pall. Astragale pois-ciche.

Tab. In Alsatia ad agros prope Ehrstein; in pratis in der Wansenau t. Mappo.

Flor. Junio, Julio. Semina Augusto matura. 7

216 DIADELPHIA. DECANDRIA.

Radix repens, ramosa, teres. Caules a radice plures, cubitales, tripedales, decumbentes, subangulati, striati, piloso-villofi. *Folia* alterna, impari-pinnata. *Foliola* subpedicellata, opposita alternave 23-25, ovali-oblonga, obtusa, aristata, integra, piloso-villosa, saturate viridia. *Flores* axillares, in pedunculis solitariis, in racemam subrotundam densam congesti. *Corolla* ochroleuca. *Legumina* rotunda, inflata, compressa, mucronata, hirsuta, polysperma. *Semina* compressa, reniformia, glabra, fusco-vescentia.

Ad rariores omnino Florae cives pertinet,
Bonum pecoribus pabulum subministrat.

1118. *ASTRAGALUS arenarius* caule

cens debilis, pedunculis folio longioribus, leguminibus ovato-acuminatis, non-villosis. *Pall. Astrag. p. 43. n. 46. t. 34*

Astragalus arenarius. Hudf. fl. Angl. I. p. 323. — Relhan. flor. Cantabr. p. 27. — Leyffer. fl. Halens. n. 732.

Astragalus hypoglottis caulescens, procumbens, foliolis oblongis obtusis, spicis ovatis pedunculatis folio longioribus, leguminibus erectis ovatis canaliculatis pilosis. *Willden. Linn. Sp. pl. 3. p. 122. — Linn. Mant. 474. — Lamarck et DeCand. fl. Franc. 4. p. 570.*

Astragalus hypoglottis caulescens prostratus floribus capitatis, leguminibus ovatis dorso canaliculatis pilosis apice aduncis. *Smith. fl. Britan. 2. p. 779.*

Astragalus danicus caulescens, decumbens hirtus: capitulis longe pedunculatis, leguminibus brevibus inflatis, pilosis. *Retz*

DIADELPHIA. DECANDRIA. 317

*obl. bot. 3. p. 41. Hoffm. fl. germ. 263. —
Roth. germ. II. p. 193.*

*Astragalus Onobrychis. Pollich. fl. pal. n.
696. (exclus. synonym.)*

*Astragalus floribus vicie dilute purpureis aut
coeruleis. Mapp. fl. Alf. p. 34. (exclusis
Synonym.)*

Icon. Oeder. fl. Dan. t. 614. (sub A. arenario.)

Berm. Sand-Astragalus.

Sell. Astragale hypoglotte.

*Tab. In Palatinatu inter Mainz et Nierstein
ad vias post Pollichium vidi, nec non in
pratis siccis am Holzhof. In Alsatia prope
Argentiniam auf der Metzgerau bey dem
alten Schiefsberg t. Mappo.*

*Flor. Majo, Junio. Semina Julio, Augusto
matura. 7*

*Radix lignosa, fibrosa. Caulis palmaris,
teretis, decumbens, seu adscendens angulatus hir-
sutus, subramosus. Folia alterna, petiolata, im-
pari-pinnata, foliolis oppositis, sessilibus, su-
perioribus confertioribus, quatuor lineas lon-
gis, duas lineas latis, ovatis, integris, superne
marginatis, utrinque pilosis. Stipulae obtusae.
Flores terminales, spicato-capitati, dense con-
gesti. Calyces cylindrici, hispidi, quinquefidi.
Petallum oblongum, obtusum, margine sub-
reflexum, violaceum, medio album. Alae ve-
llo duplo breviores, obtusae, purpureo-vio-
laceae. Carina acuminata, brevis. Legumina
inflata, inflata, brevia, dorso canaliculata, pi-
stis, apice adunca.*

*Nota. Mappi planta omnino hic, nec ad Astra-
galum austriacum pertinet. Sic etiam Pol-
lichius sub Astragalo Onobrychis, (qui
in alpinis herbis Helvetiae et Austriae crescit,*

218 DIADELPHIA. DECANDRIA.

et vix hucusque in Germania sponte obv
omnino nostram plantam comprehendeba

Obs. In horto botanico ex seminibus culta, non mutavit.

CCCL. TRIFOLIUM

Linn. Gen. pl. ed. Schreb. n. 1311.

Umbellula f. *Capitulum* receptaculo muni. *Calyx* monophyllus, tubulatus, quedentatus, persistens. *Corolla* papilion saepius persistens, marcescens. *Vexillum* flexum. *Alae* vexillo breviores, *Carini* brevior. *Filamenta* diadelpha. *Legumen* v lyce longius, univalve, non dehiscens, daum. *Semina* paucissima, subrotunda.

† *Meliloti*, *Leguminibus* 1-2 *spermi*

1119. TRIFOLIUM *M. officinalis*

minibus racemosis nudis dispermis acutis, caule erecto. *Linn. Sp. p.* — *Pollich. fl. pal. n. 697.*

Melilotus filiquis racemosis nudis disp rugosis acutis. *Hall. stirp. helv. n.*

Melilotus officinarum germaniae. Bauh 331. — Mapp. fl. Alf. p. 190.

Melilotus vera. Tabernæm. p. 895. Ico

β. *Melilotus officinarum germaniae flore Bauh. pin. 331. — Mapp. fl. Alf. p*

Melilotus sylvestris flore albo. Taber p. 893.

Ic on. Oeder. fl. Dan. t. 934. — Bulliard. t. 255. — Blackw. t. 80.

Ger m. Steinklee. Meliloten.

Gall. Melilot officinale.

DIADELPHIA. DECANDRIA. 319

hab. Utrique ad agrorum marginalia, vias, prata, pascua, et arva passim frequens cum β .

flor. Julio, Augusto, Septembri. **Semina** Augusto, Septembri, Octobri matura. ♂

Caules 2-3 pedales et longiores, herbacei, recti aut decumbentes, adscendentes, subangulosi, ramosi, glabri. **Rami** alterni. **Folia** alba, petiolata, ternata, *foliis* ovali-oblongis, obtusis, serratis, glabris, extimo petiolato, ceteris reliquis subsessilibus. **Stipulae** binæ, oppositæ, lineari-subulatæ, parvæ. **Flores** axillares, in pedunculis solitariis, erectis, nudis, villosculis, racemosi. **Racemi** secundi, multiflori. **bracteæ** lineari-subulatæ ad pedicellos, parvæ, solitariae. **Corollæ** luteæ in β . albæ, calyce triplo longiores. **Legumina** pendula, calyce longiora, ovali lanceolata, utrinque acuta, undique rupta, unilocularia, di-rarius monosperma. **Semina** suborbicularia.

Pharm. MELILOTI *Herba; Flores; Semina.*

Planta ficcata peculiari odore valido prædita, amarissima, resolvens, subemolliens, anodyna.

Siccata tineas pellit; pulverisatam plantam cum sabaci pulvere sæpius miscent.

Egregium pecoribus pabulum præbet, in pratis et silvis magis propaganda.

Flores apibus melliferi.

1120. TRIFOLIUM *M. altissimum* leguminibus ramosis nudis, monospermis, caule altissimo fricto.

Trifolium Melilotus officinalis γ . Linn.

Melilotus vulgaris altissima frutescens, flore luteo. *Tournef. inst. Mapp. fl. Alf. p.* 191.

Melilotus vulgaris altissima, frutescens flore luteo. *Dillen. Giesf. app. p.* 7.

220 **DIADELPHIA. DECANDRIA**

β. *Melilotus vulgaris* altissima frutescens flos albo. *Dillen. Gies. p.* 147.

Ger m. *Hoher Steinklee,*

Gall. *Melilot élevé.*

Hab. Utrunque in Rheni regionibus passim copiose. Prope *Dachsland, Knielingen, Eckenstein, Schroek, Linkenheim, Kehl* et alibi frequens. In Alsatia circa Argentoratum a canale prope pontem versus *Eckolsheim* prope *Illkirch* et alibi passim frequens.

Flor. Julio, Augusto. *Semina* Septembri maturat. ☞

Facile distinguitur a praecedente specie *Caulibus* multo semper altioribus, 5-6-9 pedibus, erectis, strictis, durioribus, quasi festulentibus, striato-fulcatis, inferne fusco-rubellis, aut purpurascens, *Foliis* majoribus *Racemis* longioribus, *Floribus* subminoribus *Leguminibus* rotundioribus, monospermis, Odore remissione.

Obs. Planta ex feminibus per plures annos in hortis culta, semper praecedenti multo procerior, imo saepe 10-12 pedalis, stricta, pulcherrima. *Legumina* semper monosperma.

Idem observavi circa variationem **β.** corollis albis instructam.

Caules in plantis cultis non tenaces duri, sed herbacei praecipue in variatione **β.** quam nuper apud oecanum quendam in agro plantatam speciosissimam sub nomine neuer Spanischer Klee vidi. Cultus ejus praecipue in agris pratisque Rheno vicinis, inundationibus saepe expositis, maxime commendanda hucusque plane adhuc ibi neglecta.

Egregium acceptissimumque pabulum pecoribus praebet.

Flores numerosissimi, apibus mel largiuntur.

Lotoidea leguminibus tectis, polyspermis.

21. **TRIFOLIUM** *hybridum* capitulis umbellaribus, leguminibus tetraspermis, caule adscendente. *Linn. Sp. pl. 2. p. 1079.* — *Pollich. fl. pal. n. 698.*

Trifolium caulibus subrectis, spicis depressis, filiquis dispermis. *Hall. stirp. helv. n. 368.*

Trifolium pratense corymbiferum. *Michx. Gen. 28. t. 25. f. 2. 6.*

m. Bastard-Klee.

11. *Trefle batard.*

b. Utrinque in Rheno praesertim vicinia, passim non infrequens in locis aquosis, pratisque, nec non ad aquas stagnantes. ♀

Caules ex radice plures, pedales sesquipedes et longiores, striato-fulcati, glabri, rari. *Folia* alterna, longe petiolata, ternata, ovata breviter pedicellata, ovalia, subserrata, obtusa, in foliis inferioribus saepe emarginata, glabra. *Stipulae* binae, oppositae, ovato-reolatae, acuminatae, aristatae, reticulatae, basi adae, glabrae, integrae. *Flores* ex alis foliorum prioribus in longis pedunculis solitariis striatiglabris, umbellato-capitati. *Capitulum* metrum, inferne floribus reflexis, carneis vel rubris, demum fuscis, superne albidis, albidoincarnatis patenti-erectis instructum. *Legumina* reflexa 2-3-4 sperma.

Pecorum pabulo eximie inservit; Trifolio pratensi inferius, pari ratione coli meretur. Parcius tamen maturatur. *Schreb. Spic. fl. Lipf. p. 25.*

nota. In pratis agrisque cis et transrhenanis, inundationibus saepe expositis, omni studio colenda et propaganda planta.

Flores apibus melliferi.

1122. TRIFOLIUM repens capitulis bellaribus, leguminibus tetraspermis, caule repente. *Linn. Sp. pl.* 2. p. 1080. *Pollich. fl. pal.* n. 699.

Trifolium caule repente, spicis depressis, filiquis tetraspermis. *Hall. stirp. helv.* 367.

Trifolium pratense, flore albo, minus et firmius, glabrum. *Bauh. hist.* 2. p. 280. *Mapp. fl. Als.* p. 311.

Icon. Curtis. fl. Lond. — *Oeder. fl. Dan.* t. 99. *Germ. Kriechender Klee. Weisser Honigklee. Gall. Trefle rampante.*

Hab. Utrunque in cultis, pratis, aggeribus, vicis, ad vias graminosas passim frequenter. *Flor.* Junio, Julio. *Semina* Augusto maturantur.

Caules repentes, radicales, digitales, pedales, teretes, glabriusculi, simpliciusculi ramosive. *Rami* diffusi, alterni. *Folia* alterna, petiolata, terna. *Foliola* subpedicellata, variabilia, nunc obovata, obtusa, subemarginata, nunc obcordata, subrotunda, vel subrhomboidea, firiata, argute ferrulata, glabra, laete viridia, seu fusco-purpurascens, maculata, lunulata albicante sæpe notata. *Stipulae* ovatae, lanceolatae, venosae, margine membranaceae, venis sæpe pallide purpureis pictae. *Flores* axillares, in longissimis pedunculis solitariis, sulcatis, glabris, erectis, capitati. *Capitulum* sphaericum, majusculum, praesertim in cultis plantis. *Corollae* albæ, quandoque albido-incarnatae, junioriores erectae, adultae deflexae, fuscae. *Legumina* oblongo-teretia, torulosa, 2-4 sperma. *Semina* subrotunda, compressiuscula, glabra, fusco-flavicantia.

Prat Capitulis majoribus et minoribus, Foliolis quandoque quatuor ad sex.

Harm. TRIFOLII ALBI Flores.

Hodie in medicina fere exoleti.

Hinc inde magno cum successu in pratis artificiali-
colitur. Egregium pecoribus, et in montosis, ovi-
bulum subministrat.

Flores apibus maxime melliferi.

††† *Lagopoda, calycibus villosis.*

123. TRIFOLIUM rubens spicis villosis
longis, corollis monopetalis, caule ere-
cto, foliis ferrulatis. *Linn. Sp. pl. 2. p.*
1081. — Pollich. fl. pal. n. 700.

Trifolium foliis nervosis ciliatis, spicis ova-
tis, obesis, calycis denticulis plumosis,
imo longissimo. *Hall. stirp. helv. n. 375.*

Trifolium montanum spica longissima ruben-
te. *Bauh. pin. 328. — Mapp. fl. Als. p.*
311.

Lea. Jacqu. fl. Austr. t. 385.

Herm. Großer rother Klee.

Hall. Trefle grande rougeatre.

Hab. In Marggr. superiore inter *Badenweiler*
et *Niederweiler* ad viam montosam argilla-
ceo calcaream am *Schloßberg*; prope
Oberweiler in monte *die Schwärtze*, nec
non prope *Rheinthal* et *Krenzach*, et prope
Loerrach. In Ditione Basileensi prope
Diegten et in monte *Dietisberg*. In Pala-
tinatu circa *Steinbach* in montibus lapidosis
sylvosis quibus in montem *Donnersberg*
adscenditur; Circa *Hartenburg* in collibus
et rupibus gramineis cum *Anthemide tin-*
ctoria post *Pollichium* vidi. In Alsatia prope

1123. PHADELPHIA DECANDENS

Ejchau in fylva, nec non ante dem *Jauer Wald*. t. *Mappo*.

Flor. Junio, Julio. *Semina* Augusto, Septembris matura.

Caulis sesquipedalis, cubitalis, erectus, crassiusculus, firmus, teres, striatus, glaberrimus, simpliciusculus. *Folia* alterna, petiolis vaginatis, ternata. *Foliola* lanceolata, argute serrulata, firma, nervoso-striata, glabra, marginibus villulosa. *Stipulae* praemagnae, vaginatae inferius erectae. *Flores* in spica terminali speciosae, deorsum ovato-cylindrica 2-3 uncias longa, crassa, densa. *Calycis* dentes pilosi, infimus longior, corollae. *Corollae* rubentes. *Legumina* fere tunda, compressiuscula, glabra, calyce corollae cincta, monosperma.

Planta omnino speciosa, ad rariores Flores pertinet.

Flores apibus melliferi.

1124. TRIFOLIUM *alpestre* spicis

densis, corollis subaequalibus, stipulis lanceolatis divergentibus, foliolis lanceolatis, caulibus strictis, simplicissimis. *Act. Soc. Linn. Lond.* 1. p. 234.

Trifolium alpestre spicis subglobosis vix terminalibus, sessilibus, caule erecto, foliis lanceolatis ferrulatis. *Linn. Sp.* 2. p. 1082.

Trifolium montanum purpureum majus. *Bauh. pin.* 328. — *Mapp. fl. Alf. p.*

Trifolium majus secundum. *Clus. hist.* 245.

Icon. Jacqu. fl. Austr. t. 433.

Germ. Aufrechter rother Bergklee.

G.

11. *Trefle rouge étroit de montagne.*

l. Utrique passim in apricis dumetosis præsertim montanis gramineis, ad sylvarum marginalia, nec non ad vias montanas argillaceo-calcareas dumetosas frequens. Retro *Durlach*, *Grözingen*, prope *Weingarten*, *Bruchsal* et alibi abunde.

nr. Junio, Julio. *Semina* Augusto matura. ¶

Caulis palmaris, pedalis et altior, strictus, *lex*. *Foliola* lanceolata, venoso - liriata, *pine* minutissime ferrulata, villosa. *Stipulæ* *matæ*, semiamplexicaules, angustatae, *se-* *mæ*, longæ, villosæ, margine piloso-ciliatæ. *æ* terminalis, sæpissime geminata, ovata, *h*, altera minore. *Calyces* villosissimi, ochro- *â*. *Corollæ* saturate purpureæ.

Folia magis dura quam in *Trifolio* flexuoso et *tensi*, bonum pecoribus alimentum.

Flores apibus melliferi.

25. *TRIFOLIUM medium* spicis laxis, corollis subaequalibus, stipulis subulatis conniventibus, caulibus flexuosis ramosis. *Afzel. Act. Soc. Linn. Lond.* 1. p. 237. — *Linn. fl. Succ. ed.* 2. p. 558.

Trifolium flexuosum spicis subglobosis, villosis terminalibus, caule adscendente flexuoso ramosissimo, foliolis ovato-lanceolatis ferrulatis. *Jacqu. fl. Austr.* 4. p. 45. *Roth. fl. germ.* 2. p. 201.

Trifolium alpestre. *Pollich. fl. pal. n.* 702. — *Leers. fl. Herb.* p. 575.

Trifolium foliis ovatis nervosis, supremis conjugatis, vaginis lanceolatis. *Hall. Juirp. helv. n.* 376.

226. DIADELPHIA. DECANDRIA.

Trifolium majus tertium. *Clus. hist. 2. p. 141.*

Trifolium spica subrotunda rubra, et oblonga rubra. *Mapp. fl. Als. p. 311.*

Icon. Jacqu. fl. Austr. t. 386.

Germ. Gebogener rother Bergklee.

Gall. Trèfle rouge intermédiaire.

Hab. Utrunque in pascuis apricis montosis colliculosis, ad viarum excavatarum dumtosarum marginalia graminosa, nec non pratis montosis. Retro *Durlach*, *Gröden* prope *Berghausen* et alibi passim infrequens; copiose in pratis montosis thermas Badenses.

Flor. Junio, Julio. *Semina* Augusto matura.

Differt a *T. alpestri* cum quo sæpius confundebatur, *Caule* ad genicula flexuoso, cumbente, ramoso, *Foliolis* et *Stipulis* largis, *Spica* sæpissime unica, rarius geminata ovato-globosa, laxiori, *Calycum* dentibus minus villosis, infimo tubum corollæ monopetalæ æquante, *Corollis* dilutius purpureis.

Medium quasi tenet locum inter *T. alpestre* et *pratense*.

Bonum pecoribus pabulum.

Flores apibus mel concedunt.

1126. *TRIFOLIUM pratense* spicis globosis subvillosis, cinctis stipulis oppositis membranaceis, corollis monopetalis *Linn. Sp. pl. 2. p. 1082. — Pollich. fl. pal. n. 701.*

Trifolium spicis densis, corollis inæqualibus dentibus calycinis quatuor æqualibus stipulis aristatis caulibus adscendentibus *Afzel. Act. Soc. Linn. Lond. 1. p. 240.*

Trifolium caule obliquo, foliis ovatis hirsutis, supremis conjugatis, vaginis aristatis. *Hall. stirp. helv. n. 377.*

Trifolium pratense flore monopetalo. *Tournef. inst. Mapp. fl. Als. p. 310.*

Trifolium pratense purpureum. *Bauh. pin. 327.*

2. Trifolium pratense vulgare flore suave rubente. *Mapp. l. c. et ej. dilute purpureum.*

7. Trifolium pratense album. *Bauh. pin. 327. — Mapp. fl. Als. p. 311.*

con. *Oeder. fl. Dan. t. 989. — Blackw. herb. t. 20.*

Germ. Gemeiner Wiesenkle. Futterkle.

Gall. Trefle commun des prés.

Hab. Utrique cum variationibus in pratis, pascuis aggeribus passim abunde.

Flor. Junio, Julio. *Semina* Augusto, Septembri matura. ☞

Caules adscendentes, palmares, pedales, cubitales et altiores, superne striati, subramosi. *Folia* alterna, petiolata, ternata. *Foliola* ovata, obtusa, integra, seu margine subcrenulata, subvillosa, supra macula sæpe centrali pallida aut nigricante sagittata plerumque notata. *Stipulae* latae, breves, vaginantes, aristatae. *Spica* terminalis, solitaria, quandoque gemina globosa, aut ovata obtusa, densa, inter folia duobus opposita, sessilia, quorum *Stipulae* membranaceae, dilatatae, calycem quasi communem constituunt. *Dentes calycini* pilosi quatuor aequales, quintus longior. *Corollae* amoene purpureae, papillae, incarnatae, rosae, et albae, odoratae.

Pharm. TRIFOLII PRATENSIS Flores; Semina.

228 DIADELPHIA. DECANDRIA.

Trifolium hoc apud nos vulgatissimum, frequentissime colitur, copiosissimum, et pergratum pecoribus pabulum præbet.

In agro nonnisi per triennium læte viget plantæ, quo elapso ager aratro pervertendus, et hordeo, avenis, pis, fabis, lino aut cannabi conferendus. Schreb. Samml. I. p. 193.

Si nimium aut recens, solum porrigitur, facile pecorum Emphysema producit.

Flores apibus maxime melliferi, olim cum semibus in medicina recepti.

Obs. In Hispania, cum primis in Catalonia ubi agricultura summum perfectionis et industriae gradum attingit, abunde Trifolium incarnatum colitur, quod amoenissimo colore oculos mire delectat.

Nota. Trifolium angustifolium Linn. sp. villosis conico-oblongis, dentibus calycinis æqualibus setaceis, foliis linearibus, nullis. Germania sponte crescit. Planta a Leerio agnoscata, minime ad T. angustifolium Linn. in Italia et Hispania obveniens, sed ad levissimam variationem Trifolii arvensis Linn. pertinet. Caulis in T. angustifolio Linn. non lævis, Folia non lævia, sed rigidiuscula, margine pilosa, Calycis dentes non purpurei nudi, ut in Leerii planta, sed semper longe fetoso-pilosi.

1127. *TRIFOLIUM ochroleucum* spicis villosis, caule erecto, pubescente, foliolis infimis obcordatis. Linn. Syst. veg. p. 583.

Trifolium caule erecto, foliis hirsutis, foliis premis conjugatis, spicis oblongis. *Habstirp. helv. n. 378.*

Trifolium *ochroleucum* spicis villosis globosis, corollis monopetalis, calycum immo dente longissimo, recto, caule subdeclinato, foliisque villosis, infimis obcordatis. *Huds. fl. Angl. 1. p. 325.*

DIADELPHIA, DECANDRIA. 229

Trifolium pratense hirsutum majus, flore albo-sulphureo seu ochroleuco. *Raj. hist.* 1. p. 943.

Icon. Jacqu. fl. Austr. t. 40.

Germ. Blasgelber Klee.

Gall. Trefle ochreux.

Hab. In pratis uliginosis inter *Dachsland* et *Knielingen* rarius. In pratis montosis humidis sylvaticis *auf dem Kaltenbrunnen*. In *Marggr.* superiore in prato die *Hexmatte* retro *Lippurg* non infrequens. In *Ditione Basileensi* in pascuis montis *Vogelberg* rarius; in pascuis et pratis montium *Muteti, Schauenburg, Hauenstein* frequens.

Fr. Junio, Julio. *Semina* Augusto matura. 7.

Caulis pedalis, sesquipedalis, cubitalis, basi inclinatus, demum erectus, teretiusculus, hirsutus, simpliciusculus, quandoque ramosus. *Folia* alterna, petiolata, ternata. *Foliola* subsessilis, in foliis inferioribus, obcordata, aut ovalis, reliqua oblonga, obtusa; *suprema* lanceolata, angustiora, omnia integra villosa. *Stipulae* amplexicaules, lineari-lanceolatae, subulatae, villosae. *Spica* terminalis, solitaria, erecta, oblongo-ovata, uncialis et major. *Calyces* firriati, pilosi, dente infimo reliquis longiori. *Corollae* monopetalae, albido sulphureae.

Planta elegans, bonum pecoribus pabulum praebet.

Flores apibus melliferi.

Ad rariores Florae cives pertinet, (ni fallor etiam in pratis aquosis circa Argentinam Rhenum versus vidi).

1128. TRIFOLIUM *arvense* spicis villosis ovalibus, dentibus calycinis setaceis aequalibus. *Linn. Sp. pl.* 2. p. 1083. — *Pollich. fl. pal.* n. 703.

230 DIADELPHIA. DECANDRIA.

Trifolium caule ramoso, foliis lanceolatis
ferratis, spicis villosis ovatis. *Hall. stirp.*
helv. n. 373.

Trifolium arvense humile spicatum f. *Lap.*
pus. *Bauh. pin.* 328. — *Mapp. fl. Alf.*
312.

Lagopodium. *Tabernæmont.* p. 908. *Icon*
bona.

Icon. *Oeder. fl. Dan.* t. 724. — *Blackw. her.*
t. 450. (mala).

Germ. *Hasenklee.* *Ackerklee.*

Gall. *Pied de Lievre.*

Hab. Utrunque in agris et arvis passim abunde
præsertim in sterilioribus sabulosis.

Flor. Junio, Julio. *Semina* Augusto matura.

Caulis digitalis, palmaris pedalis et altior
erectus, villosus, sæpe rubellus, maxime ramosus.
Folia alterna, petiolata, ternata. *Fol.*
liola in foliis inferioribus cuneiformia, obtusa
emarginata, superiora sublineari-lanceolata, acutiuscula,
aut sæpius obtusa, quasi truncata, longiora,
integra, omnia villosa, margine ciliato.
Stipulae lineari subulatae, membranaceae, pileosae.
Spicae caulem ramosque terminantes, ovales,
ovali-cylindricae, densae, villosae, incanae vel pallide
rubellae. *Calycis* pilosissimi dentes subaequales
subulati, ultra corollam eminentes. *Corollae* parvae,
albidae. *Alae* intus macula rubella notatae. *Legumina*
subrotunde ovata, glabra, monosperma.

Bonum quibus in arvis pabulum praebet.

Nota. *Trifolium* gemellum *Wilden.* *Lin.*
Sp. pl. 3. p. 1376. ex *Cl. Pourretto* receptum,
in Hispania lectum. est variatio *T. arven-*
sis in sterilioribus vulgarissima, caulibus demissioribus
simpliciusculis, Spicis superne duabus gemellis instructa.

DIADELPHIA. DECANDRIA. 231

¶ Semina ex talibus emarcidis individuis in Catalonia legi. in horto culta, exhibuerunt individua majora, quammaxime ramosa, spicis copiosis instructa, nostratibus simillima.

Trifolium phleoides. Willden. l. c. a Cl. Pourretto in Hispania allegatum, est variatio levidensis Trifolii angustifolii Linnaei. promiscue in sterilibus duris Cataloniae et Aragoniae obveniens.

4129. *TRIFOLIUM scabrum* capitulis sessilibus lateralibus ovatis, calycibus inaequalibus rigidis, persistentibus. Linn. *Sp. pl.* 2. p. 1084.

Trifolium caulibus prostratis, capitulis ovatis, in alis sessilibus, calycibus rigidis, segmentis erectis. Hall. *stirp. helv.* n. 371.

Trifolium cujus caules è geniculis glomerulos oblongos producant. Bauh. *hist.* 2. p. 378. — Mapp. *fl. Als.* p. 312.

Trifolium flosculis albis in glomerulis asperis cauliculis adnatis. Vaillant. *bot. par.* 196.

Leon. Vaillant. l. c. t. 33. f. 1.

Germ. Rauher Klee.

Gall. Trefle apre.

Hab. In pascuis siccis apricis passim. In Marggr. superiore in pascuis fabulosis ad *Wiesam* inter *Steinen* et *Loerrach*. In ditione Basileensi copiose passim in pascuis ad *Birsek* et *Birsam*. In Palatinatus pascuis tenui gramine vestitis ad vias arborum circa *Lautern* t. *Pollichio*.

Flor. Junio, Julio. Semina Augusto maturat. ☉

Caules e radice plures, procumbentes, digitales, palmares et longiores, teretes, villosi.

fiusculi, subramosi. *Folia* alterna, petiolata, ternata. *Foliola* subpedicellata, obovata, obtusa, integra, pubescentia. *Stipulae* subrotundae, saepe aristato-villosae, venis rubellis pictae. *Capitula* ovato oblonga, axillaria, terminaliaque sessilia, solitaria. *Calyces* glabri, dentibus recurvis, pilis adpressis instructi, infimo longioribus. *Corollae* albae, parvae, angustae.

1130. *TRIFOLIUM striatum* capitulis sessilibus sublateralibus, ovatis, calycibus striatis rotundatis. *Linn. Sp. pl.* 2. p. 1085.

Trifolium capitulis sessilibus lateralibus ovatis, calycibus subrotundis villosis striatis. *Linn. fl. Succ.* 614.

Trifolium parvum hirsutum, flore parvo luteo purpureo, in glomerulis mollioribus oblongis, semine magno. *Vaill. bot. parisi.* p. 196.

Icon. Vaillant. l. c. t. 33. f. 2.

Germ. Gestreifter Klee.

Gall. Trefle strié.

Hab. In Marggraviatu superiore in pascuis aridis apricis tenui gramine tectis circa *Loersrach* ad Wiesam, nec non in ditione *Besileensi* ultra *Schorenbruch* ad Wiesam. & *Lachenalio*.

Flor. Junio, Julio. *Semina* Augusto matura. ©

Differt a praecedente, *Foliolis* cuneiformi-obovatis, superiora versus obsolete crenulatis, pubescenti-sericeis, *Stipulis* longioribus, ovato-acuminatis, setaceo-aristatis, villosis, *Capitulis* ovatis, acutiusculis, fructificatis longioribus, *Calycibus* totis pilosis, decemstriatis, *Corollis* carneis.

III *Vesicaria calycibus inflatis ventricosis.*

431. *TRIFOLIUM fragiferum* capitulis subrotundis, calycibus inflatis bidentatis, reflexis, caulibus repentibus. *Linn. Sp. pl. 2. p. 1086. — Pollich. fl. pal. n. 705.*

Trifolium caule repente, spicis glabris, calycibus sericeis, ampullescentibus. *Hall. stirp. helv. n. 370.*

Trifolium fragiferum, *Frisicum*, folio cordato, flore rubro. *Morif. hist. 2. p. 144. — Mapp. fl. Als. p. 312.*

Trifolium fragiferum, *friscum*. *Bauh. pin. 329.*

Leon. Curtis. fl. Lond. — Oeder. fl. Dan. t. 1042. Vaillant. bot. par. t. 22. f. 2.

erm. Erdbeeren - Klee.

Gall. Trefle Fraisor.

Lab. Utrunque in pascuis demissioribus graminosis humidis, aquosis, rhenanis præsertim, passim frequens. Circa *Dachsland*, *Knielingen*, *Schröck*, *Linkenheim*, *Ruffheim*, et alibi.

Flor. Junio, Julio. *Semina* Augusto, Septembri matura. ¶

Caules palmares, pedales, cubitales et longiores, repentes, teretiusculi, læves, purpurecentes, ramosi. *Folia* alterna, petiolata, ternata. *Foliola* ovali-subrotunda, aut obovata, obtusa, emarginata, superiora acutiuscula, subvillosa, quandoque glabra, argute ferrulata, venulosa. *Stipulae* ovato-acuminatæ, venosoreticulatæ. *Pedunculi* floriferi longi, solitarii, recti, teretes, glabriusculi. *Capitula* florifera subrotunda, parva, *Trifolii* repentis emula, sed

minora, purpurascencia; *fructifera* multo i
jora, *Fragariæ* fructum æmulantia, rotund
carnea. *Involucrum* sub singulo capitulo p
phyllum, *foliolis* setaceis. *Calyx* tubulosus, l
olus, supra gibbosus, quinquedentatus, d
tibus tribus inferioribus æqualibus, acumin
viridibus, duobus superioribus paulo longi
bus subulatis, rigidulis, apice rufis. *Floris* fru
ficati calycis pars gibba, inflatur, reticulata fit,
pericarpium obtegit. *Corollae* pallide purpuri
Legumina ovata, compressa, disperma, aut n
nolperma. *Semen* ovato - reniforme, nitidū
flavicans.

Bonum pecoribus pabulum subministrat.

Nota. Trifolium capitulo spumoso aspero ma
Bauh. Prodr. p. 140. — Mapp. fl. All
312. Hall. stirp. helv. n. 366.

Est planta dubia a solo C. Bauhino in salt
Michelsfeldae, Huningam versus in dumetis se sufful
repta. Ea dumeta dudum destructa sunt et fortis
castello Huningenfi cesserunt. Caules bicubitales
nues, flexiles, folia rotunda, crenata, vulgaris
folii similia, in praelongis pedicellis. Scapi palmar
pedales. In singulis scapis capitulum subrotundum,
vae avellanae mole: flosculis aureis, his succo
folliculi membranacei, sibi incumbentes, filamentis
busdam exasperati. Ita C. Bauhinus.

De hoc Trifolio ne conjecturam habeo, qu
offeram. Hall. l. c.

++++ Lupulina, *vexillis corollae inflexis*

1132. **TRIFOLIUM montanum** spi
subimbricatis subtribus, vexillis subul
emarcescentibus, calycibus nudis, cau
erecto. *Linn. Sp. pl. 2. p. 1087.* — *P
lich. fl. pal. n. 706.*

Trifolium caule erecto paucifloro, foliis
liptico-lanceolatis, nervosis, subtus si
hirsutis. *Hall. stirp. helv. n. 372.*

DIADELPHIA. DECANDRIA. 235

* *Trifolium montanum album.* *Bauh. pin.* 328.
Mapp fl. Alf. p. 311.

Trifolium pratense album, *Tabernamont. p.*
 907. *Icon bona.*

con. Oeder. fl. Dan. t. 1172.

erm. Weisser aufrechter Berghles.

all. Trefle blanc de montagne.

ab. Utrique in pratis montosis siccis, apri-
cis, juniperetis, passim non infrequens,
nec prata in plantis posita sicca ad Rhe-
num respuit.

lor. Junio, Julio. Semina Augusto, Septem-
bri matura. ¶

Caulis pedalis, sesquipedalis et altior, ere-
ctus, firmus, teres, hirsutus, canescens, sim-
plicifolius, apicem versus subramosus. Folia
ovata, petiolata, superiora subsessilia, ternata.
loba lanceolata, acuta, aristata terminata, mi-
ne argute serrulata, transverse venoso-striata,
villosa, subtus pallidiora. Stipulae lanceolatae,
minutae, villosae. Flores in spicis 2-3, pedun-
culatis, terminalibus, densis, ovatis, obtusis. Ca-
lis dentes subaequales, subpubescentes. Corol-
lae semper albae. Vexillum dimidia parte inflex-
um. Legumina ovato-subrotunda, planiuscula,
glabra, monosperma.

Flores apibus melliferi.

133. *TRIFOLIUM spadiceum* spicis
 ovalibus imbricatis, vexillis deflexis per-
 sistentibus, calycibus pilosis, caule erecto.
Linn. Sp. pl. 2. p. 1087.

Trifolium caule erecto, spicis strepentibus,
calycibus capillaribus pilosis. Hall. stirp.
helv. n. 365.

236 **DIADELPHIA. DECANDRIA.**

Trifolium montanum lupulinum. *Bauh pi*
328. *Prodr.* 140. — *Mapp. fl. Als.* p. 31

Lotus montanus aureus, amplo **Lupuli** e
pitulo, annuus. *Barrel. lc.*

Icon. Barrel. lc. 1024.

Germ. Gelbbrauner Hopfenklee.

Gall. Trefle houblonet.

Hab. Passim in pratis montosis apricis. *Pro*
Pforzheim et *Huchensfeld.* In *Marggr.* 1
periore prope *Lippurg auf der Hexma*
et alibi. In *Brisgovia* prope *Tryberg*
graminosis montosis versus *dem Wasserfa*
In *Ditione Basileensi* prope *Bottming*
In *Alsatia* retro *Hochandlau* in pratis me
tosis t. *Mappo.*

Flor. Junio, Julio. *Semina* Augusto matura.

Caulis palmaris, pedalis, sesquipedalis
erectus, simpliciusculus, quandoque ramosus
Rami 1-2-3, alterni, erecti, striati, glabri
Folia alterna, petiolata, ternata. *Petiol* u
ciam ad sesquiunciam longi. *Foliola* brevissim
pedicellata, infima ovalia, emarginata, obtu
quandoque obcordata; superiora ovalia, obtu
omnia argute minutim serrulata, a costa tra
versim striato-venulosa, glabra, 6-7 lineas lo
ga, 4 lineas lata. *Stipulae* lineari-lanceolatae
acutae, glabrae, novem lineas longae, erectae
Spicae caulem ramosque terminantes, fae
binae, pedunculatae, erectae, ovali-oblongae
6-8 lineas longae, 5-6 latae, floribus dense im
bricatae, luteo-spadiceae. *Calyces* ante flor
centiam, pilosi: dentibus binis superioribus c
foletis. *Corollae* desfloratae, deflexae, ferru
neae, persistentes.

Ad rariores Florae cives pertinet. Non conf
denda cum sequente specie, primo intuitu affini.

34. *TRIFOLIUM campestre* spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, stipulis petiolo longioribus, lanceolatis, foliolis subsessilibus, obovati-lanceolatis, obtusis emarginatis, caule erecto.

Trifolium aureum spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, stipulis lanceolatis nudis, foliolo impari sessili, caule erecto. *Pollich. fl. pal. n.* 708.

Trifolium agrarium. *Roth. fl. germ.* 2. p. 207. (excluso *Linnaei* synonymo.)

Trifolium pratense luteum semina, flore pulchriore live lupulino. *Bauh. hist.* 2. p. 381. (descr. bona, *Icon pessima.*

Icon. Oeder. fl. Dan. t. 558. bona. (sub *T. agrario*).

Germ. Großer gelber Hopfenklee.

Gall. Trefle houblonet grande.

Hab. In Marggr. superiore in dumetosis graminosis prope *Steinen* ad *Wiesam*. In *Brisgovia* prope *Tryberg* promiscue cum *T. spadiceo* nuper legi. In Palatinatu circa *Lautern* in pratis siccis, pascuis tenui gramine vestitis frequens, nunquam in agris reperitur. Cum *Malva moschata*, *Campanulis*, *Centaurea nigra* passim crescit; in montibus sylvestribus circa *Steinbach* teste *Polluchio*. In *Alsatia* in margine agrorum montis *Rosberg* prope *Maszmünster* t. *Joh. Bauhino*.

Flor. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ☉.

Differt a praecedenti specie quoad habitum, staturam, Folia, Stipularum formam et mag-

DIADELPHIA. DECANDREA.

nitudinem, maxime affini, Petiolis foliorum 1-2 uncialibus, sed tantum 4-5 lineas longis. *Stipulis* petiolo longioribus, lanceolatis, aminatis, glabris novem lineas longis, erectis. *Capitulis* seu *Spicis* ovalibus, obtusis, 7 lineis longis, semiunciam latis, caulem ramosque terminantibus, speciosis aureis, nec luteo-spiceis; defloratis non ferrugineis. *Calycibus* glabris.

Nota. Sub *T. agrario* Linnaei et aliorum, sim botanicorum duae species distinctae hucque militabant; Haec scilicet quam sub *T. campestri* adduxi, et quae nunquam in agris, semper in locis siccis graminosis, dumetosis, et pестribus crescit; altera cum primis segetibus agrisque dicata, sub *T. agrario* designata.

Obs. Trifolium campestre nostrum in Horto Botanicum, mutationi non succubuit.

Nota. In Sturmii Deutschlands Flora mihi non ad manus est, Perillustris praefes de Schreber Trifolia lupulina disposuit.

1135. TRIFOLIUM agrarium spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus, subpubescentibus stipulis ovato-acuminatis, ciliatis, foliis cuneiformi-obovatis subsessilibus, impari longius pedicellato, infracto, caule subdiffuso.

Trifolium agrarium. Pollich. fl. pal. n. 76 (omnino).

Trifolium agrarium spicis ovalibus, imbricatis vexillis deflexis, persistentibus, calycibus nudis, caule erecto. *Linn. pl. 2. p. 1087.*

Trifolium spicis ovatis densissimis strepens, caulibus diffusis. *Hall. stirp. helv. 363.*

Trifolium pratense luteum capitulo Lupuli
 f. agrarium. *Bauh. pin.* 328. — *Mapp.*
fl. Alf. p. 311. — *Vaillant. bot. par. t.* 22.
 f. 3. (omnino) bona.

Curtis. fl. Lond. t. 161. (optima sub *T.*
agrario).

Acker-Hopfenklee. Großer Acker-
Hopfenklee.

Trefle de champs.

Utrinque passim in agris cultis, inter se-
 getes ac stipulas, nec non in graminosis
 colliculosis siccioribus frequens.

Junio, Julio, Augusto. Semina Augusto,
 Septembri matura. ☉

Caulis palmaris, pedalis et longior, fac-
 iens ad terram diffusus, quandoque plus mi-
 nus erectus, teres, villosus, ramosus. *Rami*
simpliciter, aut erectiusculi, alterni, cauli analogi.
Folia alterna, petiolata, ternata. *Petioli* 5-7
 lineas longi. *Foliola* cuneiformi-obovata,
 a medio ad finem argute ferrulata, gla-
 briata, *impari* longius pedicellato, *Pedi-*
cellula 2-3 lineari. *Stipulae* ovatae, acuminatae,
 4 ad 5 lineas longae. *Capitula* axilla-
 ria, in pedunculis solitariis, erectis, semiun-
 cialibus, subrotundo-ovalia, ob-
 imbricata, densiuscula, 40 ad 55 flora,
 aurea, antiquata fusca, semiuncialia, spe-
 cia. *Calyces* quinquedentati, dentibus duo-
 bus superioribus brevissimis, tribus inferioribus
 prioribus, setaceis, albidis, pilosulis. *Legu-*
mina ovali-subrotunda, planiuscula glabra, par-
 monosperma.

In Horto botanico culta, omnino annua, faciem
 non mutavit.

Bonum ovibus pabulum.

1136. *TRIFOLIUM pseudo-procumbens* spicis ovali-oblongis, imbricatis vexillis deflexis persistentibus, calycibus villosiusculis, stipulis ovatis latissimis liatis, foliolis omnibus pedicellatis et ovatis.

Icon. Oeder. *fl. Dan.* t. 796. (*sub procumbente*).

Germ. Mittlerer Hopfenklee.

Gall. Trefle houblonnet intermediaire.

Hab. Utrunque in agris arvisque nec non vineis et locis graminosis aridis passim.

Flor. Junio, Julio, Augusto. *Semina* Augusto Septembri matura. ☉

Medium omnino tenet locum inter *T. agrarium* et *T. procumbens*. *Linn.* Distinguitur a *T. agrario*, *Foliorum* foliolis minoribus, brevioribus, omnibus semper manifeste pedicellatis, impari quidem etiam reliquis longius pedicellato, sed pedicello semper breviori, *Stipulis* duplo fere latioribus, brevioribus, ovatis, ovatusculis, non longe acuminatis, *Capitulis* foliorum sub minoribus, ovali-oblongis, magis dense imbricatis, croceo-luteis, nec sulphureis. In singulo capitulo *Flores* 40-54 numeravi.

Differt a *T. procumbente* *Linn.* sub quo hucusque militabat, *Foliorum* foliolis non cuneiformi-obcordatis, semper majoribus, *Stipulis* duplo latioribus, minus acutis, *Capitulis* nec ovali-subrotundis obtusis, sed ovali-oblongis duplo fere longioribus, densioribus, *Floribus* in singulo capitulo semper copiosioribus, densius imbricatis.

Nota. Oederi *Icon* citata, optime plantam nostram repraesentat.

In horto culta, faciem non mutavit.

Semper magis loca culta, et praesertim agros et
amat, dum *T. procumbens* in graminosis,
primis in pratis ubique fere abundat.

Bonum praesertim ovibus pabulum.

57. *TRIFOLIUM procumbens* spicis
ovali-subrotundis imbricatis, vexillis de-
flexis persistentibus, foliolis cuneiformi-
obcordatis pedicellatis, stipulis ovatis
acutis, caulibus subprocumbentibus.

Trifolium procumbens spicis ovalibus imbricatis, vexillis deflexis persistentibus caulibus procumbentibus. *Linn. Sp. pl.* 2. p. 1088. — *Pollich. fl. pal. n.* 709.

Trifolium lupulinum alterum minus. *Raj. angl.* 3. p. 330. t. 14. f. 3.

Trifolium filiforme spicis subimbricatis, vexillis deflexis persistentibus, calycibus pedicellatis, caulibus procumbentibus. *Linn. Sp. pl.* 2. p. 1088.

Trifolium lupulinum minimum. *Raj. angl.* 3. p. 331. t. 14. f. 4.

a. *Curtis fl. Lond. t.* 307.

m. *Kleiner Hopfenklee.* β. *Kleinsten Hopfenklee.*

b. Utrunque in pascuis, pratis macilentis aliisque, locisque graminolis passim abunde, nec agros respuit.

r. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Planta pro diverso solo natali, erectiuscula, endens, et procumbens, digitalis, palmapedalis, et altior, simpliciuscula, saepissime in ramosa, glabriuscula, aut villosula. *Volia*

1136. TRIFOLIUM *pse* es, 3-4 lin
bens spicis ovali ania cuneifo
vexillis deflexis cellato, striata,
villosiusculis, late viridia. Stipi
liatis, folioli atæ, glabriusculæ, i
ovatis. petiolo subbreiores.
Icon. Oeder. f. us axillaribus, erectis,
bente). culis, uncialibus, sesquiu
Germ. Mitt. subrotunda, obtusa, multifi
Gall. Tr. lutea. Flores in singulo capi
Hab. V. pedicellati, imbricati, 28-36, rariu
vi. speciminibus exsuccis id est in variat
Flo. 15. Corollae Vexillum deflexum,
fulcatum, luteum, demum fusi
ovata, compressa, monosperma.

Semina ex speciminibus paucifloris exsuccis
est ex T. filiformi lecta, in horto botanico
ta, exhibuerunt individua, omnino capitulis
tisoris prædita, variationi a. aut T. proc
benti simillima. Hinc minime pro peculiari f
declarandum.

Egregium ovibus pabulum præbet.

CCCL. LOTUS.

Linn. Gen. pl. ed Schreb. n. 1212.

Umbella simplex. Calyx monophyllus,
bulosus, femiquinquefidus: dentibus ac
æqualibus, erectis. Corolla papilionacea.
xillum subrotundum, deorsum flexum: U
oblongo, concavo. Alae subrotundæ, ve
breviores, latæ, sursum conniventes. Ca
inferne gibba, superne clausa, acuminata,
scendens, brevis. Stamina diadelpha. Legu

DELPHIA

L. filiquoso tetrangulum), stric-
ce longius, multiloculare,
cylindrica.

quosus leguminibus fo-
branaceo-quadrangulis cau-
cumbentibus, foliis subtus pu-
bentibus. *Linn.Sp. pl. 2. p. 1089.* —
Willch. fl. pal. n. 710.

Lotus caule prostrato, scapis unifloris, fili-
quis alatis. *Hall. stirp. helv. n. 386.*

Tetragonolobus filiquosus. *Roth. fl. germ.*
2. p. 226. Scopol. fl. Carn. n. 938.

Lotus pratensis filiquosus luteus major et
durior. Bauh. pin. 332. — Mapp. fl. Alf.
p. 181.

Trifolium filiquosum. Tabernæmont. p. 904.
Icon.

Ja. Jacqu. fl. Austr. t. 361.

erm. Einblumiger Wiesen-Hülfsenkle.

all. Lotier filiqueux.

ab. In pratis humidis cis et transrhenanis,
pascuisque passim non infrequens. Prope
Dachsland, Knielingen, Ekkenstein et alibi.
In Principatu superiore in pascuis grami-
nosus ad Lacum Bodamicum passim abunde.

er. Majo, Junio. Semina Julio, Augusto
matura. 7

Caulis palmaris, pedalis, decumbens, te-
tangulatus, pubescens, simpliciusculus. Fo-
liolterna, petiolata, ternata. *Foliola* subfes-
sa, obcordato-ovata, obtusa, arisia termina-
ta, integerrima, superne glabra, inferne pubes-
centi-sericea, albida. *Stipulae* binæ, cordatæ.

242 DIADELPHIA. DECANDRIA.

alterna, petiolata. *Petiol*i tenues, 3-4 longi. *Foliola* pedicellata, omnia cuneis exacte obcordata, superiora versus ferrimpari reliquis longius pedicellato, striatibra, quandoque villosa, læte viridia. *St* rhomboideo-ovatæ, striatæ, glabriusculæ, gine ciliatæ, acutæ, petiolo subbreuiore: *pitula*, in pedunculis axillaribus, erecti litariis, villosiusculis, uncialibus, sesquialibus, ovali-subrotunda, obtusa, multi 3-4 linearia, lutea. *Flores* in singulo capbreuissime pedicellati, imbricati, 28-36, rari 25, in speciminibus exsuccis id est in variis β. 5-13-15: *Corollae Vexillum* deflexum rinatum, sulcatum, luteum, demum fuf *Legumina* ovata, compressa, monosperma flexa.

Obf. Semina ex speciminibus paucifloris exsucc est ex T. filiformi lecta, in horto botanica, exhibuerunt individua, omnino capitulitifloris prædita, variationi α. aut T. probenti simillima. Hinc minime pro peculiari declarandum.

Egregium ovibus pabulum præbet.

CCCLI. LOTUS.

Linn. Gen. pl. ed Schreb. n. 1212.

Umbella simplex. Calyx monophyllus bulosus, semiquinquefidus: dentibus æqualibus, erectis. *Corolla* papilionacea. *xillum* subrotundum, deorsum flexum: *L* oblongo, concavo. *Alae* subrotundæ, vobreviores, latæ, sursum conniventes. *C* inferne gibba, superne clausa, acuminata scendens, brevis. *Stamina* diadelpa. *Leg*

Madricum (in *L. filiquoso* tetrangulum), stric-
tum, fæctum, calyce longius, multiloculare,
valve. *Semina* plura, cylindrica.

1138. *Lotus filiquosus* leguminibus fo-
litariis, membranaceo-quadrangulis cau-
libus procumbentibus, foliis subtus pu-
bescentibus. *Linn. Sp. pl.* 2. p. 1089. —
Pollich. fl. pal. n. 710.

Lotus caule prostrato, scapis unifloris, fili-
quis alatis. *Hall. stirp. helv.* n. 386.

Tetragonolobus filiquosus. *Roth. fl. germ.*
2. p. 226. *Scopol. fl. Carn.* n. 938.

Lotus pratensis filiquosus luteus major et
durior. *Bauh. pin.* 332. — *Mapp. fl. Alf.*
p. 181.

Trifolium filiquosum. *Tabernæmont.* p. 904.
Icon.

Icon. Jacqu. fl. Austr. t. 361.

germ. Einblumiger Wiesen-Hülfsenkle.

Gill. Lotier filiqueux.

Det. In pratis humidis cis et transrhenanis,
pascuisque passim non infrequens. Prope
Dachsland, Knielingen, Ekkenstein et alibi.
In Principatu superiore in pascuis grami-
nolis ad Lacum Bodamicum passim abunde.

Flor. Majo, Junio. *Semina* Julio, Augusto
matura. 7

Caulis palmaris, pedalis, decumbens, te-
trangulatus, pubescens, simpliciusculus. Fo-
lium alterna, petiolata, ternata. *Foliola* subsef-
ta, obcordato-ovata, obtusa, arista termina-
ta, integerrima, superne glabra, inferne pubes-
centi-sericea, albida. *Stipulae* binæ, cordatæ.

444 **DIADELPHIA. DECANDRIA.**

Flos terminalis, solitarius, foliolis ternis lanceolatis suffultus. *Calyx* sericeus. *Corolla* speciosa, pallide sulphurea. *Legumen* tetragulum angulis membranaceis, polyspermum.

Bonum pecoribus pabulum.

Flores apibus melliferi.

139. *Lotus corniculatus* capitulis compressis, caulibus decumbentibus, leguminibus cylindricis, patentibus. *Lin. Sp. pl.* 2. p. 1092. — *Pollich. fl. pal.* 711.

Lotus floribus umbellatis, filiquis cylindricis, rectissimis. *Hall. stirp. helv.* n. 36.

Lotus arvensis caule angulato, pedunculis non tubulosis, calycum dentibus cinctis, leguminibus cylindricis, patentibus. *Schkuhr. bot. Handb.* n. 4139. tab. 2.

Lotus sive melilotus pentaphyllos minus glabra. *Bauh. pin.* 332. *Mapp. fl. Alp.* p. 180.

Trifolium filiquosum minus. *Tabernæm.* f. 905. *Icon.*

β. *Lotus corniculata* hirsuta minor. *Bauh. hist.* 2. p. 356. *Mapp. l. c.* 181.

γ. *Lotus tenuifolius* pentaphyllos frutescens tenuissimis glabris foliis. *Bauh. pin.* 332. *Pollich. l. c. n.* 711. ♀

Icon. *Curtis fl. Lond.* — *Oeder. fl. Dan.* t. 991
Germ. Gehörnter Schotenklee.

Gall. Lotier corniculée.

Hab. Utrinque in pratis, agris, pascuis, vineis ad viarum marginalia passim frequens. γ in pratis ad Salinas circa *Dürkheim t. Pollichio.*

DIADELPHIA. DECANDRIA 245

Junio, Julio. *Semina* Augusto, Septembri matura. ☞

Caules saepe ex radice plures, palmares, acaules, decumbentes aut adscendentes, angulati, ramosi. *Folia* alterna, petiolata, terrena. *Foliola* ovali-lanceolata, acuta, aristata, serrata, glabra, aut plus minusve hirsuta. *Stipulae* duae foliolis similes, sed iis latiores. *Flores* subumbellati 5-12 petiolis longis nudis insertis. *Calycum* dentes quinque setacei, erecti. *Corollae* flavae. *Vexillum* superne aurantium. *Legumina* cylindrica, isthmis quasi interrupta, glabra, polysperma. *Semina* subreniformia, fusca, maculata.

Variat pro diverso solo sicciore et humidior. *Caules* procumbentibus et erectiusculis, *Foliis* glabris et hirsutis, *Foliolis* *Stipulis*que latioribus robustioribus.

Bonum armentis, et in montosis pascuis siccioribus et pabulum praebet.

Flores apibus mel concedunt,

10. *LOTUS uliginosus* caulibus pedunculisque teretibus tubulosis, calycibus apice recurvis laciniatis, leguminibus cylindricis, patentibus. *Schkuhr. bot. Hand. n. 4138. tab. 211.*

Lotus corniculatus β. *Linn Sp. pl. 2. p. 1092.*

Pollich. l. c. var. α. n. 711. major magis erecta,

Lotus pentaphyllos, flore majore, luteo, splendente. *Bauh. pin. 181. — Mapp. fl. Alf. p. 181.*

Loti corniculatae major species. *Bauh. hist. 2. p. 355.*

n. Rivin. tetr. t. 76.

246 DIABELPHIA. DECANDEIA.

Germ. Gehörnter Sumpf • Schotenklees.

Gall. Lotier corniculée des marais.

Hab. Utrunque in umbrosis sylvaticis humidis ad vias, in pratis udis, ad fossas aquaticas passim frequens.

Flor. Junio Julio, Augusto, *Semina* Augusto Septembri matura. 7

Diffinguitur a praecedenti specie, *Calycibus* longioribus saepe bipedalibus et altioribus erectis, teretibus, *Pedunculisque* cavis, *Foliorum* *Stipulis* et *Floribus* majoribus. *Calycinis* lobis ante anthesin praecipue recurvis, *Leguminibus* fere duplo longioribus, *Seminibus* minoribus.

Hanc speciem omni jure a Loto corniculato acutissimus Schkuhr separavit.

Planta in horto culta, mutationi non succubunda.

Bonum pecoribus pabulum.

Flores apibus melliferi.

CCCLII. M E D I C A G O.

Linn. Gen. plant. ed. Schreb. n. 1214.

Calyx monophyllus, rectus, campanula cylindraceus, semiquinquefidus, acuminatus aequalis. *Corolla* papilionacea. *Vexillum* obtusum, integrum, marginibus inflexis, totum inflexum. *Alae* ovato-oblongae, appendice carinae affixae, lateribus conniventibus sub caetera. *Carina* oblonga, bifida, patens, obtusius deflexa a pistillo, hiansque a vexillo. *Filamenta* diadelpha. *Legumen* compressum, longum, inflexum. *Semina* plura reniformia.

DIADELPHIA. DECANDRIA. 247

811. Cochleatae Siliqua spiraliter cochleata.
Falcatae Siliqua arcuata, falcata.

8141. **MEDICAGO** *fativa* pedunculis racemosis, leguminibus contortis, caule erecto glabro. *Linn. Sp. pl.* 2. p. 1096. — *Pollich. fl. pal.* n. 712.

Medica caule erecto, foliis oblongis serratis, racemis erectis, filiquis planis iterato contortis. *Hall. stirp. helv.* n. 382.

Medica legitima. *Clus. hist.* 1. p. 242.

Foenum burgundicum. *Lobel. Icon.* 2. p. 36.

Corollis luteo-violaceis.

Icon. *Regnault bot.*

Germ. *Lucerne. Lucernerklée.*

Gall. *Sainfoin. Luferne cultivée.*

Hab. Utrunque in agris, pratis cultis, vineis passim. In arvis et pascuis cis et transrhenanis spontanea. Colitur passim copiose. β . in arvis ad Rhenum prope *Dachsland* et *Knielingen* non infrequens.

Flor. Julio, Augusto. Semina Augusto Septembri matura. γ

Caules ex una radice plures, sesquipedales, cubitales et altiores, erectiusculi, quandoque subprocumbentes, angulato-fulcati, glabriusculi aut villosuli, ramosi. *Folia* alterna, breviter petiolata, ternata. *Foliola* lanceolata, obtusa, aristata, a medio ad apicem serrulata, striata, glabriuscula: impari pedicellato. *Stipulae* binae, lanceolato acuminatae, setaceae, basi subdenticulatae. *Flores* in pedunculis angulatis, axillaribus, solitariis racemosi. *Corollae* coeruleo-violaceae, *Alae* dilutiores. *Legumina* glabra, plana, iterato contorta.

248 **DIADELPHIA. DECANDRIA.**

Variatio β . memorabilis ob corollas subminores, semper, et id jam primo exortu, violaceo-luteas. Haec ex feminibus in horto culta, faciem non mutavit. forsan non immerito pro peculiari specie declaranda.

Obf. In Catalonia legi specimina longiora, tenuiora, semper ad terram decumbentia subvillosa-incana. Foliolis angustioribus Corollis amoenissime amethystinis praedita. Semina ex his lecta, in Horto Botanico culta, exhibuerunt individua Catalonica simillima pulcherrima, decumbentia.

Bonum pecoribus alimentum, hinc praecipue in sola calcareo-argillaceo passim abunde colitur.

Flores apibus mel largiuntur.

1142. MEDICAGO *falcata* pedunculis racemosis, leguminibus lunatis, caule prostrato. *Linn. Sp. pl.* 2. p. 1096. — *Polidich. fl. pal.* n. 713.

Medica foliis oblongis serratis, filiquis racemosis. *Hall. stirp. helv.* n. 381.

Medica sylvestris floribus croceis. *Bauh. hist.* 2. p. 383. — *Mapp. fl. Alf.* p. 189.

Falcata. Rivin. tetr. 6. 8.

nitatem erecti, obsolete angulati, villosiusculi
ut glabri, ramosi. *Rami* alterni, tetragoni,
longati. *Folia* alterna, petiolata, ternata. *Fo-*
liola lineari-lanceolata, obtusa, subemarginata,
ristata, integriuscula, verticaliter striata gla-
briuscula: impari pedicellato. *Stipulae* duae,
parvae, lanceolato setaceae. *Flores* in peduncu-
lis axillaribus, solitariis, folio longioribus, ra-
cemosi. *Corollae* saturate luteae. *Legumina* fal-
cata, villosiuscula, nigra, 7-9 sperma. *Semina*
subreniformia, fusca.

Bonum pecoribus pabulum.

Flores apibus melliferi.

1143. *MEDICAGO lupulina* spicis ovali-
bus, leguminibus reniformibus mono-
spermis, caule procumbente. *Linn. Sp.*
pl. 2. p. 1097. — *Pollich. fl. pal.* n. 714.

Medica caule diffuso, capitulis hemisphæri-
cis, siliquis reniformibus. *Hall. stirp.*
helv. n. 380.

Melilotus capsulis reni similibus in capitu-
lum congestis. *Mapp. fl. Als.* p. 191.

Trifolium pratense luteum capitulo brevior.

Bauh. pin. 328.

Trifolium arvense. *Tabernæm. p.* 908. *Icon.*

con. Curtis. fl. Lond. t. 120. — *Oeder. fl. Dan.*
t. 992. — *Schkuhr. bot. Handb. t.* 212. a.

erm. Gelber Hopfen-Schneckenklee.

Gall. Luferne-houblonet.

Tab. Utrique in cultis, hortis, agris, arvis,
pratis, pascuis, vineis et viis passim fre-
quens.

lor. Junio, Julio, Augusto. *Semina* Augusto,
Septembri matura. ☉ ☿

Caules palmares, pedales et longiores, procumbentes, subangulosi, villosiusculi, ramuli alterni diffusi. *Folia* alterna breviter ovata; ternata. *Foliola* obovata, aut obcordata, obsolete denticulata, aristata, villosa mollia late viridia; impari pedicellato. *Stipulae* duae, ovato-lanceolatae, acuminatae, sinu denticulatae. *Flores* in pedunculis axillaribus solitariis, folio longioribus, capitato-spicae parvae initio subrotundae, dein ovali basi ad unum latus nudiusculae, densae, multiflorae, luteae. *Calycini* dentes quinque inaequales, tres inferiores reliquis longiores. *Corolla* parvae luteae. *Legumina* reniformia, compressa rugosa, spiraliter cochleata, nigra; monosperma. *Semen* ovatum, flavicans, glabrum.

Bonum armentis et ovibus acceptum et veniens pabulum, hinc passim in anglia colitur.

Nota. Planta in campestri solo ovibus dicato, propaganda.

1144. *MEDICAGO minima* pedunculata multifloris, leguminibus cochleatis pilulis aculeis subulatis, rectis uncinatis, stipulis integerrimis, foliolis obovatis dentatis. *Linn. Sp. pl. ed. Willden. 3. p. 1418.*

Medicago polymorpha minima. Linn. Pl. pal. n. 715.

Medicago minima pedunculis multifloris, leguminibus cochleatis: spinulis hamatis stipulis integris. *Schreb. Spic. fl. Lips. p. 1.*

Medica echinata minima. Bauh. hist. 2. p. 3. — Mapp. fl. Alf. p. 190.

Cochleata echinata minima. Rivin. t. 88

Icon. Oeder. fl. Dan. t. 24. — Schkuhr. l. t. 212. a.

herm. Kleinster Schneckenklee.

Call. Luferne petite.

Lab. In Margraviatus superioris collibus aridis prope *Efringen*, nec non circa *Basileam* in locis sterilibus ad *Birsam* et prope *St. Jacob*. In Palatinatu inter *Hartenberg* et *Dürkheim* in monte cui vetustum Monasterium *Limburgense* inædificatum est, locis siccis lapideis. In collibus juniperinis siccis circa *Callstadt*, *Herzheim* post *Pollichium* vidi. In Alsatia prope *Argentnam* in campestribus aridis asperis in *der Ruprechtsau*.

Flor. Majo, Junio. Semina Augustio matura. ☉

Caules ex radice plures, digitales, palmatis et longiores, procumbentes, quandoque ascendentes, subflexuosi, teretiusculi, villosi, tosi. *Folia* alterna, petiolata, ternata. *Folia* cuneiformi-obovata, obtusa, quandoque marginata, superne denticulata, subincana, pilosa; impari pedicellato. *Stipulae* duæ, ovato-lanceolatae, acutæ, integræ. *Flores* in pedunculis axillaribus, solitariis, folio subbrevis, spicati. *Spiculae* subrotundæ, 3-5-6 floreae, parvæ, luteæ. *Legumina* pedicellata, sæpissime tria, parva, cochleata, pilosa; aculeis tribus, rectis, apice uncinatis, imbuta, 6-7-merma. *Semina* reniformia, glabra, flavicantia.

Plantula pulchella ad rariores Floræ cives pertinet, piose in Galliæ, Italiæ et Hispaniæ aridis crescit.

CLASSIS XVIII.
POLYADELPHIA
POLYANDRIA.

CCCLIII. HYPERICUM.

Linn. Gen. pl. ed. Schreb. n. 1224.

Calyx quinquepartitus: *laciniis* subovati concavis persistentibus. *Corolla* petala quinque oblongo-ovata, obtusa, patentia. *Stamina* *Filamenta* numerosa capillaria, libera, basi tantum in 5 seu 3 phalanges coalita. *Antherae* parvae. *Styli* tres (interdum in exoticis 1, 2, 5) longitudine staminum. *Capsula* subrotunda; loculis numero styli. *Semina* plurima, oblonga.

1145. *HYPERICUM quadrangulare* floribus trigynis, caule quadrangulo herbaceo, foliis pellucido-punctatis, foliolis calycinis lanceolatis. *Smith. fl. brit.* 2. p. 801.

Hypericum quadrangulare floribus trigynis, caule quadrato herbaceo. *Linn. Sp. pl.* 2. p. 1104.

Hypericum caule quadrangulari, foliis ovatis, perforatis, punctatis. *Hall. stirp. helv.* n. 1038.

Hypericon *Ascyron* dictum, caule quadrangulo. *Bauh. hist.* 3. p. 382. — *Mapp. fl. Alf.* p. 149.

POLYADELPHIA. POLYANDRIA. 253

Icon. Curtis. fl. Lond. t. 231. — Oeder. fl. Dan. t. 640.

erm. Viereckiges Johanniskraut.

all. Willepertuis carré.

ab. Utrunque in pratis, fylvaticisque humidis, ad fossas, paludes, sepes passim frequens, nec sicciora respuit.

lor. Junio, Julio. Semina Augusto matura. 7

Caulis sesquipedalis. cubitalis, erectus, quadratus, angulis membranaceis, fusco-rufus, glaber, ramosus. *Rami* oppositi, densati, cauli analogi. *Folia* opposita, sessilia, amplexicaulia, ovata, obtusa, lævia, septa-nervia, superficie punctis diaphanis minutis adspersa, margine glandulis nigris subtus desertim punctata. *Flores* caulem ramosque terminantes, dense paniculati. *Paniculae* ramuli quinei. *Pedunculi* brevissimi. *Calyx* quinque-lobatus: *Laciniis* lanceolatis, acutis, patentibus, integerrimis, nudis. *Corollae petala* quinque, longitudine calycis, lineata, lutca, glandulis nigris margine punctulata. *Antherae* flavæ, glandula nigra notatæ. *Styli* tres, patentes. *Capitula* trilocularis, trivalvis, fusca, polysperma.

Tota planta pulchro rubro colore tingit.

Flores apibus melliferi.

1146. HYPERICUM *Leersii* floribus trigynis, caule subquadrato herbaceo, foliis ovatis amplexicaulibus, calycisque laciniis ovatis obtusis.

Hypericum dubium floribus trigynis, caule obsolete quadrangulo, foliis obtusis im-

254 POLYADELPHIA. POLYANDRIA.

punctatis, foliolis calycinis ellipticis. *Smit fl. brit. 2. p. 802.*

Hypericum dubium floribus trigynis caule subquadrato herbaceo, foliis ovatis, calycibus obtusissimis. *Leers. fl. Herb. n. 59*

Hypericum fallax floribus trigynis, foliis ovato-oblongis rugosis, calycisque laciniis ovatis obtusis, caule subquadrangulari. *Grimm. fl. Jfen. in Nov. Act. A. 1 C. vol. 3. app. 362.*

Hypericum delphinense caule recto simpliciter tetragono, foliis reticulatis punctatis, calicium foliolis ovatis punctatis, corollae medio minoribus, floribus proliferis. *Willd. lars. fl. Delph. 3. p. 497.*

Hypericum quadrangulare. *Pollich. fl. n. 716. (exclusis synonymis).*

Germ. Leersisches Johanniskraut.

Gall. Millepertuis de Mr. Leers.

Hab. Cum præcedente passim, præsertim in pratorum fossas. In Marggr. superiore in fossas pratorum circa Thermas prope *Badenweiler* frequens.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. *¶*

Distinguitur a præcedente specie, *Caulibus* altioribus, minus manifeste quadratis, *Foliis* simplicibus, majusculis, in superficie punctatis diaphanis destitutis, marginescariosis, *Floribus* multo majoribus, *Calycinis* laciniis Corollae brevioribus ovalibus, obtusis, lineolis interruptis notatis, *Capsulis* ventricosis, triangularibus, calycibus reflexis, ad basin cinctis.

Obs. Planta in horto ex seminibus culta, constanter hinc minime dubia aut fallax species.

47. *HYPERICUM perforatum* floribus trigynis, caule ancipiti, foliis obtusis, pellucido-punctatis. *Linn. Sp. pl. p. 1105.* — *Pollich. fl. pal. n. 117.*

Hypericum caule terete alato ramosissimo, foliis ovatis perforatis. *Hall. stirp. helv. n. 1037.*

Hypericum vulgare. *Bauh. pin. 279. Mapp. fl. Alf. p. 149.*

Hypericon L. *Tabern. p. 1249 Icon.*

con. Curtis. fl. Lond. — Oeder. fl. Dan. t. 1043. — Blackw. t. 15.

erm. Gemeines Johanniskraut.

all. Mille-pertuis. Mille-pertuis commun.

ab. Utrinque ad vias, agrorum margines et sepes passim abunde.

flor. Junio, Julio. Semina Augusto maturat.

Caules ex radice plures sesquipedales, cules erecti, duri, ancipites, ramosi. *Rami* oppositi, ancipites, erecti. *Folia* opposita, sessilia, ovato-oblonga, obtusa, perforata. *Flores* caulem ramosque terminantes, paniculati. *Calycinae* laciniae lanceolatae, acutae. *Corollae* petala flava; ad unum latus crenulato-ferrulata, glandulis nigris adspersa. *Antherae* glandula nigra notatae. *Styli* tres. *Capsula* subtrigona, triangularis, fusca, polysperma. *Semina* oblonga; nigra, fusco-nigricantia.

erm. HYPERICI Herba; Flores; Semen.

Americans, balsamica, vulneraria, anthelminthica.

Floribus spiritum frumenti pingunt laete purpureum confici. *Lin. fl. Suec. p. 265. FUGA DAEMONUM* nominabatur, quoniam spectra et daemones fugare, superstitiosi credunt. *Principale* ingrediens famosi olim

Balsami Magici Pforzhemiensium, (Beaume de Pforzheim) cui vires insignes ad incantamenta adscribebantur.

1148. HYPERICUM *humifusum* floribus trigynis axillaribus solitariis, caulis ancipitibus prostratis filiformibus, foliis glabris. *Linn. Sp. pl. 2. p. 1103.* — *Pollich. fl. pal. n. 1039.*

Hypericum caule prostrato, foliis ovatis, calycibus ferratis, punctatis. *Hall. stirp. helv. n. 1039.*

Hypericum minus supinum, vel supinum glabrum. *Bauh. pin. 279.* — *Mapp. fl. Als. p. 149.*

Icon. Curtis. fl. Lond. t. 172. — *Oeder. fl. Dan. t. 141.*

Germ. Kleines hingestrecktes Johanniskraut.

Gall. Mille pertuis couché.

Hab. Utrunque in arenosis sterilibus pascuis et sylvis sabulosis cricetosis humidiusculis passim copiose. Circa *Carlsruhe* et in adjacenti regione frequens.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ¶

Caules ex eadem radice sæpe plures, digitales, palmares, pedales, procumbentes, rari erectiusculi, teretes, subancipites, glabri, rubelli, superne ramosi. *Folia* opposita, sessilia, ovalia, ovali-lanceolata, obtusa, glabra, integra, minute perforata, margine punctis glanduliferis fuscis aut purpureis notata. *Flores* ex axillis ramorum, terminalesque, pedunculatæ. *Calycinæ* laciniae inæquales, ovali-oblongæ, margine reflexæ, ferrato-glanduloso-punctatæ. *Corollæ* petala oblonga, obtusa, calyce paulo

culo longiora, flava, margine glandulifera. *Semina Filamenta* subviginti, in tres phalanges divisa. *Styli* tres. *Capsula* oblonga, trilobularis, immatura amoene rubra, polysperma. *Semina* minima.

Plantula pulchella in sabulosis humidiusculis are-
na figit.

149. *HYPERICUM montanum* flori-
bus trigynis, calycibus ferraturis glandulosis, caule tereti erecto glabro, foliis ovatis. *Linn. Sp. pl.* 2. p. 1105. — *Pollich. fl. pal.* n. 719.

Hypericum foliis ovatis per oram punctatis calycibus lanceolatis ferratis globuliferis. *Hall. stirp. helv.* n. 1042.

Hypericum elegantissimum non ramosum, folio lato. *Bauh. hist.* 3. p. 382. — *Mapp. fl. Alf.* p. 149.

na. Oeder. fl. Dan. t. 173.

na. Berg-Johanniskraut.

ill. Mille pertuis des montagnes.

ab. Utrunque passim in montosis dumetosis, et pratis editioribus apricis. Retro Durlach, Wohlfartsweyher, Ettlingen et alibi.

ar. Julio, Augusto. Semina Septembri matura. ☞

Caulis cubitalis et altior, erectus, teres, ser, simplex. *Folia* opposita sessilia, semiplexicaulia, ovata, glabra, margine punctis glanduliferis nigricantibus obsessa, integra, subvenoso-reticulata. *Flores* terminales. *Bracteae* ferrato-ciliatae, glanduliferae. *Calycinae* lanceolatae, glabrae, margine ciliato-glandulosae. *Corollae Petala* ovali-lanceolata, ob-

458 POLYADELPHIA. POLYANDRIA.

tusa, integra, pallide flava. *Styli* tres. *Stigma* purpurea.

1150. *HYPERICUM hirsutum* floribus
trigynis, calycibus ferrato-glandulosis
caule tereti erecto, foliis ovatis subpubescentibus. *Linn. Sp. pl.* 2. p. 1105.
Pollich. fl. pal. n. 720.

Hypericum foliis ovatis hirsutis, calycibus lanceolatis ferratis glanduliferis. *Hort. stirp. helv.*

Hypericum villosum erectum, caule rotundo. *Tournef. inst.* — *Mapp. fl. Als.* p. 15

Androsæmum hirsutum. *Bauh. pin.* 280.

Icon. Curtis. fl. Lond. t. 162.

Germ. Zottiges Johanniskraut.

Gall. Millepertuis velue.

Hab. et Flor. cum præcedente. 7

Caulis sesquipedalis, cubitalis, tripedalis erectus, solidus, teres, pubescens, simplicifolius. *Folia* opposita subsessilia, inferiora breviter petiolata ovali-oblonga, obtusa, septemnervia, utrinque villosula, integra, margine ciliata. *Flores* terminales in panicula erecta racemosa, ovata. *Calycinae* laciniae lanceolatae bifidae, margine glandulosae. *Corollae* petalis ovali-oblonga, obtusa, striata, patentia, flavis. *Styli* tres. *Capsula* oblonga, trilocularis, trivalvis, polysperma.

1151. *HYPERICUM pulchrum* floribus
trigynis, calycibus ferrato-glandulosis
caule tereti, foliis amplexicaulibus cordatis, glabris. *Linn. Syst. veg.* p. 702. —
Pollich. fl. pal. n. 721.

POLYADELPHIA. POLYANDRIA. 259

Hypericum foliis amplexicaulibus cordatis, calycibus ovatis serratis glanduliferis.
Hall. stirp. helv. n. 1041.

Hypericum minus erectum. *Bauh. pin. 279.*

Hypericum pulchrum. *Trag. 74.*

Icon. Curtis. fl. Lond. — Oeder. fl. Dan. t. 75.

Germ. Kleines aufrechtes Johanniskraut.

Call. Mille pertuis elegant.

Tab. Utrunque in sylvaticis siccis apricis passim. Circa Carlsruhe in sylva Hauaker et alibi non infrequens.

Flor. Junio, Julio. Semina Augusto matura. 7.

Caulis pedalis, sesquipedalis et altior, erectus, teres, glaber, simpliciusculus, quandoque ramosus. Folia opposita, sessilia, amplexicaulia, cordata, obtusa, integra, firmula, minutissime punctulis pertusa, inferiora saepius ovata, pulchella, glaberrima. Flores terminales in pediculis oppositis, erectis pedicellatis subtrifloris. Bractea sub pedicellis parva, ovali-oblonga, integra. Calycinae laciniae ovatae, acutae, serratae, margine ferrulatae, glandulis fusco rufis obsitae. Corollae petala ovali-oblonga, flava, intus aurantiaco-lineata, margine serrata, glandulifera. Styli tres. Capsula subconica, trilobata, fusca, polysperma.

Planta pulchella in hortis facile cicuratur.

Nota. In Hortis Carlsruhanis sub dio adhuc hospitantur:

HYPERICUM kalmianum floribus pentagynis, caule fruticoso, foliis lineari-lanceolatis. *Linn. ex Virginia.*

HYPERICUM monogynum floribus monogynis-staminibus corolla longioribus, calycibus coloratis, caule fruticoso. *Linn. ex China, Japonia.*

260 POLYADELPHIA. POLYANDRIA:

- * **HYPERICUM calycinum** floribus pentagynis scitariis terminalibus, caule tetragono fruticoso foliis oblongo-ovatis obtusis coriaceis. Linn. ex Graecia.
- * **HYPERICUM Ascyron** floribus pentagynis, caule tetragono herbaceo erecto simplici, foliis laevibus integerrimis. Linn. ex Sibiria, Pyrenæis.
- * **HYPERICUM Androsæum** floribus trigynis, fructibus baccatis caule fruticoso, ancipiti. Linn. ex Anglia, Italia, Gallia, Hispania.
- * **HYPERICUM elatum** floribus trigynis, calycibus lanceolato-ovatis, acutis, staminibus corolla longioribus, caule fruticoso, foliis ovato-oblongis. Aiton. hort. Kiew. 3. p. 104. ex America boreali.
- * **HYPERICUM hircinum** floribus trigynis, staminibus corolla longioribus, caule fruticoso ancipiti. Linn. ex Sicilia, Calabria, Creta.
- * **HYPERICUM prolificum** floribus trigynis, primordialibus sessilibus, caule ancipiti, fruticoso, foliis lanceolato-linearibus. Linn. ex America septentrionali.
- * **HYPERICUM canadense** floribus trigynis, foliis lineari lanceolatis, caule quadrangulo, pericarpis coloratis. Linn. ex Canada.

CLASSIS XIX. Y N G E N E S I A.

POLYGAMIA AEQUALIS.

*emiflosculosi. Tournef. Corollis ligulatis
omnibus,*

LXVIII. CICHORIUM. *Recept. subpaleaceum, Pappus sub 5-dentatus.
Cal. calyculatus,*


LXV. HYPOCHAERIS. *Recept. paleaceum. Pappus subplumosus. Cal.
imbricatus.*

LIV. TRAGOPOGON. *Recept. nudum.
Pappus plumosus, stipitatus.
Cal. simplex, polyphyllus.*

LVI. PICRIS. *Recept. nudum. Pappus
plumosus, stipitatus. Calyx
duplicatus.*

LXII. LEONTODON. *Recept. nudum. Pappus
pilosus, stipitatus. Cal.
calyculatus, imbricatus, squamis
laxiusculis.*

LVII. APARGIA. *Recept. nudum, aut
subvillosum. Pappus sessilis,
in centralibus substipitatus,
plumosus. Cal. imbricatus.*

- CCCLV. SCORZONERA.** *Recept. nudum. Pappus plumosus, stipitatus. Cal. imbricatus, margine scarioso.*
- CCCLXIV. CREPIS.** *Recept. nudum. Pappus pilosus. Cal. calyculatus, squamis difformibus.*
- CCCLX. CHONDRILLA.** *Recept. nudum. Pappus pilosus stipitatus. Cal. calyculatus, multiflorus.*
- CCCLXI. PRENANTHES.** *Recept. nudum. Pappus pilosus. Cal. calyculatus, subquinqueflorus.*
- CCCLIX. LACTUCA.** *Recept. nudum. Pappus pilosus, stipitatus. Cal. imbricatus, margine scarioso.*
- CCCLXIII. HIERACIUM.** *Recept. nudum. Pappus pilosus, sessilis. Cal. imbricatus, ovatus.*
- 

SYNGENESIA. 263

** Capitati.

LXXIV. CARLINA. *Calyx* radiatus: radiis coloratis!

LXXI. CARDUUS. *Recept.* villosum. *Pappus* capillaris. *Cor.* flosculosa. *Cal.* imbricatus, ventricosus.

LXXII. CNICUS. *Recept.* villosum. *Pappus* plumosus. *Cor.* flosculosa. *Cal.* imbricatus, ventricosus.

LXX. SERRATULA. *Recept.* paleaceum. *Pappus* plumosus. *Cor.* flosculosa. *Cal.* imbricatus oblongo-cylindricus.

LXXIII. ONOPORDON. *Recept.* favosum! *Pappus* capillaris. *Cor.* flosculosa. *Cal.* squamis ventricosus spinosis.

LXIX. ARCTIUM. *Recept.* paleaceum. *Pappus* setoso paleaceus. *Cor.* flosculosa. *Cal.* subglobosus, squamis apice incurvato-hamosis!

† *Centaurea nigra.*

*** *Discoidei.*

LXXVI. CACALIA. *Recept.* nudum. *Pappus* pilosus. *Cal.* calyculatus.

LXXVIII. CHRYSOCOMA. *Recept.* nudum. *Pappus* pilosus. *Cal.* imbricatus.

SYNGENESIA,

CCCLXXVII. *ERRATORIUM. Recept. nudum.*
Pappus plumosus. Cal. imbricatus. Pistilla longissima.

CCCLXXV. *BIDENS. Recept. paleaceum. Pappus aristatus. Cal. imbricatus.*

POLYGAMIA SUPERFLUA.

* *Discoidei.*

CCCLXXX. *ARTEMISIA Recept. subnudum.*
Pappus nullus. Cor. nulli.

CCCLXXXII. *CARPESIUM. Recept. nudum.*
Papp. nullus. Cor. radii 5-fidi.

CCCLXXIX. *TANACETUM. Recept. nudum.*
Pappus submarginatus. Cor. radii trifidæ.

CCCLXXXIII. *CONYZA. Recept. nudum. Pappus pilosus. Cor. radii trifidi.*

CCCLXXXI. *GNAPHALIUM. Recept. nudum.*
Pappus plumosus. Cal. scariosus, squamis concavis.

** *Radiati.*

CCCXCIII. *BELLIS. Recept. nudum. Pappus nullus. Cal. squamis æqualibus simplex.*

CCCXCV. *MATRICARIA. Recept. nudum. Pappus nullus. Cal. squamis imbricatis, acutis.*

SYNGENESIA. 265

- CIV. CHRYSANTHEMUM.** *Recept. nudum. Pappus nullus. Cal. squamis intimis scariosis.*
- CII. DONONICUM.** *Recept. nudum. Pappus pilosus. Papp. radii nullus.*
- CI. ANEMONE.** *Recept. subnudum. Pappus pilosus. Stamina radii castrata.*
- C. INULA.** *Recept. nudum. Pappus pilosus. Antherae basi bifidae,*
- XXXIV. ERIGERON.** *Recept. nudum. Pappus pilosus. Cor. radii capillares,*
- XXXVIII. SOLIDAGO.** *Recept. nudum. Pappus pilosus. Cor. radii subfeni, remoti.*
- XXXIX. CINERARIA.** *Recept. nudum. Pappus pilosus. Cal. æqualis, simplex.*
- XXXVI. SENEIO.** *Recept. nudum. Pappus pilosus. Cal. apice squamis sphacelatis.*
- XXXV. TUSSILAGO.** *Recept. nudum. Pappus pilosus. Cal. squamis submembranaceis.*
- XXXVII. ASTER.** *Recept. nudum. Pappus pilosus. Cal. subquarrosus.*

CCCXCVI. ANTHEMISIA. *Recept. paleaceus*
Pappus nullus. Cal. hemisphaericus.

CCCXCVII. ACHILLEA. *Recept. paleaceus*
Pappus nullus. Radius
quinqueflorus. Cal. oblongus.

CCCXCIX. BUPHTHALMUM. *Recept. paleaceum.*
Pappus marginatus. Stamina hermaphroditis simplicia.

CCCXCVIII. GALINSOGA. *Recept. paleaceus*
Pappus polyphyllus, paleaceus. Cal. imbricatus.

POLYGAMIA. FRUSTRANEA.

CCCCI. CENTAUREA. *Recept. setosum. Pappus*
pilosus. Radius corollae tubulosus.

CCCC. HELIANTHUS. *Recept. paleaceus*
Pappus aristatus. Cal. squarrosus.

POLYGAMIA. NECESSARIA.

CCCCII. CALENDULA. *Recept. nudum. Pappus*
nullus. Semina membranacea.

POLYGAMIA. SEGREGATA.

CCCCIII. ECHINOPS. *Perianth. 1-florus*
Pappus pubescens. Polyaequalis.

S Y N G E N E S I A. 267

*** MONOGAMIA.**

CIV. JASIONE. *Cal. communis. Cor. quinquepetala, irregularis. Caps. infera, 2-locularis.*

XV. VIOLA. *Cal. 5-phyllus. Cor. 5-petala, irregularis. Caps. superæ, 3-valvis.*

XVI. IMPATIENS. *Cal. 2-phyllus. Cor. 4-5petala, irregularis, calcarata. Caps. superæ, 5-valvis.*

CLASSIS XIX.
SYNGENESI
POLYGAMIA. ÆQUALIS.

CCCLIV. TRAGOPOGON.

Linn. Gen. plant. ed. Schreb. n. 1229

Calyx communis simplex suboctophyllus, *foliis* lanceolatis, æqualibus, omnibus coalitis. *Corolla* composita, imbricata, unimembra. *Corollulis* hermaphroditis multis, exte-
rioribus paulo longioribus. *Propria* modice
ligulata, truncata, quinquedentata. *Calyx*
etiferus connivens, longitudine seminum, tri-
triculofus. *Semina* oblonga, utrinque attenu-
angulata, scabra, stipite subulato longo per-
gastro terminata. *Pappus* plumosus. *Recepta-
culum* nudum, planum, scabrum.

1152. *TRAGOPOGON pratense* cal-
bus corollæ radium æquantibus, foliis
tegris strictis. *Linn. Sp. pl. 2. p. 1109.*
Pollich. fl. pal. n. 722.

Tragopogon foliis gramineis, carinatis,
plexicaulibus caudatis subcrispis, cal-
lore æquali. *Hall. stirp. helv. n. 8.*

Tragopogon pratense luteum majus. *Be-
pin. 274. — Mapp. fl. Alf. p. 307.*

Tragopogon luteum. *Tabernæmont. p. 1.
Icon.*

*Icon. Oeder. fl. Dan. t. 906. — Bulliard. H.
L. 209.*

INGENESIA. POLYG. ÆQUALIS. 269

m. Gelber Wiesenboksbart. Hafermark.

ll. Barbe de Bouc ordinaire. Salsifix des près.

b. Utrinque passim in pratis præsertim montosis frequens. Copiose in pratis, pomariis Brisgovie ad montem Kaiserstuhl, nec non in Margraviatu superiori prope Schopshcim abunde.

r. Majo, Junio. Semina Julio, Augusto matura. ♂

Radix fusiformis, simplex, extus fusca, inlbida. Caulis sesquipedalis, cubitalis, erectus, glaberrimus, subramosus. Folia alba, amplexicaulia, lanceolata, superne anthera, integra, glabra. Pedunculi terminantur nudi uniflori, superne incrassati. Calyx phyllus, basi coalitus, corollæ æqualis aut revior, glaber. Corolla universalis speciosa,

m. TRAGOPOGI f. BARBAE HIRCI Radix.

Radix sapida, dulcis. Caules juniores dulci lactescentes, passim in cibum recipiuntur.

Flores apibus melliferi, utilagine sæpe corrupti

bonum pecoribus in pratis alimentum præbet.

Totam plantam pueruli legunt, comedunt, et sumaxime delectantur.

3. TRAGOPOGON majus calycibus subtredecimphyllis corollæ radio longioribus, foliis radicalibus linearibus angustis, pedunculis superne incrassatis.

tragopogon majus calycibus corollæ radio longioribus, foliis integris strictis, pe

SYNGENESIA, POLYG. ÆQUIV.

dunculis superne incrassatis, corollæ ad apicem rotundatis. *Linn. Syst. ed. Murr.* 14. p. 710.

Icon. Jacqu. fl. Austr. 1. t. 29.

Germ. Großer gelber Bocksbart.

Gall. Barbe de Bous majeure.

Hab. In collibus graminosis basalticis apud Brigoviam am *Kaiserstuhl* prope arcem *Lauburg* cum *Panico dactylo* rarius, nec in vineis adjacentibus nuper vidi cum *nerofo Ittnero*.

Flor. Junio, Julio. *Semina* Augusto maturantur.

Differt a præcedenti specie *Caule* alio. *Foliis* radicalibus caulinisque infimis exactis linearibus, angustis, *Pedunculis* superne incrassatis, *Calycibus* non octophyllis sed 13 phyllis: *Foliolis* angustioribus, corollæ rariore semper multo longioribus, *Corolla* univerni pallide sulphurea.

Obs. Individua loco citato nascentia, et alia in Hispania et Gallia meridionali a me lecta simillima sunt. Planta ex seminibus in horto culta, mutationi non succubuit.

Ad rariores omnino Floræ nostræ cives pertinet.

1154. *TRAGOPOGON porrifolium* calycibus corollæ radio sesquilingioribus foliis integris strictis, pedunculis superne incrassatis. *Linn. Sp. pl.* 2. p. 1110.

Tragopogon foliis gramineis amplexicaulis, calyce florem superante. *Hall. stir. helv.* n. 9.

Tragopogon purpureo-cæruleum porrifolium quod *Artesi* vulgo. *Bauh. pin.* p. 274.

LYNGENESIA. POLYG. AQUALIS. 272

non. Oeder. fl. Dan. t. 797. — Schkuhr. bot. Handb. t. 214.

erm. Purpurrother Bocksbart.

all. Salfiz commun.

a b. In Sponhemia prope Kreuznach versus salinas non infrequens.

lor. Majo, Junio. Semina Julio Augusto matura. ☉

Caulis 3-4 pedalis, erectus teres, glaber. Folia alterna, amplexicaulia, linearia, erecta, integra, glabra. Pedunculi ad florem incrassati. Calyx 12-13 phyllus, basi coalitus, corolla duplo major. Corolla purpureo-coerulea. Corollulae angustae, dentatae. Semina angulata, scaberrima.

Individua allegata cum his, quae in Hispania lecta consensu simillima sunt.

Hinc inde, et id praesertim in Gallia colitur et in horti loco in culinis adhibetur.

CCCLV. S C O R Z O N E R A.

Linn. Gen. pl. ed. Schreb. n. 1230.

Calyx communis imbricatus, longus, subcylindraceus: squamis subquindecim, margine auriculatis. Corolla compolita, imbricata, unilobata. Corollulis hermaphroditis plurimis, anterioribus paulo longioribus. Propria monolobata, ligulata, linearis, truncata, quinquefidata. Calyx fructiferus ovato-oblongus, immixtus, tandem patenti-reflexus. Semina longa, cylindracea, striata, calyce dimidio breviora. Pappus plumosus. Receptaculum nudum.

272 SYNGENESIA. POLYG. ÆQUAL

1155. *SCORZONERA humilis* caule nudo subunifloro, calycinis squamatis, foliis oblongo-lanceolatis nervosis planis. *Linn. Sp. pl. ed. Willd. p. 1599.*

Scorzonera humilis caule subnudo unifloro, foliis latolanceolatis nervosis, p. *Linn. Sp. pl. 2. p. 1112. Pollich. fl. n. 723.*

Scorzonera lanata. Schrank. Bavar. 2. p.

Scorzonera caule nudo, unifloro, foliis lanceolatis. *Hall. stirp. helv. n. 10.*

Scorzonera humilis latifolia pannonica. hist. 2. p. 138.

β. *Scorzonera humilis* foliis angustioribus vosis.

Scorzonera austriaca caule subnudo unifloro, calycinis squamis glabris, foliis linearibus lanceolatis. *Willden. Linn. Sp. pl. 1498.*

Scorzonera caule nudo unifloro, foliis angustis. *Hall. stirp. helv. n. 11.*

Scorzonera humilis angustifolia, ponno Clus. hist. 2. p. 139.

Icon. Jacqu. fl. Austr. 1. t. 36.

Germ. Niedere Scorzonere.

Gall. Scorzonere nain.

Hab. In pratis montosis apricis siccioribus prope *Pforzheim, Dillstein, Huchenfeld* prope *Weissenstein*. In Palatinatu transitorio in pratis siccioribus inter *Franken-Lambsheim* et *dem Holzhorf*. In Principatu *Fürstenbergensi* in pratis et collibus
mi

SYNGENESIA. POLYG. ÆQUALIS. 279

minofis ficcis prope *Donaueschingen* et alibi cum variatione β . frequens.

Flor. Majo, Junio, Julio. *Semina* Julio Augusto matura. γ

Radix fusiformi-cylindrica, extus nigricans, intus alba. *Caulis* palmaris, pedalis et altior, erectus, simpliciusculus, teres, glaber, sæpe pilosus, superne villosus - incanus. *Folia* radicalia numerosa, petiolata, lanceolata, integra, nervosa, glabra, versus basin cum petiolis sæpe bifida, erecta; *caulina* pauca, alterna, semiplexicaulia, lineari-lanceolata, angustiora, glabra, aut albide pilosa, margine connivente undulata subcurvata. *Flos* terminalis, luteus. *Prothollarum* ligulæ lineatæ, denticulatæ. *Pappi* sessilis, plumosus.

Hæc differt foliis angustioribus.

Obs. In Horto culta folia latiora imbut, hinc cum III. Hallero et Jacquino non pro peculiari specie declaramus.

Tota planta lacte candido scatet. Utilago florum genti chinensis succedaneum esse fertur. Linn. fl. rec. p. 267.

Bonum pecoribus pabulum planta præbet.

Flores apibus mel porrigunt.

1156. SCORZONERA *hispanica* caule ramoso, foliis amplexicaulibus integris, serrulatis. Linn. *Sp. pl.* 2. p. 1112.

Scorzonera major *hispanica*. Clus. *hist.* 2. p. 137.

Icon. Blackw. t. 406. Regnault. bot.

Term. Garten-Scorzonere. Spanische Haserwurz. Schwarzwurz.

Call. Scorzonere.

SYNGENESIA. POLYG. *Aquilegia*

Hab. In Hispania, Sibiria. Colitur in herbis et vitis ex quibus in agros migrat ubi passim quasi spontanea.

Flor. Junio, Julio. *Semina* Julio, Augusto matura.

Radix fusiformi-cylindrica, extus nigra, intus albida, carnosa. *Flores* lutei.

Pars *Scorzonera Radix.*

Radices nutrientes, leniter aperientes, in calidissimis; ubi passim Caffee succedaneum.

Flores melliferi.

1157. *SCORZONERA laciniata* foliis

nearibus dentatis acutis, caule erecto squamis calycinis patulo-mucronatis.

Sp. pl. 2. p. 1114. — *Pollich. fl. pal. n. 7.*

Scorzonera caule folioso ramoso foliis angustissimis, imis rariter semipinnatis, superioribus linearibus lanceolatis. *Hall. stir. helv. n. 12.*

Tragopogon laciniatum luteum. *Bauh. pin. 274. Prodr. 129.*

Icon. *Jacqu. fl. Austr. t. 356.* — *Schkuhr. bot. Handb. t. 215.*

Germ. Zerschlitzte Scorzonere.

Gall. *Scorzonere laciniée.*

Hab. In Sponhemia ad fluvium Noh inter Sponheim et Staudernheim. Circa May ad sepes post *Pollichium* vidi.

Flor. Majo, Junio. *Semina* Julio matura.

Radix fusiformi-cylindrica, fibrosa, extus albido-flavescens, intus alba. *Caulis* palmari pedalis, erectus, teretiusculus, striatus, villulosus, quandoque glaber, ramosus. *Ram.*

SYNGENESIA. POLYG. ÆQUALIS. 275

terni, pauci, erecti, longitudine caulis. *Folia* radicalia numerosa, profunde pinnatifida: segmentis oppositis alternisve, lineari-lanceolatis, remotiusculis, integris, angustis, glabriusculis; terminali longiori. *Caulina* alterna, infima radicalibus sublimilia, superiora lineari-lanceolata, integriuscula versus basin subpinnatifida, summa aut nulla aut unum alterumque lineare, integrum, angustissimum. *Flores* caulem ramosae terminantes, solitarii, parviusculi, lutei. *Calicis foliola* lanceolata, carinata, acuminata, infra apicem denticulo parvo, aculeolato, praedita. *Corollulae* subtus rubello striatae. *Semina* versus basin decemangulata. *Receptaculum* saepe aleis brevibus instructum.

158. *Planta* lacte albo scatens in horto culta, 2-3 pedalis.

Ad rariores omnino Florae nostrae cives pertinet.

CCCLVI. PICRIS.

Linn. Gen. pl. ed. Schreb. n. 1231.

Calyx duplex, exterior brevior, laxis, paucis, suboctodecimphyllus; interior longior subduodecimphyllus: *foliis* erectis, dense adpressis, carinatis, lanceolatis, dorso piloso-hispidis. *Corolla* composita, imbricata, uniformis: *Corollulis* hermaphroditis, numerosis. *Propria* monopetala, lingulata, linearis, truncata, quinque-dentata. *Calyx* fructiferus immutatus. *Semina* oblonga, gibbula, obtusa, transversim sulcata. *Pappus* subseffilis, plumosus.

158. *PICRIS Hieracioides* perianthiis laxis, caule angulato hispido, foliis sub-

Flor. Junio, *Julie.* *Semina* Augusto matur

Radix fusiformis, extus fusca, intus al
fibrosa. *Scapus* pedalis, sesquipedalis. ere
striatus, glaber, isophyllus, simplex, unifl.
Folia radicalia plura, scapo breviora, ol
ga, magis minusve profunde sinuato-dent
glabra, toto viridia. *Foliola* calycina imbr
glabra, quandoque villosiuscula, inæqualia
rolla flava. *Semina* pappo plumoso sessil
structa.

Obf. Planta pinnis variabilis, nunc Follis lud
tegriusculis, dentatis, obtusis, vel acutis,
profunde sinuato-dentatis, imo pinnatifidis,
berrimis, quandoque rariter hirsutis, Scapi
lycibusque glatis, hirsutulisve.

Nimis sane affinis sequenti speciei, a qua nesc
sit distincta.

Planta in horto culta, abiit in A, hispi
Conf. Schkuhr. bot. Handb. 3. p. 25.

1160. *APARGIA hispida* scapo nudo
floro calyce hirtio, foliis runcinato-
tatis hirtis, pilis furcatis. *Linn. ed. W*
3. p. 1550.

Leontodon hispidum calyce toto erecto
liis dentatis integerrimis hispidis;
furcatis. *Linn. Sp. pl. 2. p. 1124.* —
lich. fl. pal. n. 737.

Pieris caule nudo unifloro, foliis asperis
tatis. *Hall. stirp. helv. n. 25.*

Hieracium dentis leonis folio subaspe
Bauh. pin. 127.

Icon. Curtis. fl. Lond. t. 314. — *Oeder*
Dan. t. 862.

Germ. Rauhe Apargie.

Gall. Apargie hispide.

SYNGENESIA. POLYG. ÆQUALIS. 279

Hab. Utrunque in pratis, pascuis pomariis et agris passim frequens.

Flor. Julio, Augusto. *Semina* Septembri maturatione. ¶

Simillima in omnibus partibus præcedenti speciei, a qua mere differre videtur: *Scapo-folius* et *calycibus* multo hispidioribus. *Setis* sæpe 2-3-furcatis.

Obf. Obvia reperias individua ut in præcedente specie, digitalia, palmaria ad bipedalia.

Folia bonum pecoribus pabulum præbent.

Flores apibus melliferi.

1161. *APARGIA alpina* calyce toto erecto et scapo infra florem hispidis, foliis dentatis subpilosis, setis simplicissimis. *Hofst. Synops.* 423.

Leontodon alpinum. Linn. Murr. *Syst. veg.* ed. XIV. p. 715. ex Jacquino.

Icon. Jacqu. *fl. Austr.* t. 93.

Germ. Alpen-Apargie.

Gall. Apargie des alpes.

Hab. In ditionis Basileensis montosis prope *Muttenz im Engenthal t. Lachenalio.*

Flor. Julio, Augusto. *Semina* Septembri maturatione. ¶

Radix fusiformi-cylindrica, extus nigricans, intus albida, fibrosa. *Scapus* palmaris et altior, erectus, striatus, simplicissimus, uniflorus, sub flore tumidiusculus, asperiusculus, squamula una alterave instructus. *Folia radicalia* oblonga, utrinque attenuata, acuta, argute dentata, utrinque setoso-pilosa. *Setae* simplicissimae. *Calyx* ovali-oblongus, angulatus, erectus, higer, albide setoso-pilosus. *Corolla* lutea. *Co-*

SYNGENESIA. POLYG. EQUIS.

rollulae lingulatae, quinquedentatae. Receptaculum nudum. Pappus sessilis, plumosus.

Nota. Individuum ab Ill. Jaesquino l. c. exhibitum, nostratibus simillimum est.

1162. *APARGIA pyrenaica* Scop. l. pedunculis tumidis squamosis, calycibus villosis; foliis ovato-lanceolatis petiolatis subintegerrimis. *Gouan. Illustr.* p. 55. n. 2.

Taraxacum foliis integris dentatis, calyce hispido, pappo plumoso. *Hall. Stirp. helv.* n. 5.?

Leontodon pyrenaicum Villars fl. Delph. 1. p. 368.

Picris saxatilis Allion. fl. Pedem. n. 766.

Icon. Gouan. l. c. t. XXII. f. 1. 2. — Allion l. c. t. 14. f. 4.

Germ. Pyrenäische Apargie.

Gall. Apargie des Pyrenées.

Hab. In Ditione Basileensi ad montem *Vogelberg* nec non in Alsatiæ jugis altioribus *Belon* et *Bloutberg* t. *Lachenalio*.

Flor. Julio, Augusto. Semina Septembri matuta. 7

Plantam a b. *Lachenalio* in citatis locis reperi-
tam, non vidi, differe autem ab alpinis individuis; Sta-
tura majori, Foliis fere glabris, latioribus, Flore
majore. Conf. *Lachenal. Nov. Act. helvet. l.*
p. 276. qui eam ab *Apargia alpina* separat.

Obs. *LEONTODON hirtum* calyce toto erecto,
foliis dentatis hirtis; pilis simplicissimis. *Lin.*
Sp. pl. 2. p. 1123. Comprehendo veram *Lin-*
naei plantam abs me in *Castilia nova*, nunquam
in *Germania* lectam omnino ad *APARGIAM*. mi-
nime autem ad *THRINCIAM ROTHII* et *Will*

SYNGENESIA, POLYG. ÆQUALIS. 281

denowii pertinet. Calyx non octophyllus, octangularis sed polyphyllus cylindricus, Pappus sessilis, plumosus. Scapus digitalis, uniflorus, superne, bracteolatus. Folia runcinato-dentata, fetoso-hispida. Setae longæ, albæ, simplicissimæ.

Leontodon hirtum Leers. fl. Hernborn. p. 168. non ad hirtum Linnaei sed sine dubio ad *Apar-giam* hastilem, non ad *Thriniciae* Genus pertinet.

Apargia hirta. Schkuhr. bot. Handb. 3. p. 25. non ad *Leontodon hirtum* Linn. sed ut ex ejus Icone saltem iudico, ad variationem intermediam *Apargiae* hispidae et hastilis pertinet.

163. *APARGIA autumnalis* caule ramoso, pedunculis squamosis, foliis lanceolatis dentatis integerrimis glabris.

Leontodon autumnale. Linn. Sp. pl. 2. p. 1123. — Pollich. fl. pal. n. 736.

Hedypnois autumnalis scapo ramoso, pedicellis squamosis, foliis lanceolatis, dentato-pinnatifidis glabriusculis. Smith. fl. Brit. 2. p. 826.

Picris caule nudo brachiato, foliis glabris semipinnatis. Hall. stirp. helv. n. 28.

Hieracium foliis Coronopi. Bauh. pin. 128. Mapp. fl. Alf. p. 144.

Hieracium Chondrillæ folio glabro, radice succisa majus. Bauh. pin. 127. Mapp. fl. Alf. p. 143.

Hieracium III. f. *nigrum*. Tabernæmont, p. 491. Icon.

con. Schkuhr. bot. handb. t. 220.

erm. Herbst. *Apargie*.

all. *Apargie d'Autome*.

SYNGENESIA. POLYG. EQUIS.

Hab. Utrunque in pratis, locis graminosis, tuis, vils passim frequens.

Flor. Julio ad autumnum.

Radix alba præmorsa, fibrosa. **Caules** radice plures, palmares, pedales, sesquipedales, erecti, quandoque subdecumbentes, teretes, striati, glabri, ramosi, nudiusculi. **Rami** alterni, longi, nudi. **Folia radicalia** in orbem congesta, lineari-lanceolata, basi petiolo dilatata, pinnatifida; **pinnis** decurrentibus alternis oppositisve, angustis, erectis, integerrimis, glabris, lacte viridibus. **Folium** in basiguli rami lineari-lanceolatum, integerrimum parvum. **Rami** uniflori, sub flore in orbem squarulis parvis ad calycem usque obfessi. **Corolla** flava. **Corollulae** ligulae, subtus rubro-purpureis notatae. **Calycinae** squamae longae, imbricatae, glabrae, aut villosiusculae apice nigricantes. **Receptaculum** nudum. **Scapam radialium** Pappus sessilis, centralium stipitatus.

Nota. Folia radicalia cumprimis in solo sterili saepe lineari-lanceolata, integerrimula, vel lanceolata inciso-dentata, glabra.

Folia bonum pecoribus pabulum.

Flores apibus mel largiuntur.

1164. APARGIA incana scapo unifloro subnudo calyceque pubescente, foliis lanceolatis acutis subdenticulatis caulis scabris, pinnis multifidis. *Willden. Lich. Sp. pl. 3. p. 1549.*

Hieracium incanum foliis integerrimis, foliis denticulatis lanceolatis scabris, scapo unifloro. *Linn. Syst. nat. p. 522. — Michx. pal. n. 738.*

SYNGENESIA. POLYG. ÆQUALIS. 483

Hieracium montanum angustifolium nonnihil incanum. *Bauh. pin.* 129.

Hieracium montonum VI. *Clus. hist.* 2. p. 141.

Hieracium montanum angustifolium II. *Tabernæmont. p.* 495. *Icon.*

non. Jacqu. fl. Austr. 3. t. 287. (*sub Hieracio*).

erm. Graue Apargie.

all. Apargie pale.

h. In *Marggravatus superioris pratis montanis* prope *Oberweiler, Serlingen* et alibi passim. In *Alsatia* in cacumine montis *Bloutberg* prope *Marienkirch* ad mineras post *Lachenalium* vidi. In *Palatinatu* circa *Lautern* in pratis, aggeribus, pascuis, pomariis abunde t. *Pollichio.*

er. Junio, Julio. *Semina* Augusto matura. 7

Radix crassa, multiceps, extus nigricans, intus albida, fibrosa. *Scapi* ex radice sæpe plures, pedales, sesquipedales, erecti, teretes, simplices, aphylli, bracteola una alte vestiti, pilosuli, uniflori. *Folia radicalia* lanceolata, basi attenuata, obtusa, vel acutiuscula, obiter denticulata, crassiuscula, erecta, cinque tomentosa, incana: *Pili* albidi, apice latissimi. *Calyx* nigricans, hispidus. *Corolla* amara, flava. *Receptaculum* villis brevissimis vestitum. *Seminum* *Pappus* sessilis, plumosus.

CCCLVIII. S O N C H U S.

Linn. Gen. plant. ed. Schreb. n. 1233.

Calyx communis imbricatus, ventricosus: *lamis* plurimis linearibus, inæqualibus. *Corolla* composita, imbricata, uniformis.

SYNGENESIA. POLYG. AQUATIC.

lulis hermaphroditis numerosis, aequalibus. Petala monopetala, ligulata, linearis, truncata, quinqueidentata. Calyx fructiferus connatus in globum depressum, acuminatum. Semina longiuscula. Pappus capillaris, sessilis. Receptaculum nudum.

1165. SONCHUS *palustris* pedunculatus, calycibusque hispidis subumbellatis, foliis runcinatis, basi sagittatis. *Linn. pl. 2. p. 1116.*

Sonchus asper arborescens. Bauh. pin. Mapp. fl. Alf. p. 293.

Sonchus laevior austriacus, vel altissimus. Clus. hist. 2. p. 147.

Sonchus orborescens. Tabernemont. p. Icon mala.

Germ. Grofse Sumpf-Gänsefußel.

Gall. Laitron des marais.

Hab. Circa Argentinam in dem Ostwinke, ad aquas; ad Canalem in foveis humidis, in Mappo.

Flor. Junio, Julio. Semina Augusto maturant.

Plantam satis magnam, humanæ stature non sæpe superantem, in regionibus a Mappo citatis, non in aliis regionibus cis et transrhenanis frustra sivi. Vix spontanea in plagis rhenanis, et helvetiis in Belgii tamen, Angliæ, Galliæ et Hungariæ paludosis obveniens.

1166. SONCHUS *arvensis* pedunculatus, calycibusque hispidis, subumbellatis, foliis runcinatis basi cordatis. *Linn. Sp. 1116. — Pollich. fl. pal. n. 726.*

SYNGENESIA. POLYG. ÆQUALIS 284

Sonchus foliis amplexicaulibus, semipinnatis, serratis, calycibus hispidis. *Hall. stirp. helv. n. 23.*

Hieracium oleraceum. *Scopol. fl. Carn. n. 974.*

Sonchus repens, multis **Hieracium majus**.
Bauh. hist. 2. 1017.

on. Curtis. fl. Lond. t. 265. Oeder. fl. Dan. t. 606.

erm. Acker-Gänsefistel.

all. Laitron des champs.

ab. Utrinque in agris, vineis et oleraceis,
Sequitur præsertim terram argillaceam.

lor. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. \mathcal{P}

Radix subcylindrica longe lateque repens, diffusua. **Caulis** 2-3 pedalis et altior, erectus, bifidus, teretiusculus, lævis, lactescens, summe ramosus. **Folia** alterna, amplexicaulia, cordata, runcinata, lævia, margine spinu-
mollibus prædita. **Flores** subumbellati. **Pet-
ioli** longiusculi, teretes, cum **Calyce** setis
buligeris luteis vestiti. **Corolla** sat speciosa,
et. **Semina** oblonga, fulcata. **Pappus** pilo-
s, sessilis.

Planta ob radicem nimium reptatricem, difficile
radam, in agris hortisque et vineis præsertim noxia,
si studio diminuenda.

Bonum et acceptum suis pabulum.

Flores melliferi.

167. SONCHUS lævis foliis amplexicaulibus dentatis integris et semipinnatis, calycibus lævibus. *Hall. stirp. helv. n. 21.*

SYNGENESIA, FOLYG. EQUALE

Sonchus oleraceus *a. laevis*. Linn. Sp.
2. p. 1117. — Pollich. fl. pal. n. 727.

Sonchus laevis laciniatus latifolius. Bauh. p.
124. — Mapp. fl. Alf. p. 293.

Sonchus laevis III. IV. V. *Tabernaemontana*
p. 499. Icon.

β. *Sonchus laevis minor paucioribus laciniis*
Bauh. pin. 124. — Mapp. fl. Alf. p. 293.

Icon. α. Curtis. fl. Lond. t. 123. — Black
herb. t. 130.

Germ. Glatte Saudistel. Glatte Gänsedistel.
Gall. Laitron sans épins. Laitron doux.

Hab. Utrinque, in agris, vineis, oleraceis
ruderalis passim abunde.

Flor. Junio, Julio. *Semina* Augusto matura.

Radix simplex, fibrosa, albida, lactescens.
Caulis 1-3 pedalis, erectus, fistulosus, laevis,
teretiusculus, superne ramosus. *Folia* alternata,
amplexicaulia, inferiora pinnatifida, inaequaliter
sinuato-denticulata, terminali triangulata,
majori, omnia, laevia, glauca. *Pedunculi* teretes,
bri, aut tomentosi, per aetatem nudj. *Capitula*
laevis, glaucus. *Corolla* lutea. *Semina* per longum
tem observata, duobus denticulis truncatis
obliquis instructa. *Pappus* sessilis pilosus.

Planta glaberrima valde variabilis, magnitudine
figura et foliorum respectu.

Pharm. SONCHI Herba.

Lactescens, saponacea, amara, subnarcotica,
hodie in medicina recepta.

Bonnum anseribus suisque pabulum, nisi quod
nem ingrato inficit sapore.

Nimiam saepe ob abundantiam pessimum agrosti
livitium, quod tamen abscissione ante florescentiam
cum non difficile diminuitur.

168. SONCHUS asper foliis amplexicaulibus rigidiusculis integris aut semipinnatis, dentatis, calycibus laevibus. *Hall. stirp. helv. n. 22.*

Sonchus oleraceus. γ. Linn. Sp. pl. 2. p. 1117. Pollich. fl. pal. n. 227.

Sonchus asper laciniatus folio dentis leonis. Bauh. pin. 124. Mapp. fl. Als. p. 293.

Sonchus asper L. Tabernæmont. p. 468. Icon.

Sonchus asper non laciniatus. Bauh. pin. 123. — Mapp. fl. Als. 293.

Leon. = Blackw. herb. t. 39. — Oeder. fl. Dan. t. 843.

erm. Rauhe Gänsefistel.

all. Laitron épineux.

ab. Cum præcedente utrinque passim.

or. et Semina cum ea matura. ☉

Planta aspera, variabilis. Distinguitur a præcedente. *Caule aspero. Foliorum* dentibus asperis pungentibus. *Seminibus* ad latera membranis appendiculatis, quo signo optime accuratissimus *Schkuhr* distinxit.

In horto culta, constans, non abiit in præcedentem speciem.

Qualitates præcedentis forsan magis narcoticas habet.

169. SONCHUS alpinus foliis amplexicaulibus cordato sagittatis runcinatis, sinuato-dentatis glabris, lobo terminali deltodeo maximo, floribus racemosis, pedunculis hispidis.

Sonchus alpinus pedunculis bracteisque longioribus hirsutis, floribus racemosis, foliis

LYNGENESIA. POLYG. ACUT.

runcinatis basi sagittatis. *Frœlich in l
bot. Annal. 1. p. 26.*

Sonchus montanus Lamark. encycl. 3. p

*Sonchus foliis amplexicaulibus petiolat
angularibus, petiolo pinnato, floribu
catis. Hall. stirp. helv. n. 20.*

*Sonchus laevis laciniatus*coeruleus. f.
chus alpinus coeruleus. Bauh. pin.*

*Lactuca montana latifolia laciniata flore
ruleo. Mapp. fl. Alf. p. 160.*

*Sonchus laevis quartus flore coeruleo.
bernæm. p. 500. Icon.*

Icon. Oeder. fl. Dan. t. 182.

Germ: Alpen-Gänsefistel.

Gall. Laitron des alpes.

*Hab. In Marggraviatus superioris monte
chen et Feldberg. In Monte Muteto et N
serfallen. t. C. Bauhino. In Alfatiae m
Ballon.*

*Flor. Julio, Augusto. Semina Augusto
tembri matura. 7*

*Caulis 3-4 pedalis et altior, erectus,
cis rubellis subhirsutis præditus, fistulosus,
plex. Folia alterna, amplexicaulia, corda
hastata, runcinata, inaequaliter sinuato-de
culata, glabra, subtus pallida, lobo termi
maximo deltoideo, acuminato; *suprema* an
stiora, cordato-lanceolata, acuminata, arg
dentato-ciliata. Flores caulem terminantes,
cemosi. Pedunculi subunciales, subuniflori, c
suprema caulis parte piloso-hispidi fusco-
belli, squamula una alterave angustissima,*

SYNGENESIA. POLYG. ÆQUALIS. 289

petali pilosa præditi, *Bractea* lineari, angusta, folia, suffulti. *Calyx* fusco-rubellus, pilosulus. *Corolla* mediocris, coeruleo-rubella. *Semina* valil-linearia, compressa. *Pappus* sessilis, pilosus.

Variationem Corollis niveis in pyrenæis vidi.

Nota. b. Linnaeus sub Soncho canadensi Sp. pl. 2. p. 1115. sine dubio longe aliam plantam quam Sonchum alpinum nostrum comprehendebat ut ex ulteriori descriptioni patet: Folia, inquit, Sonchi floridani, Pedunculi longi — Flores coerulei, magnitudine Sonchi palustris. Sonchus noster alpinus, Pedunculos habet breves, vix unciales, Folia minime Sonchi floridani, Flores coeruleo-rubellos, Soncho palustri duplo minores.

Sub Soncho alpino Linn. Sp. pl. 2. p. 1117. species diversae militabant, alter verus noster alpinus. pedunculis piloso-hispidis, alter lapponicus annuus aut biennis, pedunculis squamosis, quem sub alpino descripsit et delineavit. Smith. Icon. pl. edit. p. 21. t. 21.

CCCLIX. L A C T U C A.

Linn. Gen. pl. ed. Schreb. n. 1234.

Calyx communis imbricatus, cylindricus. *Sepalis* plurimis, acuminatis, margine membranaceis. *Corolla composita*, imbricata, unilobis: *Corollulis* hermaphroditis plurimis, aequalibus. *Propria* monopetala, ligulata, truncata, quadri-quinquedentata. *Calyx* fructiferus unilobis, ovato-cylindricus. *Semina* ovata, acuminata, lævia, compressa. *Pappus* capillaris: *Stipite* longo, inferne attenuato. *Receptaculum* nudum.

Flora Bod. Tom. III,

T

1170. *LACTUCA fativa* foliis rotunda
caule corymboso. *Linn. Sp. pl. 2. p. 111.*

Lactuca fativa. Bauh. pin. 122.

β. *Lactuca capitata. Bauh. pin. 123.*

Germ. Gartensalat. β. Kopfsalat.

Gall. Laitue cultivée.

*Hab. Colitur in hortis, oleraceis et
neis gum β.*

*Flor. Julio, Augusto. Semina Augusto, Se-
tembri matura. ☉*

Plures sane variationes in hortis coluntur.

Pharm. LACTUCÆ Semina.

Ufus plantae culinaris notissimus.

Semina aviculis accepta.

1171. *LACTUCA Scariola* foliis verti-
libus carina aculeatis. *Linn. Sp. pl. 2. p.
1119. — Pollich. fl. pal. n. 728.*

*Lactuca foliis semipinnatis, nervo spinoso.
Hall. stirp. helv. n. 14.*

*Lactuca sylvestris, costa spinosa. Bauh. pin.
123. — Mapp. fl. Alf. p. 160. — Mor-
hist. 3. p. 58. Sect. 7. t. 2. f. 16.*

Germ. Wilder Lattig.

Gall. Laitue Scariole.

*Hab. Utrinque passim in ruderatis, ad agrorum
vias, in vineis et collibus asperis. Circa
Carlsruhe am Landgraben versus Müll-
berg rarius. In montosis dumetosis asperis
calcareo-argillaceis retro Durlach, Grä-
zingen, prope Weingarten, Bruchsal, Nuß-
loch et alibi. In Margraviatu superiore ad
Kaiserstuhl frequens, nec non in vineis
prope Badenweiler, Oberweiler et alibi.*

SYNGENESIA. POLYG. ÆQUALIS. 291

Flor. Junio, Julio, Augusto. **Semina** Augusto, Septembri matura. ♂

Caulis 2-3 pedalis et altior, erectus, laevigatus, setis raris passim præditus, lactescens, ramosus. **Folia** alterna, amplexicaulia, hastata, uncinato-pinnatifida, verticalia, seu obliqua, non horizontalia, dentato ciliata, costa media longitudine aculeata; superiora lanceolata. **Flores** paniculati, parvi, sulphurei. **Pappus** longepilatus, pilosus.

Planta narcotica sequentis speciei vires suspicari permittit.

1172. LACTUCA virofa foliis horizontalibus carina aculeatis dentatis. *Linn Sp. pl.* 2. p. 1119.

Lactuca foliis ovatis patulis, nervo spinoso. *Hall. stirp. helv.* n. 15.

Lactuca sylvestris, odore viroso. *Bauh. pin.* 123. — *Mapp. fl. Als.* p. 160. — *Morif. hist.* 3. p. 58. *Sect.* 7. t. 2. f. 16.

Don. Schkuhr. bot. handb. t. 217.

erm. Betäubender-Lattic.

Call. Laitue virose.

Tab. In Margraviatu superiore am *Kaiserstuhl* Brisgoviae, prope *Altbreisach*, *Burgheim*, *Sponck*, *Rothweil* et *Limburg* in rupibus apricis ex basalte, et lava constructis. In ditione Basileensi inter *Muttenz* et *Gempen* juxta montanam semitam im *Stierenwald-boden*. In Alsatia prope Argentinam *vor dem Steinstrasserthor auf der Contrescarpe* et alibi. In Alsatia superiore retro *Rappolsweiler*, et prope *Marienkirch* ad metallifodinas in rupibus montosis. In Sponhemia

892 SYNGENESIA, POLYG. ~~ÆQUALIS~~

ad euprifodinam prope *Fischbach* in rupi
fissuris copiose.

Flor. Junio, Julio. *Semina* Julio, Augu
matura. ♂

Affinis præcedenti, sed major sæpe 4-5 p
dalis, et crassior. *Folia* multo majora, latior
non verticalia, sed fere horizontalia: inferius
non runcinata, sed sinuato-inæqualiter den
ta, subundulata.

Pharm. LACTUCA VIROSA Herba:

Planta præcedente magis virosa, narcotica, lact
cens, lacte amaro, acri, viroso, in Hydrope a no
nullis laudatur. Collin. obs. circa morbos. 178

Nota. A. pharmacopoeis non cum Soncho aspe
aut laevi confundenda.

1173. *LACTUCA augustana* foliis int
gris dentatis acute hamatis, costa læ
Allion. Mifs. Taur. 5. p. 72. n. 74.

Lactuca longo et valde angusto folio. *Bau
hist. 2. p. 999.*

Icon. Allion. fl. Pedem. t. 52. f. 1.

Germ. Augustaner Lattic.

Gall. Laitue Augustane.

Hab. In Marggraviatu superiore in Brisgov
prope *Altbreisach* in agris. Circa Basilea
in agris cultisque passim non infrequens.

Flor. Junio, Julio. *Semina* Augusto matura. (

Radix fusiformis, fibrofula, recta. *Caul*
cubitalis, sesquicubitalis, erectus, laevis, firi
tus, ramosus: *Rami* alterni, erecti, subdivi
multiflori. *Folia* caulina alterna, semiamplex
caulia, basi hamata, margine inæqualiter argu
spinuloso-denticulata, utrinque cum costa m
dia glaberrima, lactescentia, acuminata, saturat

india, superiora minora, angustiora, minus lenticulata. Flores breviter pedunculati, paniculati. Flosculi in singulo flore subduodecim, sulphurei. Semina cinereo albida.

Differt ab affini *Lactuca virosa*, Costis foliorum aculeatis, foliis inferioribus non sinuatis. Odore viroso.

1174. *LACTUCA saligna* foliis hastato-nearibus sessilibus, carina aculeatis. Linn. *Sp. pl.* 2. p. 1119. *Pollich. fl. pal.* n. 729.

Lactuca foliis imis semipinnatis, superioribus amplexicaulibus hamatis, nervis spinosis. Hall. *fl. stirp. helv.* n. 13.

Lactuca sylvestris altera, angusto saligno folio, costa albicante. Rupp. *fl. Jen.* 167. ed. Hall. t. 4.

Chondrilla viscosa humilis. Bauh. *pin.* 130. *Prodr.* p. 68. Icon.

Icon. Jacqu. *fl. Austr.* t. 250.

Ger m. *Weidenblättriger Salat.*

Gall. *Laitue a fevilles des Saules.*

Hab. In Brisgovia am *Kaiserstuhl* in collibus vulcanicis, asperis, nec non ad vinearum marginalia prope *Burgheim am Lützelberg* et *Limbürg* passim. In Ditione Basileenli in agris ad pontem *Birsæ*, nec non in agris inter *Oberwieler* et *Neuwjeler* t. *Lachenalio*. In Palatinatu in aggeribus rhenum versus inter *Oppenheim* et *Gernsheim* poli *Pollichium* vidi. In pratorum fossis ad *Salinas* circa *Dürkheim* t. *Pollichio*.

Flor. Julio, Augusto. Semina Augusto, Septembri matura. ☞

294 SYNGENESIA. POLYG. AQUATILIS.

Radix teres albido-lutescens, fibrosa, *Caulis* sesquipedalis, bipedalis, erectus, simpliciusculus, quandoque a radice ramosus, tere glaberrimus, nitidus, albido-cinereascens. *Ran* alterni, patentiusculi. *Folia* alterna, verticali sessilia, hastata, sublinearia, glabra, integra inferiora interdum rariter profunde dentata; ac argute pinnatifida, glauco-viridia, costa medi albida subinermi, quandoque, et quidem praesertim inferiora, subtus aculeis cuspulis albidimunita. *Flores* sparsi, ramulosque terminantes. *Flosculi* in flore subtredecim, ligulati, *Ligulae* quinquedentatae, patentissimae, pallide sulphureae, subtus fusco-purpurascens. *Pappus* breviter stipitatus, pilosus. *Semina* utriusque acuminata, nigricantia.

Tota planta lactescit.

1175. *LACTUCA perennis* foliis linearibus dentato-pinnatis: laciniis sursum dentatis. *Linn. Sp. pl.* 2. p. 1120. *Pollich. fl. pal.* n. 730.

Lactuca foliis glabris pinnatis, caule brachiato. *Hall. stirp. helv.* n. 16.

Lactuca perennis humilior, flore coeruleo. *Mapp. fl. Als.* p. 160.

Chondrilla coerulea. *Tabernæmont.* p. 486 *Icon.*

Chondrilla coerulea latifolia laciniata. *Bauh. pin.* 130.

β. Flore albo.

Icon. *Miller. Dict.* t. 157.

Ger m. *Blauer Lattic.* *Bergsalat.*

Gall. *Laitue vivace.*

HYNGENESIA. POLYCL. ÆQUALIS. 295

ss. In ditione Basileensi in collibus argillofis
t. *Joh. Bauhino*. hodie exul. In vineis lapido-
fis prope *Neustadt* t. *Zeyhero*. In Alfatiz
collibus asperis prope *Buchsweiler*, in monte
Pafsberg in agris t. *Mappo*. In Palatinatu
in vineis circa *Umstein Dürkheim*, in col-
libus sylvaticis circa *Neustadt an der Hardt*
post *Pollichium* vidi. In Sponhemia prope
Winningen in rupibus præaltis schiffaceo-
argillaceis, nec non circa *Naumburg, Fisch-*
bach et *Oberstein* cum var. β . in rupibus
dumetosis gramineis frequens.

lor. Junio, Julio. *Semina* Augusto matura. γ

Caulis sesquipedalis, bipedalis et altior,
rectus, sulcatus, glaber, ramosus. *Rami* al-
mi, patentiusculi. *Folia* alterna, amplexicau-
li, pinnatifida; *laciniis* lineari-lanceolatis, ra-
ter dentatis, vel integris glabris; *summa* inte-
na, glabra, glauco-viridia. *Flores* paniculati,
eciosi, coerulei, ex coeruleo-violacei, ra-
mis albi. *Semina* longe stipitata.

Folia plantæ tenera, in Sponhemia, mensibus Fe-
uario et Martio ad *Acetaria* sapidissima colliguntur.

Flores apibus mel porrigunt.

CCCLX. C H O N D R I L L A.

Linn. Gen. pl. ed. Schreb. n. 1235.

Calyx communis calyculatus, cylindræ-
; *squamis cylindri* plurimis, parallelis, linea-
tus, æqualibus; *baseos* paucis, brevissimis.
rolla composita, imbricata, uniformis: *Co-*
llulis hermaphroditis, æqualibus. *Propria* mo-
petala, ligulata, linearis, truncata, 4-5-den-
a. *Calyx fructiferus* cylindræceus, oblongus.

SYNGENESIA POLYG. EQUALITER

*Semina ovata, compressa, muricata. Pappus
pilleris, Stipite longo, superne attenuato.
receptaculum nudum.*

1176. *CHONDRILLA juncea* foliis rac-
calibus runcinatis, caulinis linearibus
tegris, caule ramoso, junceo.

Chondrilla juncea. Linn. *Sp. pl.* 2. p. 112
— Pollich. *fl. pal.* n. 731. — Hall. *fl.
helv.* n. 17.

*Chondrilla juncea viscosa, arvensis, quæ
prima Dioscoridis.* Bauh. *pin.* 130.
Mapp. fl. Als. p. 72.

Chondrilla juncea. Tabernæmont. p. 487.
Icon. Jacqu. fl. Austr. t. 427. — Schkuhr. *fl.
Handb.* t. 218.

Germ. Chondrille, Binsenartige Acker-Chondrille.

Gall. Chondrille joncière.

Hab. Utrunque passim. Circa *Carlsruhe* jux-
dem *Landgraben* versus *Mühlberg*; prope
Ettlingen in arvis ad viam publicam co-
piose. Retro *Durlach* et *Grözingen* in vi-
neis passim. In *Brisgovia* am *Kaiserstuhl*
prope *Altbreisach* in collibus ex cinere *Vu-*
cani, et *Lava*, *Augite*, et *Zeolite* con-
stitiis abunde. In *Margraviatu* superiori pa-
ssim in agris et vineis, præsertim argilloso
In *Alsatiæ* arvis, agris et vineis hinc inde
ut in *Palatinatu*.

Flo r. Junio, Julio. Augusto *Semina* matura.

Caulis 3-4 pedalis et altior, erectus, ba-
pilofo hirsutus, superne glaberrimus, ramosus
Rami alterni, copiosi, juncei. *Folia* radicali

SYNGENESIA. POLYG. ÆQUALIS. 297

malinaque infima pinnatifida, sinuato-dentata, scabra, caulina superiora ramæaque, linearia, glabra, viridi-glaucæ, integriuscula, vel subdenticulata. *Flores* terminales lateralesque, sessiles, parvuli, lutei.

Tota planta lacte viscido, albo turget.

Magnum sæpe præsertim in vineis et agris argillæ vitium, probe eradicandum.

CCCLXL P R E N A N T H E S.

Linn. Gen. pl. ed. Schreb. n. 1236.

Calyx communis calyculatus, cylindraceus, lobis: *squamis cylindri* numero corollularum; lobis paucis, inæqualibus, brevissimis. *Corolla* imposita fere ex simplici orbe flosculorum: *corollulis* hermaphroditis, 5-8 et pluribus, inæqualibus, in orbem simplicem positis. *Procorolla* monopetala, ligulata, truncata, quadrilobata. *Calyx* fructiferus cylindraceus, ore brevissime connivens. *Semina* cordata. *Pappus* capillaris, sessilis. *Receptaculum* nudum.

Obs. *P. muralis* pappum habet stipitatum.

1177. *PRENANTHES tenuifolia* flosculis quinis, foliis linearibus amplexicaulis integris, caule ramoso.

Prenanthes tenuifolia foliis linearibus integerrimis. *Linn. Sp. pl. 2. p. 1120.*

Prenanthes angustifolius, flore purpureo. *Vaill. act. 1721. p. 193.*

Chondrilla angustissimo longissimo integroque folio. Juss. Act. 1709.

Icon. Allion. fl. Pedemont. t. 33. f. 2.

Germ. Schmalblättrige Prenanthe.

Gall. Prenanthe à sévilles étroites.

Hab. In montosis umbrosis sylvaticis humis prope *Frauenalb*, auf dem *Tobel*, auf der *Herrenwiese* et auf dem *Kaltenbrunn*. In convallibus im *Oppenauer*, *Griechen* et *Petersthal* passim non infrequenter. In sylvis montanis circa *Muttenz* t. *Lachnatio*.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. 7

Affinis sequenti speciei, a qua facile distinguitur. *Caulis* graciliore, ramosiore. *Folii* linearibus, integris, aut saltem ad basin alterave denticulo instructis, multo angustioribus et longioribus. *Flores* ut in *P. purpurea*.

Nota. Planta hucusque Collectores Florae Germaniae fecisse videtur, pro patria nova.

Obs. In Horto botanico Carlsruhano culta, nullam mutationem passa est.

In omnibus partibus exacte cum specimenibus Galliae meridionali et Hispaniae observatis, ut et cum Icone allegata, convenit.

1178. *PRENANTHES purpurea* flosculis quinis, foliis lanceolatis, denticulatis
Linn. Sp. pl. 2. p. 1121. — Pollich. f. pal. n. 732.

Prenanthes foliis amplexicaulibus, petiolatis, lanceolatis, serratis. *Hall. stirp. hel. n. 19.*

Chondrilla Sonchifolio, flore purpurascens minor. *Mapp. fl. Alf. p. 72.*

SYNGENESIA. POLYG. ÆQUALIS. 299

Lactuca montana, purpureo - coerulea minor. *Bauh. pin.* 123.

Chondrilla Sonchifolio flore purpurascente major. *Mapp. fl. Alf. p.* 72.

Lactuca montana purpureo - coerulea, major. *Bauh. pin.* 123.

n. *Jacqu. fl. Austr. t.* 317.

m. *Purpurfarbige Prenanthe.*

b. Utrinque passim in sylvis montosis rupesribus humidiusculis abunde.

r. Julio, Augusto. *Semina* Augusto, Septembri matura. ¶

Caulis 2-3-5 pedalis, erectus, teres, glabrus, superne ramoso-paniculatus. *Folia* altera oblonga, subtus glauca, glabra, *inferiora* sessilia, denticulata; *superiora* amplexicauminus denticulata, quandoque integriuscula. *Flores* copiosi, paniculati, cernui. *Flosculi* paucissimi quinque, quandoque quatuor, rarius sex, purpureo-rubentes, interdum pallide albi, rarius albi. *Pappus* sessilis.

Nota. Bonum in montosis sylvaticis et abundans praesertim sylvae nigrae pecoribus pabulum.

Flores apibus melliferi.

179. PRENANTHES muralis flosculis quinis, foliis runcinatis. *Linn. Sp. pl.* 2. p. 1121. — *Pollich. fl. pal. n.* 733.

Prenanthes foliis ferratis pinnatis: pinna suprema triangulari, trilobata. *Hall. stirp. helv. n.* 18.

Chondrilla Sonchifolio, flore luteo pallescente. *Mapp. fl. Alf. p.* 73.

900 SYNGNESIA. POLYG. ÆQUALI

Sonchus lævis laciniatus muralis, parvis
ribus. *Bauh. pin.* 124.

Sonchus sylvaticus quartus. *Tabernaem.*
501. *Icon.*

Icon. Curtis. fl. Lond. t. 306. — *Oeder.*
Dan. t. 509. — *Schkuhr. bot. handb. t.* 2.

Germ. Gelbe Mauerprenanthe.

Gall. Prenanthe des murs.

Hab. Utrique in sylvaticis rupibus, muris
sylvis cæduis passim frequens.

Flor. Junio, Julio. *Semina* Julio, Augusto
matura. 7.

Caulis 1-2 pedalis et altior, erectus, ter-
glaber, ramosus. *Rami* alterni, paniculati.
lia alterna, amplexicaulia, pinnatifida, ac-
nata, glabra, lacinia suprema triloba, trian-
lata, margine sæpe rubella. *Stipulae* parvae,
pedunculorum basin. *Flores* paniculati, lutei,
parviusculi. *Pappus* stipitatus.

Bonum pecoribus alimentum.

Flores apibus mel concedunt.

1180. *PRENANTHES viminea* foliorum
ramentis cauli adnatis. *Linn. Sp. pl.* 2.
1120.

Prenanthes foliorum laciniis extrorsum de-
tatis, ramentis foliaceis cauli ramisque
adnatis. *Linn. vir. Cliff.* 78.

Lactuca viscosa, caule foliis obducto. *Toe-
nef. inst.* 472.

Chondrilla viscosa, caule foliis obduc-
Bauh. pin. 130.

Chondrilla viminea viscosa monspelia
Bauh. prodr. 68.

EGGENESIA. POLYG. ÆQUALIS. 302

Jacqu. fl. Austr. t. 9. Allion. fl. Pedem. 52. f. 2.

Klebrige Prenanthe.

Prenanthe visquese.

In monte Muteto. t. Lachenalia.

Julio. Semina Augusto matura. ♂

radiculis sesquipedalis, cubitalis, erectus, glaber, albidus, superne ramosus. *Rami*, patentes. *Folia* alterna sessilia, caulina *superiora* et media, profunde pinnatifida, laciniabus infimis linearibus, obtusis, integris arcte adnatis, reliquis lineari-falcatis, integris, quandoque rariter denticulata. *Infima* recta elongata; *suprema* rameaque *media*, linearia, acuta, integriuscula, basi *inferiore* appendiculo cauli adnato. *Flores* teretes, lateralesque, subsessiles, lutei. *Flos* quinque. *Semina* lanceolata, nigra, subviscosa. *Pappus* capillaris.

1. Ad rarissimas Florae cives omnino pertinet, planta alias magis in regionibus Austriæ Galliæ et Hispaniæ calidioribus obveniens.

2. Angularis omnino ob foliorum appendices virides albo arcte adnatos.

CCCLXII. LEONTODON.

nn. Gen. plant. ed. Schreb. n. 1237.

Calyx communis imbricatus, oblongus: *lobis interioribus* linearibus, parallelis, æqualibus: *exterioribus* paucioribus sæpe reflexis asinis. *Corolla* composita, imbricata, unilobis: *Corollulis* hermaphroditis, numerosis, subulibus. *Propria* monopetala, ligulata, linearis, truncata, quinquedentata. *Calyx* fructu-

208 SYNGENESIA. POLYG. EQUALIS.

ctiferus oblongus, rectus, tandem reflexus.
mina oblonga, scabra. *Pappus* capillaris, fir-
tatus. *Receptaculum* nudum, punctatum.

1181. LEONTODON *Taraxacum*
ce inferne reflexo, foliis runcinatis di-
ticulatis lævibus. *Linn. Sp. pl.* 2. p. 111.
Pollich. fl. pal. n. 734.

Taraxacum calycibus glabris, styaminibus
reflexis. *Hall. stirp. helv.* n. 56.

Dens Leonis latiore folio. *Bauh. pin.* 126.
Mapp. fl. Alf. p. 91.

Dens Leonis I. II. *Tabernaemont. p.* 481. *Id.*

8. Dens Leonis angustiore folio. *Bauh. p.*
126. — *Mapp. l. c.* p. 92.

Icon. *Curtis. fl. Lond.* — *Bulliard. herb.* t. 21.
— *Regnault. bot.* — *Schkuhr. bot. hand.*
t. 219.

Germ. Löwenzahn. *Pfaffenröhrlein.*

Gall. Dent de Lion. *Pissenlit commune.*

Hab. Utrinq̃ue in hortis, cultis, aggeribus
vineis, pratis, pascuis, sylvis, sepibus, ri-
deratis, ad vias passim frequens.

Flor. Martio, Aprili, Majo. *Semina* Majo
Junio matura. 7.

Radix cylindrica, albido-cinerea, rugis tra-
versis subannulata, subramosa, fibrosa, carnos-
lactescens. *Scapi* ex radice sæpe plures, di-
tales, palmares, pedales et altiores, erecti
glabri, fistulosi, uniflori. *Folia* radicalia plura
congesta, petiolata, runcinata, laciniis recti-
acutis, triangularibus, dentatis, glabriusculi
quandoque villosis. *Flores* lutei.

INGENESIA. POLYGE. EQUALIS. 303

Tota planta lactescit.

Herb. TARAXACI Radix; Herba.

amara, aperientes, diureticas, resolventes, et rores vires habet.

Folia juniora in cibum et ad Acetaria adhibentur.

Stemibus bonum in pascuis pabulum.

Stemibus apibus mel praebent.

2. *LEONTODON salinum* calyce toto erecto, laevi, foliis carnosiss, oblongis, obtusis, denticulatis, glabris, scapo unifloro laevi. *Pollich. fl. pal. n. 735.*

Leontodon lividum foliis lanceolatis nudis, squamis calycinis exterioribus ovatis adpressis. *Waldst et Kitaibl. pl. rar. Hung. 2. p. 120.*

Leontodon palustre squamis calycinis exterioribus brevioribus erectis, ovatis, foliis sinuatis dentatis, glabriusculis. *Smith. fl. Brit. 2. p. 823.*

Leontodon erectum calyce toto erecto marginato, foliis pinnatis acutis denticulatis, scapo unifloro, superne villosa. *Hoffm. fl. germ. p. 278.*

Apargia salina. *Flora der Wetterau. 3. p. 146.*

Hedypnois paludosa. *Scopol. fl. Carn. ed. 2. p. 100. tab. 48.*

Leon. Waldst et Kitaibl. l. c. t. 115.

Herb. Salz-Löwenzahn.

Call. Dent de Lion salée.

Tab. In ditionis Basileensis pratis prope Michelsfelden. t. *Lachenalio.* In Palatinatu in pratis ad Salinas circa Dürkheim t. *Polli-*

804 SYNGENESIA POLYÆ AQUATILIS

chlo; nec non in pratis humidis ad Salin
prope *Krautzenach* in Spandemia.

Flo r. Majo, Junio, *Semina* Junio, Julio
tura. ♀

Scapi ex radice sæpe plures, digitales, es
eti, fistulosi, teretes, rariter tomentosuli, us
flori. *Folia* radicalia plura, lanceolata, sinu
dentata, glabra, carnosæ, longitudine sca
Calycinae squamæ exteriores, ovali-lanceo
latae, 10-12, erectae, adpressæ, glabræ, inter
res parallelæ, lineares, erectae; angustæ, le
giore, 14, glabræ. *Flor* lætus.

CCCLXIII. HIERACIUM.

A. Linn. Gen. plant. ed. Schreb. n. 1238.

Calyx communis imbricatus, ovatus: squ
mis pluribus linearibus, valde inaequalibus lon
gitudinalibus et incumbentibus. *Corolla* com
posita, imbricata, uniformis: *Corollulis* he
maphroditis, numerosis, æqualibus. *Propr*
monopetala, ligulata, linearis, truncata, qui
quedentata. *Calyx* fructiferus connivens, ov
tus. *Semina* obtuse tetragona, brevissima. *Papp*
capillaris, sessilis. *Receptaculum* nudum.

† *Scapo unifloro.*

1183. *HIERACIUM aureum* scapo us
floro recto, supernè hirsuto, foliis fo
spatulatis sinuato-dentatis obtusiuscul
calyce erecto hispido, flore aureo.

Hieracium aureum foliis runcinatis cau
subunifolio, calyce hispido, radice pr
mor

ÆGENESIA. POLYG. ÆQUALIS. 305

morfa. *Villars. fl. Delph.* 3. p. 96. t. 33.

Scopol. fl. Carn. 2. p. 104.

æontodon aureum foliis runcinatis caule
subunifolio, calyce hispido. *Linn. Sp.*
pl. 2. p. 1122. — *Jacqu. vind.* 268.

araxacum calycibus hirsutis, squamis ere-
ctis. *Hall. stirp. helv.* n. 57.

ens Leonis foliis extremo subrotundis, flo-
ribus aureis. *Raj. eur.* 399. *Sequ. pl. ver.*
3. p. 265.

1. *Jacqu. fl. Austr.* t. 297.

m. *Goldgelbes Habichtskraut.*

l. *Eperviere d'orée.*

2. In graminosis montis *Feldberg* Brisgo-
viæ palcuis infrequens. In Principatu *Für-
stembergenſi* in montosis prope *Krügenthal*
cum *Anemone narcissiflora* L. nuper vidi.

3. Junio, Julio. *Semina* Julio, Augusto ma-
tura. 7

Radix præmorfa, fibris aucta, extus fusco-
icans, intus alba. **Scapus** ex radice solita-
r, simplex, aut nonnulli, semipedales et lon-
ges, erecti, fistulosi, teretes, striati, infer-
glabriusculi, superne hispidi, squama foliacea
altera pluribusve, muniti, uniflori. **Folia**
icalia sæpe plura, oblongo-spathulata, ver-
basin sensim angustiora, subinæqualiter sinua-
toſſe dentata, obtusa, utrinque glabriuscula,
viridia. **Flos** speciosus, intus ex flammeo au-
ro, extus croceus. **Calycinae** squamæ interiores
longo-lanceolatae, acutæ, erectæ, saturate ex
ro viridescentes, pilis asperis atris, hispi-
bus; exteriores 5-7, interioribus multo brevio-
res. **Semina** linearia, nigra. **Pappus** sessilis, ca-
ris, niveus. **Receptaculum** nudum.

306 SYNGENESIA. POLYG. ÆQUALIS.

Ob pappum capillarem sessilem omnino a L. todo removendum erat.

Obs. Individua in Pyrenæorum pascuis a me lecta stratibus hispidiora, imo foliis utrinque his instructa sunt.

Planta ob flores amoenos elegans.

184. *HIERACIUM alpinum* foliis longis integris dentatis scapo subnudo, floro, calyce piloso. Linn. Sp. pl. 1124. Jacqu. vind. p. 141.

Hieracium caule unifloro, foliis lingulatis, hirsutis, calyce hirsuto. H. p. helv. n. 49.

Hieracium alpinum scapo unifloro foliis lingulatis mollibus; calyce piloso. Scopol. fl. Carn. ed. 2. p. 106. n. 967.

Hieracium villosum alpinum, flore magnifungulari. Raj. angl. 3. p. 169. t. 6. f. 1.

Icon. Jacqu. fl. Austr. t. 191. (sub *H. alpestre*) Allion. fl. Pedem. n. 771. t. 14. f. 2.

Germ. Alpen-Habichtskraut.

Gall. Eperviere des Alpes.

Hab. In Brisgovix monte altissimo Feldberrarius.

Flor. Junio, Julio. Semina Augusto matura.

Radix subfusiformis, sæpe præmorsa, fibrosa, extus fusca, intus albida. *Caulis* digitaliter pedalis, erectus sæpissime simplicissimus subaphyllus, uniflorus, quandoque uno altero folio instructus, tunc ex axilla, ramo florifero auctus, teres, striatus, inferne glabriusculus superne villosus, sæpe totus plus minusve villosus. *Folia radicalia* plura, oblonga, obtusa vel acuta, sæpe lanceolata, versus basin præ-

inequaliter plus minusve denticulata, quaque integra, glabriuscula, aut sæpissime plus minusve hirsuta. *Calycinae* squamæ erectæ, liri-acutæ, virides, longe pilosæ, et id præcipue in individuis quorum folia longius hirsuta sunt; in aliis autem quorum folia glabriuscula, ex breviori hirsutie, imbutus. *Corolla* mecris lutea. *Pappus* sessilis, capillaris. *Receptaculum nudum.*

Planta pro diverso solo natali alpestri et alpino altiori, rupestri, et gramineo, maxime polymorpha. Possideo specimina a me in Hispaniæ, Galliæ, Sabaudia et Helvetiæ alpinis lecta, aliaque ab amicis ex alpihus Austriæ et Lapponiæ communicata, quæ separatim in herbario sicco servatæ, fere tres ad quatuor species mentiuntur, collatæ multis variationibus intermediis, hinc ita committuntur ut dubitem an tuto vix separarentur. Hinc sine dubio *Hieracium alpestre* Host. Synops. 429. Jacqu. fl. Austr. t. 191. porro *Hieracium pumilum* et *glabratum* Willden. Linn. Sp. pl. p. 1562. vix ab *H. alpino* separandum.

85. *HIERACIUM Pilosella* foliis integerrimis ovatis, subtus tomentosis, caule repente, scapo unifloro. Linn. Sp. pl. 2. p. 1125. Pollich. fl. pal. n. 740.

Hieracium caule unifloro, stolonibus reptantibus, foliis petiolatis, ovatis, longe pilosis, subtus tomentosis. Hall. stirp. helv. n. 55.

Pilosella major repens hirsuta. Bauh. pin. 262.

Pilosella majore flore s. vulgaris repens. Bauh. hist. 2. p. 1039.

Pilosella L. *Tabernæmont.* p. 506. Icon.

308 SYNGENESIA. POLYG. ÆQUA

Icon. Curtis. fl. Lond. t. 279. — Oed. Dan. t. 1110. — Bulliard. herb. t. 2. Blackw. t. 365.

Germ. Gemeines Mausöhrlein.

Gall. Epervière Piloselle. La Piloselle.

Hab. Utrunque in pascuis sylvisque sab ericetis, juniperitis, ad sylvarum et v marginalia passim frequens.

Flor. Majo, Junio, Julio. *Semina* Julii gusto matura. ♀

Radix fibrosa. *Scapus* palmaris, spitha erectus, teres, nudus, inferne pilosus, si hirsutus, uniflorus. *Folia* radicalia pet ovali-oblonga, integerrima, superne læ dia, scabriuscula, inferne alba, tomentosa, que pilis raris longis obsita. *Stolones* rep e centro foliorum, foliosi, hirsuti, *foli* lanceolatis, obtusis pilosis instructi. *Flos* nalis, pallide luteus. *Flosculi* extimi subti purascentes. *Calycinae* squamae lineares quales, hirsutæ, pilosæ, per lentem glol *Receptaculum* setis brevibus conformib structum. *Semen* striatum, nigricans. *P* sessilis, capillaris.

Variat. Scapis interdum 2-3 floris.

Pharm. PILOSELLÆ s. AURICULÆ MURI ba; Flores.

Planta amariuscula, adstringens hodie in m fere exoleta.

Flores apibus melliferi.

In sabulosis arenam figit.

†† *Scapo nudo multifloro.*

1186. *HIERACTIUM dubium* foliis gris ovato-oblongis, stolonibus re

INGENESIA. POLYG. ÆQUALIS. 309

bus, scapo nudo multifloro. *Linn. Sp. pl.*
2. p. 1125. — *Pollich. fl. pal. n.* 741.
omnino!

lieracium caule nudo paucifloro, stoloni-
bus reptantibus, foliis longe pilosis lan-
ceolatis. *Hall. stirp. helv. n.* 53.

ilosella major, repens, minus hirsuta. *Bauh.*
pin. 262.

ilosella major II. *Tabernæm. p.* 507. *Icon.*
n. Oeder. fl. Dan. t. 1111. (*Individuum*
iusto majus),

m. Zweifelhafte Mausöhrlein.

L. Epervière rampante.

. Utrunque in pratis humidis pascuisque
paludosis, udis cum *Melica coerulea* passim
frequens.

. Junio Julio, Augusto. *Semina* Augusto,
Septembri matura. 7

Scapus digitalis, palmaris, quandoque sed
s pedalis, erectus, simplex, teres, glaber,
us, superne villosiusculus. *Folia* radicalia
ma, obverse lanceolata obtusa, rarius acu-
ula, in petiolum abeuntia, integra, basi
ertim et ad marginem pilosa, utrinque læte
ia, glabriuscula, aut pilis raris, albis, mu-
Stolones repentes, e basi caulis, foliosi.
es terminales 2-3-4-5 breviter pedicellati,
de sulphurei. *Calycis* subcylindrici squamæ
tri-lanceolatæ, nigris pilis per longitudinem
itæ. *Receptaculum* setis oblitum.

Distinguitur a præcedente specie, Caule glabrius-
versus calycem squamis 2-3 obfesso, apice 2-3
2-3-4-5 floro, Foliis radicalibus semper lon-
bus. angustioribus, obverse lanceolatis, obtusis,
que viridibus, Floribus utrinque pallide luteis,
ribus.

Flores apibus mel præbent.

§10 SYNGENESIA. POLYG. ÆQUALI

1187, *HIERACIUM Auricula* foliis
 serrimis lanceolatis, scapo nudo
 floro, stolonibus reptantibus. *Lin.*
pl. 2. p. 1126. — Pollich. fl. pal. n.
ommino!

Hieracium caule nudo, stolonibus rep-
 bus, foliis longe pilosis lanceolatis
 ribus umbellatis. *Hall. stirp. helv.*

Pilosella major erecta altera, *Bayh. pin.*

Icon. Oeder. fl. Dan. t. 1044.

Germ. Großes Mausohrlein,

Gall. Epervière Auricule.

Hab. Utrinque in collibus pascuisque
 graminosis, nec non in vinearum
 talibus ad vias et muros passim.

Flor. Junio, Julio. *Semina* Julio, Augusti
 tura. ☞

Differt a praecedente specie, cum qua
 confunditur, *Culmo* semper multo altiori
 trapedali, sesquipedali, piloso, crassiori
 medium vel infra medium folio instructo,
 longioribus, lanceolatis, acutis, (nec ob-
 rigide pilosis, margine integerrimis, *Flc.*
 4-5-9, cymoso-umbellatis, pedicellatis,
 minoribus, magis confertis, saturatius luteis
 pro loco natali sicco aprico.

Flores apibus melliferi.

Nota. *HIERACIUM florentinum.* *Allic.*
pedem. n. 775. seu *Hieracium* caule bra-
 paucifloro, foliis lanceolatis longe pilosis,
 ribus glabris. *Hall. stirp. helv. n. 54.*
Hallerus hanc plantam, auctoritate *uter-*
chenalii, ad *Rhenum Basileæ*, ipse autem
echenalius Basileæ eandem nunquam, se

que in arenosis ad Ticinum per totam Helvetiam transalpinam se legisse affirmat. Act. nov. helvet. I. p. 304.

38. HIERACIUM cymosum foliis lanceolatis integris pilosis, scapo subnudo, basi piloso, floribus subumbellatis. *Linn. Sp. pl. 2. p. 1126. Pollich. fl. pal. n. 743.*

Hieracium caule subnudo, foliis hirsutis lanceolatis, floribus umbellatis, petiolis ramosis. *Hall. stirp. helv. n. 51.*

Hieracium Pilosellæ folio, erectum majus. *Mapp. fl. Als. p. 145.*

Hieracium murorum angustifolium non sinuatum. *Bauh. pin. 129. Prodr. p. 67. Icon.*

Pilosella major altera III. *Tabernæmont. p. 507. Icon.*

Pilosella minori flore hirsutior et elatior non repens. *Bauh. hist. 2. p. 1040.*

n. Oeder. fl. Dan. t. 810.

rm. Vielblumiges aufrechtes Habichtskraut.

ll. Epervière a Bouquet.

b. Utrunque passim in muris præsertim vetustis frequens, nec non in rupibus et locis graminosis elevatis. In muris prope Durlach, Ettlingen, Weingarten, Pforzheim, Heidelberg, Freyburg Brisgovix et alibi.

or. Junio, Julio. Semina Julio, Augusto matura. 7

Radix teres, præmorsa, albicans, fibrosa. *ulis* sesquipedalis, cubitalis, et altior, ereus, nudiusculus, inferne et versus medium pius uno alterove folio præditus, teres, striatus, tuberculis purpurascentibus piliferis asper. *tolones* nulli. *Folia* radicalia plurima, caulina

313 SYNGENESIA. POLYG. EQUASTRI

sub duo, tria, quatuor, alterna, sessilia, omnia erecta, lineari-lanceolata, acutiuscula quandoque obtusiuscula, integra, caulinis longiora, utrinque tuberculis rarioribus rubris pilis obsessa. *Flores* pallide lutei, parvi, caule terminantes, cymoso-umbellati. *Pedunculi* visi, apice uniflori, asperi, pilosi, basi *Bract.* lanceolatis, brevibus, suffulti. *Calycinae* squamae erectae, lineari-acutae, dorso pilis nigrescentibus obsessae.

Flores apibus melliferi.

189. HIERACIUM praemorsum foliis ovatis subdentatis, scapo nudo radice nuda floribus superioribus primoribus. *Lin. Sp. pl.* 2. p. 1126. — *Pollich. fl. palat.* 744.

Hieracium caule nudo, foliis petiolatis ovatis integerrimis hirsutis, floribus spicatis. *Hall. stirp. helv.* n. 51.

Hieracium latifolium praemorsa radice, caulis singulari, Pilosellae majori affini. *Bauh. hist.* 2. p. 1033.

Icon. Oeder. fl. Dan. t. 942.

Germ. Abgeissenes Habichtskraut.

Hab. In nemoribus montosis et pratis elatioribus passim. In Margraviatu superiore ad *Kaiserstuhl* Brisgoviae in collibus humidi graminosis et sylvis, nec non ad viarum excavatarum marginalia virgultosa gramine frequens. Retro *Oberweiler* in sylvaticis nec non prope *Crenzach*. In ditione Basileensi in pratis inter villas *Gundeldingen* in monte *Muteto*, nec non in pratis circum *Münchenstein*. In Principatu superiore in pratis elevatis sylvaticis prope *Mörsburg*.

SYNGENESIA. POLYG. ÆQUALIS. 313

et alibi passim. In Palatinatu circa *Niedengelheim* in pascuis et collibus gramineis siccis t. *Pollichio*.

107. Majo, Junio. *Semina* Julio matura. 7

Radix subcylindrica, præmorfa, fusca, fissula. *Scapus* pedalis, cubitalis, erectus, crassusculus, simplex, aphyllus, obsolete angulatus, pubescens, scabriusculus, læte viridis, ad apicem fusco-rubellus. *Folia* radicalia plurima, ligulata, ovalia, plus minusve elongata, integruscula, aut levissime repando-sinuato-dentulata, utrinque molliter subpubescentia, lætidia. *Flores* parvi, lutei, scapum terminantes, racemosocymosi. *Pedunculi* inferiores in cæmo subtrifidi, superiores simpliciusculi, univari, basi *Bractea* lanceolata suffulti. *Pappus* pillaris.

Flores apibus melliferi,

190. *HIERACIUM staticefolium* caule subnudo ramoso subtrifloro glabro erecto, foliis radicalibus linearibus, obtusiusculis, basi angustatis, rariter dentatis glabris.

Hieracium staticefolium Allion. fl. Pedem. n. 782.

Hieracium staticefolium caule subnudo, foliis ligulatis obtulis, flore sulphureo. *Villars. fl. Delph.* 3. p. 116. t. 27,

Hieracium staticefolium caule ramoso subnudo, foliis linearibus obtusiusculis, floribus majusculis exsiccatione virescentibus. *Lamark. Encycl.* 2. p. 360.

Hieracium caule nudo brachiato paucifloro, foliis linearibus rariter dentatis. *Hall. stirp. helv.* n. 48.

814 SYNGENESIA. POLYG. ÆQUALIS.

Hieracium caule ramoso, foliis radicalibus linearibus subdenticulatis, caulinis minimis. *Gerard. flor. Gallopr. p. 167. n.*

Chondrilla folio non dissecto, caule nudo. *Bauh. hist. 2. p. 1041.*

Icon. Allion. fl. Pedem. t. 81. f. 2.

Germ. Graselkenblättriges Habichtskraut.

Gall. Epervière a févilles de Statice.

Hab. In monte Muteto et Wasserfallen, cum Mimliswyl et Gilgenberg t. Lachenalio.

Flor. Junio, Julio. Semina Augusto matura.

Radix cylindrica, extus fusca, intus alba subfibrosa. *Caulis* pedalis vix altior, erectus, teres, glaber, nudiusculus, ultra medium ramum uno alterove unifloro præditus, superne subcrassatus. *Squamula* una alterave lineari præditus, sub rami insertionem, folio lineari acuto angustius integro munitus. *Folia* radicalia numerosa, subsessilia, lineari-lanceolata, basi angustata, rariter dentata, apice obtusiuscula, glabra pallide viridia. *Flos* majusculus, luteus, ficcitate virescens. *Calycinae* squamæ erectæ, linearis-acutæ, subvillosæ-tomentosæ. *Pappus* capillaris, sessilis, longitudine calycis.

1191. Hieracium aurantiacum foliis integris, caule subnudo simplicissimo piloso corymbifero. *Linn. Sp. pl. 2. p. 1126*

Hieracium caule subnudo, foliis ovatis integris, floribus umbellatis aurantiis. *Hall stirp. helv. n. 50.*

Hieracium hortense non laciniatum floribus atropurpureiscentibus. *Bauh. pin. 131 Prodr. p. 65.*

SYNGENESIA. POLYG. ÆQUALIS. 315

non. Jacqu. fl. Austr. t. 410. — Allion. fl. Pedem. t. 14. f. 1,

erm. Pomeranzenfarbiges Habichtskraut.

all. Epervière a fleurs d'orange.

ab. In Alfatiaë montis Ballon summis pascuis.

lor. Junio, Julio. Semina Augusto maturat. 7

Caulis pedalis, sesquipedalis, erectus, subhyllus, simplex, quandoque inferne folio uno trove, et in medio squama foliacea præditi, totus fusco-hirsutus. *Folia* radicalia ovali-lanceolata, subobliqua, integriusculum versus basin angustata, sæpe obsolete rari-dentata, utrinque hirsuta. *Flores* terminati, cymoso-umbellati, 7-11, amoene purpureo-aurantiaci. *Pedunculi* hirsuti, squamulis lim muniti. *Calycinæ* squamæ hirsutæ. *Papillæ* capillaris.

Pulchra omnino ob flores amoenos planta, passim hortis colitur, ubi non raro caule superne ramo uno trove aucto observatur.

Flores apibus melliferi.

††† *Caule folioso.*

192. HIERACIUM scorzoneraefoli-

um foliis lineari-lanceolatis glaucis et pilosis, caule oblique piloso et incano paucifloro, calyce fetuloso, farinoso.

Hieracium scorzoneraefolium. Villars. fl. Delph. 3. p. 111.

Hieracium glaucum Allion. fl. Pedemont n. 781.

Hieracium foliis lanceolatis, glaucis, caule brachiato multifloro. *Hall. Emend. II. p. 73.*

316 SYNGENESIA. POLYG. ÆQUALIS

Hieracium Nro. 48. var. *β*. *Hall. stirp. ha*
p. 21.

Hieracium folio Statices, caule foliato. *J*
guier. pl. Veron. Suppl. p. 270.

Hieracium Tragopogonis folio. *Bauh. pi*
129.

Chondrilla folio non dissecto, caule foliat
Bauh. hist. 2. p. 1041. Icon.

Icon. Allion. fl. Pedem. t. 81. f. 1.

Germ. Scorzonerenblättriges Habichtskraut
Gall. Epervière a fevilles de Scorzonere.

Hab. In Ditione Basileensi in rupibus sup
Dornach et Ramstein, nec non in mont
Muteto et Wasserfallen t. Lachenalio.

Flor. Junio, Julio. *Semina* Augusto matura.


Primo intuitu affine *H. staticefolio*, a quo
distinguitur, *Caule* sæpissime altiore sesqui-
dali, cubitali, subsolioso et ramoso, multifloro
rarius subnudo, unifloro, *Foliis* radicalibus
longe lanceolatis, acutissimis, integris et ob-
ter dentatis, glaucis, glabris, præter longiu-
culos pilos, quibus in juniore planta ad exortum
in dorso et marginibus instructa sunt, *caulin*
folia radicalibus similia, triplo semper minor
Floribus subminoribus, flavis, siccatibus non v-
ridescentibus. *Calycinae* squamæ setulosæ, bal
maxime farinaceæ.

Confer. Nov. Act. Helvet. I. p. 303.

Obs. Individua in rupestribus calcareis alpinis helveti-
cis et pyrenaicis observata, nostratibus similia

HIERACIUM porrifolium caule ramoso fo-
lioso, foliis lineari-lanceolatis elongatis, pers-
gustis pallide virentibus, glabris. remote obsolet
denticulatis, est planta mere alpina, que vix in re-
gionibus nostris crescit, quam vidi in rupestribus





Tab.



Tom. III.

YESIA. POLYG. ÆQUALIS. 317

alpium Abbatiszellanensium auf dem
Sentis et Hohenkasten. Distinctif-
ino species. vid. Jacqu. fl. Austr.

ERACIUM *bupleuroides* foliis
tri-lanceolatis, caulinis radicalibus
tribus, glaucis, integris, rariter pilo-
sis, caule erecto angulato glabro mul-
to, calyce piloso.

im *scorzoncræfolium* β. involucre
to. *Lamarck et Decand. flor. franç.*
26. — *Villars. fl. Delph. 3. p. 112.*
n?

oria.

Isenöhrchen - blättriges Habichts-

errière bupleuroide.

isdem cum præcedente regionibus,
tempore florens. 7

præcedenti speciei, a qua autem
guitur, *Caule* semper magis folioso,
Foliis radicalibus caulinisque infi-
riorem petiolum membranaceum abeun-
do fere longioribus, quinque ad sex
gis; caulinis reliquis semper duplo
us, 8-10, triplo longioribus, 3-4
gis, omnibus cæterum acuminatis,
terminatis, latiusculis, semiunciam fere
que glaucis, rariter pilosulis, inte-
quandoque rariter et vix manifeste
s. *Pedunculis* tenuioribus, unifloris,
incrassatis, pilosis. *Calycibus* longe
pilosis. *Floribus* subminoribus.

288 SYNGENESIA. POLYG. EQUALI

Nota. Planta omnino ut facile ex characteribus
tis, et habitu patet, specie diversa a *H. scorzoneraefolio*.

H. bupleuroides ob folia *Bupleurologa*, denominavi.

1194. *HIERACIUM saxatile* caule
ramoso folioso, foliis radicalibus ang
lanceolatis, utrinque acuminatis, rar
denticulatis, basi pilosis, caulinis mi
ribus angustissimis.

Hieracium saxatile caule ramoso folio
foliis anguste lanceolatis utrinque acu
natis, basi pilosis. *Jacqu. Collect.* 1
159. — *Willd. Linn. Sp. pl.* 3. p. 159

Icon. Jacqu. Icon. rar. 1. t. 163.

Germ. Stein-Habichtskraut.

Gall. Epervière des roches.

Hab. In Ditione Basileensi in rupibus calca
supra *Dornach*, *Ramstein*, *Gilgenberg*
Mimliswyl et *Wasserfallen*.

Flor. Junio, Julio. *Semina* Augusto matur

Affine primo intuitu *H. scorzoneraefolio*
a quo autem differt, *Caule* demissiore, pet
Foliis radicalibus manifestius, denticulatis,
petiolo utrinque longe et molliter pilosis; ca
linis remotioribus, multo angustioribus stric
libus fere setaceis. *Calycinae* squamae line
ranceolatae, pilosae, inferiores breviores. *Ca*
hulae ligulae luteae.

Individua nostra saepe uniflora cum rudiment
morum, simillima *Jacquini* in omnibus re
partibus.

1195. *HIERACIUM humile* foliis infe
ribus petiolatis pinnatifide runcinatis

INGENËSIA. POLYG. ÆQUALIS. 319

futis, superioribus lanceolatis integris, caule ramofo subflexuoso, subbifloro, hirsuto.

Lieracium humile caule paucifloro foliis inferioribus hirsutis et incisís vix altiore. *Hofst. Synopf.* 432.

Lieracium pumilum. *Jacqu. fl. Austr.* p. 53.

Lieracium Jacquini foliis basi pinnatifidis hirsutis, viridibus, pilis calycinis glandulosis, caule subbifloro. *Villars. fl. Delph.* 3. p. 123. t. 28.

Ilosellæ majoris seu *Pulmonariæ luteæ* laciniatæ species minor. *Bauh. hist.* 2. p. 1034. *Icon.*

n. *Jacqu. Flor. Austr.* 2. t. 189.

m. *Niederer Habichtskraut.*

l. *Epervière nain.*

o. In Ditione Basileensi in montis *Dornach* rupestribus; majora pedalia specimina *Bruntruti* ad moenia prope portam *Sancti Germani.* t. *Lachenalio.*

r. Junio, Julio. *Semina* Augusto matura. ¶

Caules ex una radice sæpe plures, digitalemipedales, pedales, assurgentes, subflexi, hirsuti, ramofo. *Rami* alterni, pauci, hirsuti, simpliciusculi. *Folia radicalia,* inaque inferiora petiolata, ovata, ovatoeolata, obtusa vel acuta, runcinato pinnata, aut profunde inæqualiter dentata, hirsuta, viscida, viridia; *suprema* minora, sublanata, integriuscula. *Pedunculi* aphylli, hirsutiflori. *Calycinae* squamæ plurium ordinis, lineares, acuminatæ, per longitudinem hirsutæ. *Flores* circiter 2-3, quandoque

plures, magni, lutei. *Semina* oblonga. *Pappi* capillaris, sessilis.

Nota. Individua nostra minora, austriacis in monte Schneeberg lectis, simillima sunt.

Minime confundendum cum *H. prunellae-folio* Gouani. Allionii et Villarsii, quod Pyrenaeis et Galliae alpinis legi, et quod est *Hieracium*. Hall. stirp. helv. n. 42. f. *Crepis pyramidea* Linn. Sp. pl. 2. p. 1131. f. *Hieracium pumilum* Linn. Mant. 107.

Ad rariores omnino florae cives pertinet, in Germania hucusque nondum observata planta.

1196. *HIERACIUM murorum* caule piloso superne paucifloro, foliis radicalibus ovatis dentatis petiolatis pilosis, caulino subsessili, minori.

Hieracium caule ramofo, foliis radicalibus ovatis dentatis: caulino minori. Linn. Sp. pl. 2. p. 1128. — Pollich. fl. pal. n. 74.

Hieracium caule subnudo paucifloro, foliis ovato-lanceolatis circa petiolum dentatis. Hall. stirp. helv. n. 46.

HYNGENESIA. POLYG. ÆQUALIS. 321.

Caule altiore ramosiore, foliis radicalibus longioribus, longiusque petiolatis, profundius dentato-incis, minus hirsutis, in sylvestribus obveniens.

Hieracium sylvaticum murorum laciniatum, minus pilosum. *Bauh. pin.* 129. *Pollich. fl. pal. n.* 745. β .

Pilosellæ majoris seu **Pulmonariæ** luteæ species magis laciniata. *Bauh. hist.* 2. p. 1034.

Pulmonaria gallica foemina. *Tabernæmont.* p. 504. *Icon bona!*

r m. **Mauer-Habichtskraut.**

all. **Epervière des Murs.**

a. b. a. et **b.** utrinque in muris, rupibus, vinerarum et sylvarum marginalibus frequens
γ. præsertim graminosa sylvatica amat.

or. Majo, Junio, Julio. *Semina* Julio, Augusto matura. γ

Caulis pedalis, fesquipedalis et altior, erectus, sulcato-striatus, hirsutus, subnudus uno crove folio versus medium instructus. **Folia** *lucalia* plurima longe petiolata, ovato-lanceolata acuta aut obtusiuscula, cumprimis versus **lin** plus minusve profunde dentata, utrinque hirsuta, viridia, quandoque maculis fusco-bellis notata. **Caulinum** folium sæpissime unum subpetiolatum, minus, argute dentatum. **ores** caulem terminantes, pedunculati, meoeres, lutei. **Pedunculi** subumbellati, hirsuti, **sciales**, fesquiunciales. uniflori. **Calycinæ** *uamæ* exteriores breviores, interiores lineari-aceolatæ, erectæ, hirsutæ.

In sylvaticis præcipue obveniens minus hirsuta cæterum variationi **a.** simillima, folio caulino unico alterove tantum instructa, minime confundenda cum **H. Lachenalii** nostro.

222 SYNGENESIA. POLYGA. ÆQUALIS.

Nota. Descriptio *H. sylvatici* Roth. fl. germ. non male quadrat cum *H. Lachenalii* Synonyma autem a Cl. Rothio ad suum *H. sylvaticum* citata, non *Lachenalii* planta sed variationi. γ . seu *Hieracio murorum* sylvatico respondent.

Obs. Variatio γ . in horto culta, semper multo longioribus pedunculis quam α . prædita et minus hirsuta est. Folia caulina nunquam unico alterove plura.

1197. *HIERACIUM Lachenalii* caul. erecto piloso folioso superne ramoso foliis radicalibus petiolatis ovato-oblongis, grosse sinuato-dentatis, caulinis alternis petiolatis ovali-lanceolatis, utrinque acutis. dentatis ciliatis supremis sessilibus.

Germ. *Lachenalisches-Habichtskraut.*

Gall. *Epervière de Mr. Lachenal.*

Hab. In Margraviatu superiore, t. *Lachenalio* in sylva *Weilensi*, durante bello interrim excisa.

Flor. Junio, Julio. *Semina* Augusto matura. γ

Caulis 3-4 pedalis, erectus, crassus, pilosus, foliosus, superne saltem ramosus. *Folia radicalia* plurima, longe petiolata, ovali-oblonga, petiolo marginato longiusculis pilis villosa, magna, subhirsuta, grosse sinuato-dentata, *Caulina* inferiora et media petiolata, alterna, radicalibus similia, potius latiora, ultra binas fere uncias lata, quatuor ad quinque uncias longa, acuta, *suprema* sessilia. *Flores* in ramosis pedunculis quasi paniculati, terminantes, mediocres, lutei. *Calyces* villosi-hirsuti.

Obs. Planta omnino maxime diversa a *H. murorum* sylvatico. a Cl. *Lachenalio* in Act. nov.

SYNGENESIA. POLYG. ÆQUALIS. 323

Helvet. I. p. 298-299 descripta, ab amicissimo viro mihi communicata, in horto culta, mutationi non succubuit.

Minime confundenda cum *H. sylvatico* Willden. Linn. Sp. pl. 3. p. 1578, quod ut ex Synonymis a Clarissimo viro allegatis, tria specie diversa *Hieracia* comprehendit, scilicet variationem nostram *H. Hieracii murorum* ut facile ex Barrelier Synonymo apparet, et *H. murorum Allionii* fl. Pedem. n. 785. t. 28. f. 1. quod omnino peculiarem speciem fuit; porro tertiam: *Pilosellæ majoris* s. *Pulmonariæ* species angustifolia Bauh. hist. 2. p. 1034, *Pulmonaria* dictum angustifolium Raj. Synops. 168. Petiv. Brit. 3. 13. f. 5. quæ speciem sequentem nostram comprehendit.

1198. *HIERACIUM angustifolium* caule erecto piloso folioso superne ramoso, foliis radicalibus caulisque inferioribus petiolatis lanceolatis, superioribus amplexicaulibus sessilibus ovali-lanceolatis sinuato-inequaliter dentatis hirsutis.

Hieracium murorum, laciniatum minus pilosum folio angustiore. Bauh. pin. 129. Mapp. fl. Als. p. 144.

Pilosellæ majoris s. *Pulmonariæ luteæ* species angustifolia. Bauh. hist. 2. p. 1034.

Pulmonaria gallica tenuifolia. Tabernaemont. p. 505. Icon bona. Dillen. Cat. pl. Gieff. p. 95. nov. pl. Spec. p. 48.

Germ. Schmalblättriges Habichtskraut.

Gall. Epervière à févilles étroites.

Hab. In Margraviatus superioris passim sylvestribus. In Brisgovia am Kaiserstuhl prope Burgheim, Sponek, ad arcem Limburg et alibi in collibus virgultosis sylvestribus. In ditionis Basileensis sylvestribus passim.

324 SYNGENESIA. POLYG. EQU

nec non in pratis udis circa *Newbachenalio*. In Alsatie muris vetustis *Rappolstein* t. *Mappo*.

Flor. Junio, Julio. *Semina* Augusto ma

Caulis pedalis, sesquipedalis, rectiosus, piloso-hirsutus, superne ramosus *radicalia* petiolata, lanceolata, utrinque minata, tres ad quatuor uncias longa, ficiam lata, margine utrinque inaequalitata, hirsuta; *Caulina* alterna, inferiora lata, radicalibus similia, suprema sessilia, lanceolata, denticulata, minora, angustiores in ramis axillaribus, erectis, hirsutifloris, lutei, mediocres. *Calyces* hirsutius capillaris.

Nota. Planta omnino diversa a precedentiibus, optime in *Tabernaemontana* lineata.

Obs. In Pyrenæorum et Helvetiæ alpinis magis stricta, hirsuta, foliis angustioribus prædita. In horto Botanico culta, mutationi non sub

1199. *HIERACIUM paludosum*

paniculato, foliis amplexicaulibus tatis glabris, calycibus hispidis.

Sp.pl. 2. p. 1129. — Pollich. fl. pal.

Hieracium foliis glabris dentatis, imis latis, superioribus amplexicaulibus licibus hirsutis. *Hall. stirp. helv. n.*

Hieracium montanum latifolium glabrum. *Bauh. pin. 129. Mapp. fl. Als.*

Hieracium montanum majus latifolium. *bernæmont. p. 494. Icon bona.*

β. *Hieracium montanum latifolium glabrum minus.* *Bauh. pin. 129. Mapp. fl. Als.*

SYNGENESIA. POLYG. ÆQUALIS. 325

Hieracium latifolium glabrum ex valle Griesbachiana. *Bauh. hist. 2. p. 1033.*

Hieracium latifolium glabrum minus. *Tabernæm. p. 495. Icon bona.*

non. Schkuhr. bot. handb. t. 221.

erm. Sumpf-Habichtskraut.

211. Epervière marécageuse.

2 b. In pratis montosis udis sylvaticis passim frequens. Auf dem Tohel, Kaltenbrunn, Herrnwiese, in convallibus prope Bühl, Oppenau, Griesbach, Antegast, Petersthal, am Kniebis. In Margraviatu superiore in pratis udis prope Schweighof, Sirnitz, Neuenweg, cum var. β. non infrequens. In Vogesi Alsatiae montosis sylvestribus pratisque retro Barr et alibi, In Palatinatu circa Lautern in pratis paludosis.

For. Junio, Julio. Semina Augusto matura. 7

Caulis sesquipedalis, cubitalis, et altior, erectus, sulcatus, glaber, superne ramosus. Semi alterni, aphylli, glabriusculi, superne divisi. Folia alterna, inferiora in pedunculum lam decurrentia, ovali-lanceolata, acuminata, sin versus præsertim profunde sinuato-dentata, superiora amplexicaulia, cordato lanceolata, acuminata, sinuato-dentata, omnia glabra, læte viridia. Flores caulem ramosque terminantes, mediocres, lutei. Calycinæ squamæ erectæ, dorso pilis nigris hispidae.

Planta bonum pecoribus pabulum præbet; indicium facie Turfæ subjacentis.

bf. Planta in horto culta, mutationi non succubuit.

200. HIERACIUM blattarioides foliis lanceolatis amplexicaulibus dentatis, flo-

326 SYNGENESIA. POLYG. ÆQUALIS

ribus solitariis, calycibus laxis. *Lin. Sp. pl. 2. p. 1129. Lamarck. Encycl. p. 368.*

Hieracium foliis amplexicaulibus ferratis glabris, hamis aristatis, calycibus villosis. *Hall. stirp. helv. n. 39.*

Crepis austriaca foliis oblongis denticulatis amplexicaulibus, involucro laxissimo calycibus hispidis. *Jacqu. Enum. Stirp. Vind. p. 140. et 270.*

Icon. *Jacqu. fl. Austr. V, t. 41.*

Ger m. *Blattariaförmiges Habichtskraut.*

Gall. *Epervière fausse-blattaire.*

Hab. In monte *Wasserfallen* t. *C. Bauhino* *Laohenalio.*

Flor. Junio, Julio. **Semina** Augusto matura.

Radix lignosa, fibris aucta, fusca, quandoque præmorsa. **Caulis** pedalis, sesquipedalis erectus, striatus, inferne rubellus, sæpe solitarius, foliosus, hirsutus, uniflorus, aut superius ramosus. **Rami** 1-3 alterni, uniflori. **Folia radicalia** nonnulla, facile in planta florente evanescentia, longe petiolata, oblonga, denticulata; **Caulina** alterna, sessilia, subamplexicaulibus, oblongo-lanceolata, denticulata, subhirsuta, superiora acutiora. **Flos** terminalis majusculus, luteus. **Calycinae** squamæ oblongæ, auctæ, exteriores laxæ, patentes, ab interioribus erectis distantes. **Pappus** capillaris, sessilis.

Nota. Planta nostra, omnino in omnibus partibus cum illa a *Jacquino* depicta, convenit.

1201. **HIERACIUM Succisaefolium**

foliis radicalibus longe petiolatis, caulinis semiamplexicaulibus ovali-oblon-

gis glabriusculis subintegris obtusis, caule erecto scabriusculo.

Hieracium Succisaefolium. Allion. *fl. ped.* n. 786. — Lamarck et Decandolle. *fl. franc.* 4. p. 28.

Hieracium integrifolium. Willden. *Linn. Sp. pl.* 3. p. 1568. Hoppe *Cent.*

Hieracium foliis lævibus integerrimis, radicalibus longe petiolatis, ellipticis, caulinis amplexicaulibus, hamis obtusis. *Hall. stirp. helv.* n. 47.

Germ. Habichtskraut mit Abbissblättern.

Gall. Epervière a fevilles de Succise.

Hab. In ditione Basileensi in monte Vogelberg, et aliis montis *Juræ* tractibus. t. *Lachenalis.*

Flor. Junio, Julio. *Semina* Augusto matura. ¶

Radix præmorsa, fusco-nigricans. *Caulis* sesquipedalis, cubitalis, erectus, foliosus, subhirsutus, striatus, superne ramo uno alterove erecto, hirsuto, unifloro munitus. *Folia radicalia*, caulinaque infima in petiolum decurrentia, ovalia, obtusa, integra, basi angustata, superiora sessilia, subhirsuta, vel glabra, laete viridia. *Corolla* saturate lutea, mediocris. *Calycinae squamae* lineari-acutae, in unica serie dispositae, erectae, obscure virescentes, hirsutae. *Pappus* capillaris, sessilis.

Nota. Maxime affine *H. molli* Jacqu. *fl. Austr.* 2. p. 12. t. 119, a quo mere differre videtur, foliis non mollibus, neque villosis, sed vel leniter subhirsutis vel penitus glabris. Conf. *Nov. Act. Helv.* 1. p. 300.

1202. *HIERACIUM amplexicaule* foliis amplexicaulibus cordatis subdentatis, pe-

328 SYNGENESIA. POLYG. ÆQUALIS.

dunculis unifloris (2-3-4 floris) hirsutis
caule ramoso. *Linn. Sp. pl.* 2. p. 1129.

Hieracium foliis ovato-lanceolatis, raris
dentatis, caulinis amplexicaulibus. *H. Helv.*
stirp. helv. n. 36.

Hieracium amplexicaule foliis radicalibus
ovato-lanceolatis, acutis, basi dentatis,
caulinis hastato-cordatis, amplexicaulibus.
Gouan. Illustr. bot. p. 58. n. 6.

Hieracium pyrenaicum longifolium amplexicaule.
Tournef. inst. p. 472.

B. Foliis omnibus brevioribus rotundioribus
subdentatis.

Hieracium pulmonarioides foliis ovatis
subdentatis, hirsutis, pedunculis proli-
feris calycibus glandulosis. *Villars. fl. Delph.*
2. p. 133. t. 34.

Hieracium pyrenaicum rotundifolium amplexicaule.
Tournef. inst. p. 472.

Icon. *Allion. fl. Pedem.* Tab. 15. f. 1. t. 30.
f. 2. (optima).

Germ. Stengelumfassendes Habichtskraut.

Gall. Epervière embrassante.

Hab. In ditione Basileensi, in montis *Dornach*,
Wallenburg, *Schauenburg* rupibus, nec
non in monte *Wasserfallen* ad novem font-
es t. *Lachenalio*.

Flor. Junio, Julio. Augusto Semina matura. 7

Radix magna, dura, fibris capillata, fusca.
Caulis cubitalis et altior, erectus, crassiuscu-
lus, sulcato-striatus, piloso-hispidus, ramosus.
Rami alterni, hirsuti. **Folia radicalia** in petio-
lum decurrentia, ovali-lato-lanceolata, basi
versus angustiora, acuta, ultra binas uncias lata,
8-9 uncias longa, margine argute inæqualiter

SYNGENESIA. POLYG. ÆQUALIS. 329

ovato-denticulata, venosa, villis flavescentibus utrinque et præsertim in costâ et petiolo arbata; *Caulina* alterna, sessilia, amplexicaula, cordato-ovata, acuta, argute denticulata, præmura integriuscula, omnia pilosulo-hirsuta, et viridi flavescentia. *Flores* caulem ramosque, terminantes, majusculi, lutei. *Calycinæ* squa-
 æ laxae, erectæ, lineari-lanceolatæ, pilis
 obiferis cum pedunculis, *Bracteolisque* hirti. *Pappus* capillaris.

b f. Individua in pyrenæis abs me lecta, nostratibus simillima sunt.

Planta junior glutinosa, grate odorata, balsamea.

In horto botanico ex seminibus culta, mutationi succubuit.

203. HIERACIUM villosum caule erecto subramoso, foliisque pilosis, radicalibus lanceolatis obsolete denticulatis integriusculis, caulinis semiamplexicaulibus ovato lanceolatis integriusculis, calycibus albide pilosis.

Hieracium villosum caule ramoso folioso, foliis hirtis, radicalibus lanceolato-ovatis, dentatis; caulinis amplexicaulibus cordatis. *Linn. Sp. pl.* 2. p. 1130.

Hieracium foliis villosis integerrimis, radicalibus ligulatis, caulinis ovatis, amplexicaulibus calycibus villosis. *Hall. stirp. helv. n.* 44.

Hieracium alpinum latifolium villosum magno flore. *Bauh. pin.* 128. — *Mapp. fl. Alf.* p. 145.

Hieracium montanum latifolium. *Tabern. n. XII. p.* 494. *Icon mala.*

330 SYNGENESIA. POLYG. ÆQU.

Icon. Jacqu. fl. Austr. t. 87.

Germ. Zottiges Habichtskraut.

Gall. Epervière vélue.

Hab. In Sylvæ nigræ montosis propet
bach, Petersthal et Antegast; et in
Vogefis t. *Tabernaemontano*; nec
Alsatia retro *Niedek* versus *Dach-*
Mappo.

Flor. Junio, Julio. *Semina* Augusto m

Radix fusco nigricans, dura. *Cauli*
ris, vix pedalis, erectus simpliciusculi
florus, aut ramo sæpe uno alterove p
teres, pilosus. *Folia radicalia* plura, c
basi attenuata, acuta, integra, aut sal
solete rariter denticulata, glauca, lon
mollibus præsertim in costa et margine
caulina alterna, subovata, suprema sub
ovata, amplexicaulia. *Flos* præmagnus,
Calyx longe albide-piloso-barbatus.

Obs. Planta in alpinis omnino maxime variab
in pyrenæorum jugis altioribus individu
cissima, uniflora, caule aphylo aut salte
alterove folio instructa; alia, caule folio
floro; alia superne ramoso; Ramis 1-2-3
ris prædita, plus minusve pilosa. Ad pri
vationem pertinet H. Schraderi. Læ
et Decandolle. *Flor. franc. 4. p. 2*

1204. *HIERACIUM sabaudum*

erecto multifloro, foliis ovato-
latis dentatis semiamplexicaulibu.
Sp. pl. 2. p. 1131. — Pollich. fl.
747.

Hieracium foliis hirsutis dentatis, in
bus ellipticis, superioribus ovato-
latis. *Hall. stirp. helv. n. 35.*

SYNGENESIA, POLYG. ÆQUALIS. 331

Hieracium caule multifloro folioso; foliis lanceolatis dentatis; dentibus glandulosis. *Scop. fl. Carn. 2. n. 972.*

Hieracium fruticosum latifolium hirsutum. *Bauh. pin. 129.*

Hieracii sabaudi varietas I. *Bauh. hist. 2. p. 1030.*

Ramosior, glabra, annosior, adulta.

Hieracium montanum latifolium, foliis dentatis glabrum. *Mapp. fl. Als. p. 145.*

o n. *Oeder. fl. Dan. t. 872.*

r m. *Savoier-Habichtskraut.*

ll. *Epervière de Savoie.*

b. Utrisque in dumetis, sylvarum marginibus, ad praecipitia virgultosa, dumetosa aspera passim frequens.

or. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ζ

Caulis 1-2-3-4 pedalis et altior, erectus, terne hirsutus, glanduloso-rigidus, superne natus, valde foliosus, durus. **Folia** alterna, inferiora sessilia, lanceolata; superiora sessilia, semiamplexicaulia, ovato-lanceolata, sinuato-dentata, duriuscula; **summa** lanceolata, integra, minora, omnia scabriuscula, villosula, quandoque glabriuscula, maculis fusco-rubellis depicta. **Flores** caulem ramosque terminantes, dioecres, flavi. **Pedunculi** incani, magis minusve tomentosuli. **Calycinae** squamae lanceolatae, acutae, exteriores breviores, apice patentes, scabriusculae, aut glabrae. **Receptaculum** albidum subsetosum, **Pappus** capillaris, sessilis.

Triat Caule magis minusve hirsuto, Foliis magis minusve, interdum vix manifeste dentatis latioribus et angustioribus, hirsutis vel subglabris.

338 SYNGENESIA. POLYG. ÆQUALIS.

Visum excavatarum marginalia condensat. Flores melliferi.

1205. *HIERACIUM umbellatum* foliis

linearibus subdentatis sparsis, floribus subumbellatis. *Linn. Sp. pl. 2. p. 113.*
Pollich. fl. pal. n. 748. Hall. stirp. helv. n. 34.

Hieracium fruticosum angustifolium majus.
Bauh. pin. 129. — Mapp. fl. Alf. p. 145.

Hieracium montanum angustifolium L. *Tabernaemont. p. 495. Icon.*


β. Foliis angustioribus, caule ramosiori, aut minus ramoso.

Hieracium saxatile. Schkuhr. bot. handb. 3. p. 32.

Hieracium fruticosum minus. Mapp. fl. Alf. p. 145.

γ. Foliis pene ovatis, vix dentatis, caule hemili paucifloro.

Icon. Oeder. fl. Dan. t. 680.



SYNGENESIA. POLYG. ÆQUALIS. 339

ice reflexis scariosis. *Receptaculo* florum mag-
profunde alveolato, piloso.

Secundum diversum locum natalem, habitu, mag-
ndine, et foliorum respectu, ut *H. sabaudum*
ltum variat.

of. *Hieracium saxatile Schkuhrii* omnino huc
pertinet, non confundendum cum *H. saxatile*
nostro.

106. HIERACIUM *chondrilloides*

foliis caulinis elongato-dentatis glabris:
radicalibus lanceolatis integris, caule ra-
moso. *Linn. Sp. pl.* 2. p. 1128. — *Jacqu.*
vind. p. 273. t. 7. — *Lamarck. et Decan-*
dolle fl. Franc. 4. p. 36.

Hieracium alpinum pumilum, chondrillae
folio. *Bauh. pin.* 129. — *Prodr.* 64.

on. *Jacqu. fl. Auftr.* 4. t. 429.

er m. *Chondrillenblättriges Habichtskraut.*

ill. *Epervière fausse-chondrille.*

ib. In Alfatizæ Vogesi monte Balon t. *Bu-*
choz.

or. Junio, Julio. 7

Nota. Planta a Bursero in alpibus altioribus
Schneeberg Austriæ inferioris lecta, ab ill.
Jacquino optime descripta et depicta, vix in
subalpinis Alfatizæ crescit. In collectione certe fic-
ca plantarum Alfatizæ superioris Lachenalii
quoad *Hieracia* completissima, omnino deest.
Hinc forsan ex Catalogo plantarum Alfaticarum
eliminanda.

CCCLXIV. C R E P I S.

Linn. Gen. pl. ed. Schreb. n. 1239.

Calyx communis, duplex: exterior brevif-
olus, patulus, deciduus: interior ovatus,

334 SYNGENESIA. POLYG. ÆQUA

simplex, fulcatus, persistens: *squamis* bus conniventibus. *Corolla* compo-
cata, uniformis: *Corollulis* hermaph-
plurimis, æqualibus. *Propria* monopet-
gulata, linearis, truncata, quinque-
dentata, fructiferus subrotundus. *Semina* o-
Pappus capillaris, stipitatus, (in aliis
Receptaculum nudum, in paucis pilosum

1207. *CREPIS foetida* foliis run-
pinnatis hirtis, petiolis dentatis,
stipitato.

Crepis foetida. Linn *Sp. pl.* 2. p. 11
Pollich. fl. pal. n. 749.

Crepis foliis scabris pinnatis, pinnis
lobis dentatis retroverfis, extrema
lobulari maxima. *Hall. stirp. helv.*

Barkhausia foetida. Lamarck et Deca-
flor. Franc. 4. p. 42.

Picris foetida. Lamarck *flor. fr.* 2. p. 1.

Hieracium luteum Cichorii sylvestris
amygdalas amaras olens. *Morif. l.*
p. 63. *Sect.* 7. *t.* 4. *f.* 4.

Hieracium foliis cichorii sylvestris vi-
odore castorei. *Magnol. monsp.* 1
Icone.

Germ. Stinkender Pippau.

Gall. Crépide foetide.

Hab. Utrunque passim ad viarum versuras
ras siccas apricas graminosas. Inter *Car-*
et Mühlberg am Landgraben et ali-
quens, nec non in vinearum margini-
bus inde.

YNGENESIA. POLYG. ÆQUALIS. 335

r. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix fusiformis. *Caulis* pedalis et altior, tus, sulcatus, hirsutus, ramosus. *Folia* *ra-*
lia numerosa, petiolata, profunde pinna-
ta, dentata; *caulina* alterna, amplexicaulia,
ato-pinnatifida: *pinna* extima majori, den-
tata, lateralibus latioribus, infimis sub-
ribus, incanis, piloso-scabris. *Flores* cau-
ramosque terminantes, pedunculis uniflo-
ngiusculis insidentes, ante explicationem
tes, lutei, mediocres. *Calycinae* squamæ
albido-cinerascentes, villosæ, molles.
us longe stipitatus, capillaris. *Receptacu-*
lveolatum.

lanta canescens. Tritæ Odor singularis, bi-
osus, ad castoreum, amygdalas amaras, vel cor-
salicis recentem accedens.

Medicatas vires promittit, hucusque in medicina
cta.

8. *CREPIS Lachenalii* foliis radica-
libus runcinato-pinnatis hirtis pinna
extrema latiori, floribus erectis, calyci-
bus subtomentosis, pappo stipitato.

crepis taurinensis foliis scabris, radicalibus
lyrato-runcinatis, caulinis lanceolatis am-
plexicaulibus basi dentatis, calycibus sub-
tomentosis, squamis exterioribus margine
membranaceis. *Willden. Linn. Sp. pl.* 3.
p. 1595.

crepis taraxacifolia *Thuil. fl. Paris.* 2. p.
409.

crepis tectorum. *Villars. fl. Delph.* 3. p. 144.
excl. *Synon.*

236 SYNGENESIA. POLYG. ÆQUAL

Crepis vesicaria. Balbis. Taur. p. 9.
Synon.

Barkhausia taraxacifolia. Lamarck
candolle. fl. franc. 4. p. 43.

Crepis foliis subscabris, semipinnatis,
extrema maxima. Hall. stirp. helv.

Hieracium asperum maximum Chon
folio. Bauh. prodr. p. 64.

Hieracium Erucaefolio hirsutum. Bau
2. p. 1025. Icon bona.

Germ. Lachenalischer Pippau.

Gall. Crépide de Mr. Lachenal.

Hab. In Marggraviatu superiore in agri
 neis, locisque graminosis siccis ad vi
 ca *Weil, Riechen, Basileam* et alib.
 quens.

Flor. Julio, Augusto. *Semina* Augusto,
 tembris matura. ☉

Praecedenti speciei maxime affinis,
 distinguitur, *Caulibus* saepissime altioribus
 fioribus, minus hirsutis. *Foliorum* extrema
 semper latiori. *Floribus* ante et post expli
 nem semper erectis. Odore nullo.

Differt a *Crepide bienni* Linn. cum qua al
 lero aliisque passim sub variatione citatur, P
 semper stipitato.

A. b. *Lachenalio* denominata, quia primu
 qui eam in Nov. ct. helv. tanquam peculiarem
 ciem distinxit.

1209. *CREPIS setosa* foliis radica
 caulisque infimis runcinatis, lob
 tremo sinuato-dentato, maximo,
 rioribus sessilibus semiamplexicau
 basi hastatis, inciso-sinuato-dentat
 tulosis, caule ramoso, calycibus se
 pappo stipitato.

C

SYNGENESIA. POLYG. ÆQUALIS. 397

Crepis setosa. Hall. fil. in Roem. Arch. 1.
St. 2. p. 1. — Suter. fl. helv. n. 2. p. 155.

Barkhausia setosa. Lamarck et Decand. fl.
franc. 4. p. 44.

Crepis calycibus muricatis. Hall. stirp. helv.
n. 32.

erm. Borstentrager Pippau.

all. Crépide hérissée.

ab. In Margraviatu superiore circa Basileam
extra portam Richanam versus Crenzach
ad vinearum semitas t. Lachenalio.

lor. Julio, Augusto. Semina Augusto, Sep-
tembri matura. ♂

Radix fibrosa, pallide fusca. *Caulis* cubi-
lis, tripedalis, erectus, striato-fulcatus, fo-
sus, ramosus, superne in ramis cum primis se-
sus. *Folia* radicalia caulinaque inferiora bre-
ter petiolata, glabra, læte viridia, runcinata,
et profunde inciso-dentata: lobo supremo
maximo; caulina superiora sessilia, semiample-
caulia, basi hastata, subinciso-sinuato dentata;
prema hastata, lanceolata, acuminata, inte-
ra, margine et costa utrinque setosa. *Flores*
ulem ramosque terminantes, mediocres, pal-
de lutei. *Calycinae* squamæ lineari-lanceolatae,
ubarinosae, dorso setis albidis exstantibus mu-
tae. *Pappus* stipitatus, mollissimus.

bf. Planta in horto culta, mutationi non succubuit,
facile setis praesertim in pedunculis et calyce ex-
stantibus numerosis, a Crepide tectorum di-
stinguenda.

Nota. Ad rarissimas Florae cives pertinet.

210. *CREPIS Dioscoridis* foliis radi-
calibus runcinatis, caulinis hastatis, ca-

338 SYNGENESIA. POLYG. ÆQUALIS.

lycibus subtomentosis. *Linn. Sp. pl.*
1133. — *Pollich. fl. pal. n.* 750.

Hieracioides vulgaris annuo cichorei foli
flore luteo subtus purpurascente. *Val.*
act. 715. *bot. parif.* 102.

Hieracium I. *Tabernam. p.* 491. *Icon bot.*

Icon. Schkuhr. bot. Handb. t. 222 an?

Germ. Dioscorids - Pippau.

Gall. Crépide de Dioscoride.

Hab. In vineis passim, nec non in agris c
careo - argillaceis lapidosus. In vineis pro
Durlach, Grözingen, Berghausen et ali
In Marggraviatu superiore circa *Basilea*
in agris lapidosus trans pontem *Wiesae*
sus *Weil.* Extra portam *Johanneam* ven
St. Louis, Huningen. t. Lachenalio. In P
latinatu in vineis circa *Dürkheim t. P.*
lichio.

Flor. Junio, Julio, Augusto. *Semina* Aug
sto, Septembri matura. ☉

Caulis palmaris, pedalis, erectus, foli
rius, striato-fulcatus, glaber, inferne sub
bescens, ad exortum ramorum angulosus. *Ram.*
alterni, distantes, patentes, simpliciusculi, v
in duos tresve ramulos subdivisi. *Folia* radical
plurima, in rosulam expansa, petiolata, ovata
lanceolata, 5, 6, ad 7 dentibus brevibus
acutis notata, quandoque profundius secta et
am semipinnata, glaberrima, subtus pallidior
nervo eminente percursa; *Caulina* alterna, i
feriora sessilia, ceteris similia, reliqua valde
gusta, longa, linearia, integerrima, raro un
alterove dente instructa, auriculis longis incur
vis caulem amplexantia, qui tamen hami
superioribus foliis saepe nulli. *Flores* caulem r

SYNGENESIA. POLYG. ÆQUALIS. 339

fque terminantes, aliquantum majores quam
equente specie. *Calyx* tomentolus vel fari-
us, *squmis* exterioribus fere capillaribus,
oribus, exiguis. *Corollularum* ligulæ flavæ,
iffime extus purpurascentes. *Semina* duplo
giora quam in sequente specie, valde graci-
Pappus capillaris, sessilis. *Receptaculum*
um.

Descriptionem plantæ debemus b. Lachenalio,
opprime eam cum affini sequenti specie enucleavit
ov. Act. Helvet. I. p. 285.

ota. *Crepis* Dioscoridis Roth. fl. germ. 3. p.
255. non huc, sed ut ex descriptione ejus facile
patet, ad variationem *C. tectorum* pertinet.

II. *CREPIS tectorum* foliis lanceola-
to-runcinatis sessilibus lævibus: inferio-
ribus dentatis, caule glabro. *Linn. Sp.*
pl. 2. p. 1135. — *Pollich. fl. pal.* n. 751.

Crepis caule ramosissimo, foliis glaberrimis
teneris integris et dentatis. *Hall. stirp.*
helv. n. 33.

Hieracium majus erectum angustifolium cau-
le lævi. *Bauh. pin.* 127.

Hieracium luteum glabrum seu minus hir-
sutum. *Bauh. hist.* 2. p. 1024.

Hieracium II. *Tabernæmont.* p. 491. *Icon*
bona.

on. *Curtis. fl. Lond.* t. 327.

rm. *Gemeiner Pippau.*

III. *Crépide des toits.*

ab. Utrunque in pascuis, pratis, viis, ruder-
atis, hortis, muris, segetibus, inter stipulas
frequens.

SYNCHENESIA. POLYG. ÆQUALIS.

Herb. Julio, Augusto, Septembri. Semina Augusto, Octobri matura. ☉

Caulis palmaris, pedalis, cubitalis et altior, sulcatus, glabriusculus, ramosus. *La radicalia* petiolata, lanceolata, pinnatifida plus minusve profunde dentata, glabra, basi subhirsuta; *caulina inferiora* radicebus familia, breviter petiolata; *media* sessilibus amplexicaulia, sagittata seu auriculata, sinu dentata; *superiora* sensim angustiora, integerrima, longe acuminata, basi dentata; *superiora minima*, auriculata. *Flores* caulem ramosque terminantes, parvisculi, sulphurei, subtus pallidiores, interdum ex luteo-purpureascentes. *Bractea* squamæ exteriores interioribus tripliciter breviores, angustæ, acutæ, laxæ, pubescentes; interiores arctiores, lineares, margine viridato dorso fuscae, pilis quasi muricatis. *Semina* parva, gracilia, striata, obtusa. *Pappus* sessilibus capillaris, mollissimus.

Planta ceterum maxime polymorpha, mire variata pro diversitate loci natalis, florentiæ tempore, magnitudine, foliorum figura, incisuris, hirsutia, calycibus glabro vel hirsuto etc. In solo pinguiori sæpe pedalis, angulata, sulcata, glabra, aut saltem radice versus hirsuta; in macri vero macra, digitalis, palmaris, sæpe villosa, canescens.

Conf. Nov. Act. Helv. I. p. 282. ibi III. L. chenalius secundum Hallerum speciem sequentem cum hacce sub variatione combinavit.

1212. CREPIS virens foliis runcinatis glabris amplexicaulibus, calycibus submentolis. *Linn. Sp. pl. 2. p. 1134.*

Lapsana capillaris. *Linn. Sp. pl. ed. I. to 2. p. 812.*

SYNGENESIA. POLYG. ÆQUALIS. 342

Hieracioides foliis variis glabris, caule ramossissimo. *Hall. enum. helv. p. 751. n. 4.*

Hieracium minus glabrum *Bauh. Prodr. p. 63.* — foliis eleganter virentibus. *Bauh. pin. 127.*

Hieracium IV. *Tabernæmont. p. 491. Icon. germ. Grüner Pippau.*

all. Crépide verdâtre.

ab. Utrunque cumprimis in Margraviatus superioris agris, vergente ætate frequens. . .

lor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ☉

Praecedenti nimis omnino affinis, a qua se differre videtur *Caulibus* humilioribus, *ra-*
floribus, glabris, *Foliis* minus incis, immo
rumque leviter saltem dentatis amoene vi-
bus, glaberrimis, *Calyce* glabro vel leniter
descendente, *Floribus* minoribus.

Omnino et haec polymorpha, varians caule sub-
plici, unifloro nudo, quae est *Crepis* uniflora
Will. fl. Paris. II. p. 410. — et foliis radicalibus
minisque infimis magis incis, subpinnatifidis, quae
est dubio *Crepis* pinnatifida. Willden. Linn.
p. pl. 3. p. 1605.

213. CREPIS biennis foliis runcinatis
pinnatifidis scabris: basi superne dentatis,
calycibus muricatis. *Linn. Sp. pl. 2. p.*
1136. — *Pollich. fl. pal. n. 752.*

Crepis foliis ad terram pinnatis, superne
amplexicaulibus, pinnatis hastatis. *Hall.*
fl. helv. n. 31.

Hieracioides foliis variis subasperis, caule
altissimo folioso et multifloro. *Hall. Enum.*
helv. p. 753. n. 3.

342 SYNGENESIA. POLYG. ÆQUA

*Hieracium majus erectum latifolium
gustifolium caule aspero. Bauh. p*

*Hieracium multis aliis Cichorium
hirsutum et glabrum vel minus hirsutum
Bauh. hist. 2. p. 1024. Icon.*

*Hieracium chondrilloides. Tabernam
Icon bona.*

Ger m. Zweyjähriger Pippau.

Gall. Crépide bisannuelle.

*Hab. Utrunque in pratis locisque fertili
gramineis subhumidis, riguis. ♂*

Caulis 3-4 pedalis, erectus, angulatus, plus minusque hirsutus, quam glabriusculus, læte viridis, striis purpureo pictus, ramosus. Rami alterni, divisi. Folia spida, quandoque glabriuscula, læte viridicalia plurima, petiolata; caulina alter filia, amplexicaulia, pinnatifida, acuminatis retroverfis sinuato-dentatis, superiorioribus, extrema triangulari, integra, inferioribus angustioribus sublinearibus; Supramea lanceolata, basi subdentata. Flores ramosque terminantes, speciosi, ut lutei. Calycinae squamæ scabræ, dorsum nigris ciliatæ. Pappus capillaris, sessilis. Receptaculum subcylindrica, incurviuscula, fusca. Receptaculum pilis tenuibus præditum.

Hujus speciei insignem variationem legi in acclivibus retro Gundeldingen, Caulibus radice pluribus, pedalis paucifoliis et paucifoliis radicalibus numerosis in orbem sparsis, ex petiolo in obtusissimum sinem exeuntibus, qualis tenerrimis, glabris præter oram subhileviter et rariter dentatis, dentibus tamen retracaulinis paucis similibus, vix hamatis. In floribus differentia apparebat. Lachenal. Nov. Act. 1. p. 281.

SYNGENESIA. POLYG. ÆQUALIS. 343

Distinguitur a *C. tectorum*, Radice bienni, foliis caulinis majoribus, pinnatifidis, segmentis re-
versis, extremo triangulari, integro, Floribus
cinque semper luteis. Receptaculo pilis tenuibus
nudo.

Flores apibus melliferi.

CCCLXV. H Y P O C H A E R I S.

Linn. Gen. pl. ed. Schreb. n. 1246.

Calyx communis subrotundus, imbricatus,
intus ventricosus: *squamis* lanceolatis, acutis.
Corolla composita imbricata, uniformis: *Co-
mmissuris* hermaphroditis, æqualibus, numerosis.
Corolla propria monopetala, ligulata, linearis, trun-
cata, quinquedentata. *Calyx* fructiferus con-
vexus, globofo-acuminatus. *Semina* oblonga.
Stylus plumosus, stipitatus. *Receptaculum* pa-
ceum. *Paleis* lanceolato-linearibus, seminum
longitudine.

214. *HYPOCHAERIS maculata* caule
subnudo: ramo solitario, foliis ovato-
oblongis integris dentatis. *Linn. Sp. pl.*
2. p. 1140. — *Pollich. fl. pal. n. 754.*

Hieracium caule unifloro, foliis dentatis,
hirsutis. *Hall. stirp. helv. n. 2.*

Hieracium alpinum, foliis dentatis flore mag-
no. *Bauh. pin. 128.* — *Prodr. 65.*

Hieracium VIII. *Tabernæmont. p. 493. Icon.*
rudis.

con. Oeder. fl. Dan. t. 149. — *Hall. l. c. tab*
1. *variatio helvetica.*

344 SYNGENESIA. POLYG. ÆQUALIS.

Germ. Geflecktes Ferkelkraut,

Gall. Porcelle tachée.

Hab. In pascuis graminosis altioribus. In montosis gramineis *auf dem Kaltenbrunn*. In Margraviatu superiore in Brisgovia in pratis altissimis a pago *Schellingen am Kaiserstuhl* denominatis. In pascuis montis *Felsberg* et *Belchen*. In Principatu Fürstenbergensi in pratis montosis passim. In Alsatia montis *Ballon* summis pascuis, t. *Lachen*. In Palatinatu circa *Lautern* in pratis et pascuis siccioribus. Prope *Hochspey* in collibus gramineis aridis. t. *Pollich*. In Sponhemia retro *Winningen* ad Mosellam in pratis montosis.

Flor. Junio, Julio, *Semina* Augusto matura.

Radix ramosa, crassa, extus fusco-nigricans, intus flavicans. *Caulis* sesquipedalis, cubitalis, erectus, crassiusculus, teres, striatus, hirsutus, sæpe maculatus, maculis fusco-purpureis, subaphyllus, folio uno alterove præditus simpliciusculus, versus florem incrassatus, in medio, ramis sæpe simplici longiori, auctus. *Folia radicalia* plurima, terræ incumbentia, in petiolum decurrentia, caulina alterna, sessilia, semiamplexicaulia, omnia ovato-lanceolata, marginem riter argute denticulata, utrinque hirsuta, late viridia, maculis sæpissime purpurascentibus picta. *Flos* solitarius, terminalis, speciosus, præmarinus, amplus, luteus. *Calycinae* squamæ lanceolatae, dorso pilosæ, scabræ.

Nota. Hypochæris helvetica a nostra planta vi specie differt, caulis in ea simplex, uniflorus, inferne magis foliosus.

Germani Herbam Costam vocant eaque in peccatoris morbis et ad phthisin utuntur. C. Hoffmann mat. med. p. 270.

215. HYPOCHAERIS *glabra* calycibus oblongis imbricatis, caule ramoso nudo, foliis dentato-sinuatis. *Linn. Sp. pl. 2. p. 1140.* — *Pollich. fl. pal. n. 755.*

Hieracium minus dentis Leonis folio oblongo glabro. *Bauh. pin. 127.*

Hypochaeris chondillæ folio, parvo flore. *Vaill. act. 1721. p. 214.*

non. Curtis. fl. Lond. t. 145. — *Oeder. fl. Dan. t. 424.* — *Schkuhr. bot. Handb. t. 225.*

Berm. Glattes Ferkelkraut.

Ball. Porcelle glabre.

Tab. In Margraviatu superiore in agris et pascuis arenosis Rhenum versus inter *Neuenburg* et *Zienken* rarius t. *Vulpio*. Prope *Basileam* in agris humidis retro pagum *Alschweiler* t. *Lachenalio*. In Palatinatu inter *Sembach* et *der Langen Meil* in pascuis glareosis siccis, nec non circa *Sambach*, *Katzweiler* in agris t. *Pollichio*.

Flor. Junio Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix fusiformis, pallide fusca, fibris paucis aucta. *Caules* sæpe plures, palmares, spinosissimi, suberecti, teretes, glabri, glauci, subnudi, in duos, rarius tres ramos simplicissimos lividi. *Folia* radicalia numerosa, supra terram expansa, subsessilia, sinuato-dentata, obtusiuscula, versus apicem latiuscula, glabriuscula, margine ciliata; *caulina* vix ulla, minima. *Flores* in pedunculis superne subincrassatis, squamosis, terminales, solitarii, parvuli, lutei. *Caryx* primum cylindricus, fructiferus oblongo-ovatus magnus. *Squamæ* ovato-lanceolatae, inaequales, apice rubellæ, læves. *Semina* sublinea-

346 SYNGENESIA. POLYG. ÆQUALIS.

ria, basi acuminata, scabra. *Receptaculum* paleaceum, paleis concavis, lanceolatis, acuminatis, nitidis, deciduis, longitudine pappi. *Pappus* plumosus, rigidulus, radii sessilis, disci longissimipitatus.

Ad rariores omnino Floræ cives pertinet.

1216. *HYPOCHAERIS radicata* foliis

runcinatis obtusis, scabris, caule ramoso nudo, lævi, pedunculis squamosis. *Lin.* *Sp. pl.* 2. p. 1140. — *Pollich. fl. pal.* n. 756.

Hypochaeris foliis semipinnatis hirsutis, caule nudo brachiato. *Hall. stirp. helv.* n. 3.

Archirophorus radicans. *Scop. fl. Carn.* n. 987.

Hieracium dentis Leonis folio obtuso majus. *Bauh. pin.* 127. — *Mapp. fl. Alf.* p. 142.

Hieracium VII. *Tabernæmont.* p. 492. *Icon.*

Icon. Curtis. fl. Lond. t. 152. — *Oeder. fl. Dan.* t. 150.

Germ. Wurzelndes Ferkelkraut.

SYNGENESIA. POLYG. ÆQUALIS. 347

bus obtusiusculis, hirsuta, scabra, lacte virida. *Pili* erecti, simplices, e punctis asperis prominentes. *Flores* caulem ramosque terminantes, compli, majusculi, lutei. *Calycinæ* squamæ ovate, acutæ, apice rufæ, carina ciliatæ. *Semen* longum, castaneum, striatum. *Pappus* stipitatus, plumosus. *Receptaculum* paleaceum, pappi membranaceis, longis, concavis.

Bonum pecoribus pabulum præbet. Flores apibus largiuntur.

CCCLXVI. ARNOSERIS.

Gaertn. de fruct. et semin. pl. 2. p. 355.

Calyx communis calyculatus ovali-subglobose: *squamis interioribus* numerosis, erectis, linearilanceolatis, acutis; *exterioribus* paucioribus, 11-12, minutis, interioribus adpressis.

Corolla composita imbricata, uniformis: *Cellulis* hermaphroditis copiosis, densis, æqualibus. *Propria* monopetala, ligulata, truncata, sinquedentata. *Calyx* fructiferus torulosus, globose connivens. *Semina* angulata, sulcata, membranacea coriacea tecta, pappo destituta. *Receptaculum* nudum, planum, excavato-punctatum, subrotundum.

217. *ARNOSERIS pusilla* caule diviso nudo, pedunculis incrassatis.

Lapsana foliis ovatis, dentatis, caulibus nudis, sub flore tumentibus. *Hall. stirp. helv. n. 4.*

Lapsana pusilla. Willden. Linn. Sp. pl. 3. p. 1623.

348 SYNGENESIA, POLYG. ÆQUALIS.

Lampfana minima. Lamarek et Decandolla
fl. franc. 4. p. 3.

Hyoseris minima. Linn. Sp. pl. 2. p. 1138. —
Pollich. fl. pal. n. 753.

Hieracium minus folio subrotundo. Bauh.
pin. 127.

Hyoseris I. Tabernemont. p. 490. Icon rud.

Hieracium minimum. Clus. hist. 2. p. 141. fig.

Icon. Oeder. fl. Dan. t. 201. — Schkuhr. bot.
handb. t. 224.

Germ. Kleines Lämmerkraut.

Gall. Arnoseride fluette.

Hab. In agris et arvis sterilioribus sabulosis
et transrhenanis passim frequens. Prope
Carlsruhe circa Mühlburg, Eckenstein.
Schreck, Linkenheim, Graben et alibi.

Flor. Julio, Augusto. Semina Septembri ma-
tura. ☉

Caules ex una radice plures, digitales, pal-
mares, spithamæi, erecti, aphylli, glabriusculi
superne incrassati, ramo uno alterove nude
superne incrassato præditi. *Folia radicalia* nu-
merosa, in orbiculum congesta, subpetiolata
terræ incumbentia, ovali-lanceolata, ob-
tusa, versus basin angustiora, dentata, subgla-
bra, crassiuscula, glauco-viridia. *Flores* cau-
lem ramosque terminantes, solitarii, pallide
lutei, mediocres.

Omnino a genere *Hyoseris* et *Lampfanae* di-
pertium fructificationis structuram longe diversam, re-
movenda erāt.

Bonum ovibus pabulum.

SYNGENESIA. POLYG. ÆQUALIS. 349

CCCLXVII. L A P S A N A.

Linn. Gen. pl. ed. Schreb. n. 1247.

Calyx simplex, octophyllus, erectus, connivens, basi calyculatus, squamulis paucis brevissimis. *Corolla composita* imbricata, uniformis: *Corollulis* hermaphroditis circiter 16, æqualibus. *Propria* monopetala, ligulata, truncata, quinquedentata. *Calyx* fructiferus, foliis angulo obtuso carinatis. *Semina* oblonga, deorsum acuminata, obsolete angulata, striata, calva. *Pappus* nullus. *Receptaculum* nudum, planum.

1218. LAPSANA communis calycibus fructus angulatis, pedunculis tenuibus ramosissimis. *Linn. Sp. pl. 2. p. 1141. — Pollich. fl. pal. n. 757.*

Lapsana caule brachiato, foliis ovatis longe petiolatis, petiolis pinnatis. *Hall. stirp. helv. n. 6.*

Soncho affinis *Lapsana domestica*. *Bauh. pin. 124.*

Lampfana. *Dodon. — Mapp. fl. Alf. p. 162.*

Sonchus sylvaticus I. II. III. *Tabernæmont. p. 500. 501. Icon.*

Icon. Curtis. fl. Lond. — Oeder. fl. Dan. t. 500.

Germ. Gemeiner Milchen. Rainkohl.

Gall. Lampfane. Herbe aux Mamelles.

Hab. Utrunque in ruderatis, muris, hortis, cultis, sylvis cæduis, ad sepes abunde,

350 SYNGENESIA. POLYG. ÆQUALIS.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix fibrosa. *Caulis* cubitalis, tripedalis et altior, erectus, fulcato - angulatus, ramoso subdichotomus, plus minusve hirsutus. *Folia* inferiora et radicalia versus basin profunde, dentuno alterove incisa, ovata, obsolete sinuato dentata, obtusiuscula, villosa; *caulina* alterna superiora ovali - lanceolata, sinuato - dentata, *ramæ* minora, lanceolata, angustiora, integriuscula. *Flores* mediocres, flavi, paniculati. *Pedunculi* divisi, tenues, glabriusculi. *Semina* subincurvata.

Pharm. LAMPSONÆ *Herba recens.*

Emollientes vires habet. Papillas exulceratas mammarum, imposita, sanat, hinc germanis olim Milche vocata, hodie fere exoleta.

Planta tenella ab omni pecorum genere amatur. Flores apibus melliferi.

CCCLXVIII. C I C H O R I U M.

Linn. Gen. plant. ed. Schreb. n. 1251.

Calyx communis calyculatus, cylindraceus, squamis subocto, anguste lanceolatis, æqualibus, cylindrum constituentibus; quinque vero incumbentibus, brevioribus. *Corolla composita*, plana, uniformis: *Corollulis* hermaphroditis viginti in orbem positis. *Propria* monopetala, ligulata, truncata, profunde quinquedentata. *Calyx fructiferus* cylindraceus, apice connivens. *Semina* compressa, angulis acutis. *Pappus* obsolete pilosus, quinquedentatus. *Receptaculum* subpaleaceum.

19. CICHORIUM *Intibus* floribus geminis sessilibus, foliis runcinatis, Linn.
Sp. pl. 2. p. 1142. — *Pollich. fl. pal. n.* 758.

Cichorium foliis pinnatis, pinnis triangularibus dentatis, floribus sessilibus. Hall.
fl. alp. helv. n. 1.

Cichorium sylvestre, f. officinarum. *Bauh. pin.* 126. — *Mapp. fl. Als. p.* 74.

Cichorium l. II. *Tabernæmont. p.* 467. *Icon. n. Curtis. fl. Lond. t.* 241. — *Oeder. fl. Dan. t.* 907. — *Blackw. herb. t.* 183. — *Schkuhr. bot. handb. t.* 226.

r m. *Wegwart. Cichorien. Hundläufte.*

ll. *Cichorée sauvage.*

b. Utrinque ad vias et agrorum margines frequens.

o r. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ¶

Radix fusiformi-cylindrica, etiam ramosa, harnæa, et longior, descendens, extus luteus, intus albida. *Caulis* sesquipedalis, cubitalis, subitalis, erectus, striato aut sulcato-angulosus, nudus, ramosus. *Rami* alterni, copiosi, subtuoli. *Folia radicalia* plurima, runcinata; *ulina* alterna, amplexicaulia, dentata, mucronata asperiuscula, *ramea* minora, angustiora, curviuscula, minus aspera. *Flores* sessiles plerumque bini, quandoque tres, etiam solitarii, caulis et ramorum latera, speciosi, cærulei, rarius rosei, et albi. *Calycinae* squamæ hispidæ.

harm. **CICHORII** *Radix; Herba; Flores; Semina.*

Planta amariuscule resolvens, detergens, tonica, viscerum obstructione, atonia intestinorum præstans.

352 SYNGENESIA. POLYG. ÆQUALIS.

Radices siccatæ, concisæ, tostæ, potum Cassiæ æmulum præbent hodie notissimum, jam ultra quingenta annos ab hortulano principis de Schwarzburg-Sondershausen L. D. Timme inventum et evulgatum.

Planta culta mitior, tenerior, præsertim oleribus et acetariis inservit.

Flores apibus melliferi.

1220. CICHORIUM *Endivia* floribus foliariis pedunculatis, foliis integris crenatis. *Linn. Sp. pl.* 2. p. 1142.

Cichorium latifolium f. *Endivia* vulgaris. *Bauh. pin.* 125.

β. *Endivia* crispa. *Bauh. pin.* 125.

Icon. Regnault. bot.

Germ. Endivie. β. krause Endivie.

Gall. Chicorée endive. β. Chicorée frisée.

Hab. Colitur in hortis et vineis passim.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Planta in culinis notissima.

CCCLXIX. A R C T I U M.

Linn. Gen. plant. ed. Schreb. n. 1253.

Calyx communis globosus, imbricatus: *squamis* lanceolatis, desinentibus in aculeos subulatos longos, apice reflexo-hamatos. *Corolla composita* tubulata, uniformis. *Corollulis* hermaphroditis, æqualibus. *Propria* monopetala, tubulata; *tubo* tenui longissimo; *limbo* ovato-quinquefido: *laciniis* linearibus, æqualib

Caly

Stylis fructiferus connivens. *Semina* verticali-
ter pyramidata, angulis duobus oppositis obli-
atis, extrorsum gibba. *Pappus* simplex, se-
ne brevior, obsoletus, rigidulus. *Recepta-
culum* paleaceum, planum: *Paleis* fetaceis.

21. *ARCTIUM tomentosum* calyci-
bus arachnoideo-tomentosis.

Arctium Bardana. Willden. Linn. Sp. pl. 3.
p. 1632.

Lappa tomentosa. Lamarck et Decandolle,
fl. franc. 4. p. 57.

Arctium Lappa β. Linn. Sp. pl. 2. p. 1143.

Lappa major montana capitulis tomentosis.
Bauh. pin. 198.

*Lappa major, montana, capitulis minoribus
rotundioribus et magis tomentosis*. *Mapp.*
fl. Alf. p. 163.

o n. *Oeder. fl. Dan. t.* 642. — *Curt. fl. Lond.*
t. 258. — *Schkuhr. t.* 227.

r m. *Spinnen-Klette*.

z ll. *Bardane a têtes cotonneuses*.

z b. Utrique in ruderatis, ad vias, pagos
et sepes. Abunde ad castella destructa prope
Phillipsburg, Kehl, Altbreisach.

l o r. Julio, Augusto. *Semina* Septembri ma-
tura. ♂

Radix fusiformi-cylindrica, pedalis, de-
currens, fibris filiformibus munita, extus nigri-
bus, intus albida. *Caulis* tripedalis, bicubita-
tus, crassius, erectus, striato-fulcatus, villosus,
tomentosus. *Folia* inferiora longe petiolata, cor-

354 SYNGENESIA. POLYG. ÆQUALIS.

data, integra, præmagna, ampla, margine lobata, undulata, superne saturate viridia, subtus cinerea, venosa, hispidula, superiora ovata, tusa, minora. *Flores* axillares terminalesque corymbosi, breviter pedunculati, soliosi, simpliciter, purpurei, rarius albidi. *Calyces* pulcherrime arachnoideo-tomentosi.

Pharm. BARDANÆ Radix; Semina.

1222. *ARCTIUM majus* calycibus glabris, floribus terminalibus solitariis pedunculatis.

Arctium Lappa. Linn Sp. pl. 2. p. 1143. Pollich. fl. pal. n. 759.

Lappa major. Lamarck et Decandolle bot. franc. 4. p. 57.

Lappa officinalis. Allion. fl. pedem. n. 59.

Lappa major, Arctium Dioscoridis. Bapin. 148. — Mapp. fl. Alf. p. 163.

Lappa major. Tabernæmont. p. 1157. Icon. Blackw. t. 117. — Schkuhr. t. 227.

Ger m. Grofse gemeine Klette.

Gall. Bardanne a grosses têtes.

Hab. Utrunque ad vias, sepes, rudrata, aggeres, frequens.

Flor. Julio, Augusto. *Semina* Septembri, Octobri matura. ♂

A præcedente facile dignoscitur *Calyces* glabris.

Pharm. BARDANÆ Radix; Folia; Semina.

Radix subdulcis, subaustera, blande resolvens, sternens.

Recens radix in taleolos secta, a monticolis Margaviatus superioris, contra oculorum palpebrarumque florem, inflammationem velut amuletum circa collum suspensum, tanquam optimum remedium applicatur.

Caules juniores s. turiones deraso cortice cocti aut par asparagorum cum sale aut aceto oleoque imbuti, satum cibum præbent, vulgo quidem neglecti.

Foliorum succus in ulceribus in Gallia laudatur.

Folia amplissima pecoribus detestata. Schreb. am l. 3.

223. *ARCTIUM minus* calycibus glabris, floribus racemoso-corymbosis, congestis.

Lappa minor. Lamarck et Decandolle fl. franc. 4. p. 57.

Arctium minus. Schkuhr. bot. handb. 3. p. 49. con. Schkuhr. l. c. t. 227.

erm. Kleine Klette.

Gall. Bardane a petites têtes.

lab. Utrinque ad viarum margines ficos, nec non in ruderatis.

lor. Julio, Augusto. Semina Septembri matuta. ☉

Dignoscitur a præcedente specie Caulibus foisque minoribus. Floribus multo semper minoribus, 5-6 in racemum congestis.

obs. Hæ omnes tres species, in horto botanico Carlsruhano, per incomparabilem hortiInspectorem Schweyckert nuper pro dolor defunctum, cultæ, mutationi non succubuerunt.

Nota. Species minor tertia, non confundenda cum Bardana minori medicorum quæ est *Xanthium strumarium*. Linn.

CCCLXX. SERRATULA.

Linn. Gen. pl. ed. Schreb. n. 1264.

Calyx communis oblongus, subcylindricus imbricatus; *squamis* lanceolatis, acutis, se obtusis muticis. *Corolla composita* tubulosa uniformis. *Corollulæ* hermaphroditæ, æquales. *Propria* monopetala, infundibuliformis: *tubo* inflexo, *limbo* ventricoso, quinquefido. *Calyx* fructiferus immutatus. *Semina* obovata. *Pappus* sessilis, plumosus. *Receptaculum* paleaceum, planum.

1224. *SERRATULA tinctoria* foliis lyrate-pinnatifidis, pinna terminali maximâ, flosculis conformibus. *Linn Sp. pl.* 2. p. 1144. — *Pollich. fl. pal. n.* 760.

Carduus inermis, foliis glabris ferratis, superioribus semipinnatis. *Hall. stirp. helv.* n. 163.

Carduus tinctorius Scop. fl. Carn. n. 1015.

SYNGENESIA. POLYG. ÆQUALIS. 357

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. 7

Radix maxime fibrosa, fusca. *Caulis* sesquipedalis, cubitalis, et altior, erectus, angulosus, gracilis, superne ramosus. *Folia* in eodem sæpe individuo varia: *inferiora* petiolata, lyrata, ovato-lanceolata, aut lyrato-pinnatifida, *superiora* sessilia, minora, integra aut pinnatifida lyrata, argute serrulata, interdum omnia subserrulata, margine serrulata. *Flores* caulem terminantes, corymbosi, mediocres, purpurei, aut purpurei, rarius albi. *Calycinae* quamvis glabræ, inermes, coloratæ, violaceo-æruleæ.

Planta cum alumine flavum colorem præbet.

1225. *SERRATULA arvensis* foliis dentatis spinosis. *Linn. Sp. pl.* 2. p. 1149. — *Pollich. fl. pal.* n. 761.

Cirsium caule anguloso ramoso, foliis semipinnatis, pinnis angulosis spinosis, calycibus longis inermibus. *Hall. stirp. helv.* n. 171.

Carduus radice repente, foliis lanceolatis dentatis margine aculeatis. *Linn. fl. Lapp.* 293.

Cirsium arvense, Sonchifolio, radice repente, flore purpurascens et albo — et caule tuberoso. *Mapp. fl. Als.* p. 76.

Carduus vinearum repens, Sonchi folio. *Bauh. pin.* 387. *Prodr.* 156.

Carduus arvensis. *Tabernæm.* p. 1079. *Icon. zoon.* *Curtis. fl. Lond.* (sub *Carduo arvensi*). *erm. Ackercharten. Felddistel.*

358 SYNGENESIA. POLYG. ÆQUALIS.

Gall. Serrete des champs. Cirse des champs

Hab. Utrunque in agris et vineis passim, fossarum aquaticarum et viarum marginibus frequens.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ♀

Radix cylindrica, profunde in terram descendens, latissime repens, sordide albida. *Caulis* tripedalis tetrapedalis et altior, erectus, hirsutus, lanuginosus, superne angulatus, lævis, ramosus. *Folia* sessilia, alterna, lanceolata, inciso-subpinnatifida, sinuata, subundulata, sæpe crispata dentibus argute spinulosis praedita, superne glabriuscula, læte viridia, subtus pallidiora; suprema parvula, integriuscula. *Flowers* caulem ramosque terminantes, pallide purpurei, quandoque albi, mediocres. *Calycis* squamæ numerosæ, imbricatæ, adpressæ, lanceolatae, apice appendiculatæ, mucrone terminatæ, cæruleo-purpureæ. *Receptaculum* paleaceum: *Paleis* longiusculis planis splendentibus præditum. *Pappus* plumosus, sessilis.

Habitus Cardui aut Cnici, a quibus autem omnino recedit, Receptaculo non piloso, sed paleaceo.

Magnum sæpe agrorum ob radices longe lateque sese propagantes vitium, sollicitè eradicandum.

Planta junior suis gratum pabulum.

Flores odorati apibus melliferi.

CCCLXXI. C A R D U U S.

Linn. Gen. pl. ed Schreb. n. 1254.

Calyx communis ventricosus, imbricatus squamis numerosis, lanceolatis, acuminatis, spinosis. *Corolla composita* tubulosa, uniformis

Corollulæ hermaphroditæ, subæquales, reflexæ. *Propria* monopetala, infundibuliformis: tubo tenuissimo, limbo erecto, basi ovato, quinque-
 lido: laciniis linearibus, æqualibus: unica profundius separata. *Calyx* fructiferus parum concin-
 uens. *Semina* obovata, tetragona, oblitis angulis oppositis duobus. *Pappus* capillaris, fessilis, longissimus. *Receptaculum* pilosum, lanum.

b. Plures CARDUI Linneani PAPPUM plumosum habent, ad CNICOS rectius referendi.

† *Foliis decurrentibus.*

226. *CARDUUS nutans* foliis semidecurrentibus spinosis, floribus cernuis: squamis calycinis superne patentibus. *Linn. Sp. pl.* 2. p. 1150. — *Pollich. fl. pal.* n. 763.

Carduus foliis semipinnatis, pinnis angulosis, spinosis, floribus nutantibus, calycibus pungentibus, patulis. *Hall. stirp. helv.* n. 167.

Carduus spinosissimus latifolius sphaerocephalus vulgaris. *Bauh. pin.* 385.

Carduus nutans flore rubro-et albo. *Bauh. hist.* 3. p. 56.

Icon. Oeder. fl. Dan. t. 675. — *Barrel. Icon.* 1116.

Germ. Ueberhängende Distel. Bisamdistel.

Gall. Chardon penché.

Hab. Utrique in pascuis graminosis et ad agrorum viarumque marginalia passim frequens.

366 SYNGENESIA. POLYG. AQUALIS.

Flor. Julio, Augusto. *Semina* Augusto Septembri matura. ☉

Caulis sesquipedalis, cubitalis et altior, erectus, subramosus. *Folia* ad caulem decurrentia, aut semidecurrentia, alterna, oblongo-lanceolata, pinnatifida, plicato-undulata, margine spinosa, utrinque scabriuscula. *Pedunculi* terminales, solitarii, subinermes, longiusculi, temontosi, superne cum floribus solitariis deorsum nutantes. *Flores* speciosi, magni, purpurei, raro albi, moschum subredolentes.

Flores apibus acceptissimi, multam mellis ceræque materiam præbent.

1227. *CARDUUS polyacanthos* foliis decurrentibus sinuato-pinnatifidis margine spinosis, calycibus solitariis pedunculatis erectis villosis. *Schreb. Spic. fl. Lips.* p. 15. *Wetterauische flora.* 3. p. 159.

Carduus Acanthoides. *Linn. Sp. pl.* 2. p. 1150. — *Poll. fl. pal. n.* 765. — *Smith. fl. brit.* 848. *Lamarck et DeCandolle. fl. franc.* 4. p. 79.

Carduus crispus foliis decurrentibus sinuatis margine spinosis, calycibus globosis pedunculatis solitariis erectis. *Huds. fl. Angl.* 2. p. 350.

Carduus polyacantha *Theophrasti.* *Tabern.* p. 1080. *Icon bona.*

Carduus caule crispo. *Bauh. hist.* 3. p. 59. videtur.

Carduus spinosissimus vulgaris, Polyacantha *Theophrasti.* *Lobel. Icon.* 21.

Icon. *Jacqu. fl. Austr. t.* 249. — *Schkuhr. bot. handb. t.* 228.

SYNGENESIA. POLYG. ÆQUALIS. 362

erm. Vielschlichte Wegdistel.

all. Chardon polyacanthé.

ab. Utrunque in ruderatis ad vias passim frequens, præsertim circa *Carlsruhe, Bruchsal, Mannheim, Frankenthal, Worms* et alibi.

or. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caulis tripedalis, tetrapedalis, et altior, rectus, angulosus, subtomentosus, utrinque iis decurrentibus alatus, dentato-spinosus. *Folia* alterna, decurrentia, lanceolata, natifida: *laciniis* undulato-plicatis, subcrisis, margine spinulis numerosis rigidis punctibus obsessis, superne saturate viridia, subpallidiora, costa media subpubescente. *Flo-*
caulem ramosque terminantes, breviter pedunculati, erecti, uniflori, carduo crispo paulo jores, purpurei, raro albidii. *Calyx* subglossus, imbricatus: *Squamæ* lineares, acutae, vula terminatæ, erecto-patentes, reflexiuscæ, villis tenuibus præditæ.

A. Carduo crispo facile dignoscitur Calycibus gius pedunculatis, solitariis, non aggregatis, Flo-
us semper majoribus.

Nota. Carduus Acanthoides Joh. Bauh. hist. pl. 3. p. 59. a Linnaeo ad nostram plantam citata longe alia planta exotica est, hinc Perill. Schrebero auctore, nomen triviale immutavimus, ne cum Bauhiniana confundatur.

Flores apibus melliferi.

128. CARDUUS *crispus* foliis decurrentibus sinuatis, margine spinosis, floribus aggregatis terminalibus inermibus, *Linn. Sp. pl. 2. p. 1150. — Pollich. fl. pal. n. 765.*

362 SYNGENESIA. POLYG. ÆQUALIS.

Carduus Acanthoides foliis decurrentibus sinuato-dentatis marginē spinosis, calycibus ovatis terminalibus aggregatis sessilibus, squamis acuminatis erectiusculis
Hudsf. fl. Angl. 2. p. 351. — Allion. fl. ped. n. 146.

Carduus foliis ovatis et semipinnatis, subtomentosis spinosis, calycibus mollibus spinosis reflexis. *Hall. stirp. helv. n. 160 omnino (exclus. J. Bauh. Synonymo).*

Icon. Oeder. fl. Dan. t. 621.

Germ. Krause Distel.

Gall. Chardon frisé.

Hab. Utrunque in ruderatis ad sepes, juxta vias, in cultis, hortis, vineis, præsertim in pagorum rejectamentis, passim frequens.

Flor. Junio, Julio. *Semina* Julio, Augusto maturantur. ☉

Sæpe cum præcedenti affini specie confunditur, a qua autem differt, *Caulibus* magis tomentosis, *Foliis* non tam profunde pinnatifidis supra rariter hirtis, saturate viridibus, subtus albidis tomentosis, margine spinulosis, *Floribus* terminalibus, in capitulum aut fasciculum congestis, minoribus, *Calycibus* sessilibus, aut brevissimo pedunculo tomentoso, niveo, imbructis, subrotundo-ovatis, *Squamis* calycinæ lineari-acuminatis, erectiusculis, rarius inferioribus subreflexis, aculeolo miti præditis, mollibus.

Flores vulgo purpurei, rarius albid.

Nota. Cl. Hudsonii descriptio omni ex parte cum nostra planta convenit.

Carduus tenuiflorus foliis decurrentibus sinuatis spinosis tomentosis, calycibus subcylindricis

gregatis sessilibus, squamis lanceolatis erectiusculis. Smith. fl. Britann. 849. ad quem Cl. vir *Carduum acanthoidem* Hudsonii l. c. citat, vix specie diversus a *C. crispo* Linnaei. De calycibus subcylindricis Hudsonus nil dicit, sed de ovatis.

1229. CARDUUS *defloratus* foliis decurrentibus lanceolatis ferratis subspinoso-ciliatis nudis, pedunculis longissimis lanuginosis unifloris. *Linn. Sp. pl. 2. p. 1152.*

Carduus inermis foliis ciliatis, scapo nudo, paucifloro. *Hall. stirp. helv. n. 164. t. 4.*

Cirsium angustifolium. *Bauh. Prodr. p. 155. Mapp. fl. Als. p. 77.*

Cirsium tertium montanum. *Clus. hist. 2. p. 149.*

Icon. Jacqu. fl. Austr. 1. t. 89.

Ger. m. Bergwiesendistel.

Fall. Chardon terne.

Hab. In Margraviatus superioris pascuis graminosis montosis Brisgoviae in monte *Feldberg*. In ditionis Basileensis monte *Wasserfallen* et *Dornach* non infrequens. In Principatus superioris pratis sylvaticis montosis ad *Lacum Bodamicum* prope *Oeningen*, *Stein* et alibi frequens; nec non in Principatu *Fürstenbergensi* in pratis prope *Donaueschingen*, *Fürstenberg*, *Engen* et alibi.

Flor. Junio, Julio. *Semina* Julio, Augusto matura. ☞

Radix fusco-nigricans, longa, fibrosa. *Caulis* bipedales, tripedales et altiores, erecti, angulati, glabriusculi, simpliciusculi, foliosi, abeun-

364 SYNGENESIA. POLYG. EQUALIS.

tes in pedunculum unum, alterumve, longum, erectum, aphyllum, lanuginosum, superne uniflorum. *Folia* per caulem decurrentia, alternatim oblongo-lanceolata, argute serrata, spinulosa, ciliata, acuta, superne pallide viridia, subtus glauca. *Flores* majusculi, erecti, quandoque nutantes, purpurei, raro albidi. *Calycinae* squamae imbricatae, ovato-lanceolatae, acutae, glabrae, virides, apice fusco-purpureascentes, persistenti-erectae.

Variat Foliis plus minusve profunde serratis.

Flores apibus melliferi.

1230. *CARDUUS Personata* foliis decurrentibus ciliato-spinosis: inferioribus profunde pinnatifidis, floribus terminalibus subsessilibus congestis, calycinis squamis reflexis.

Carduus Personata. Jacqu. Lamarck et DeCandolle. fl. franc. 4. p. 85.

Carduus arctioides. Vill. fl. Delph. 3. p. 22.

Arctium Personata. Linn. Sp. pl. 2. p. 1144. Jacqu. vind. 274.

Carduus inermis foliis imis pinnatis, superioribus ovato-lanceolatis serratis, squamis calycinis reflexis. *Hall. stirp. helv. n. 162. t. 3.*

Carduus mollis latifolius Lappæ capitulis Bauh. pin. 377. Prodr. p. 155.

Icon. Jacqu. fl. Austr. t. 348.

Germ. Klettenartige Distel.

Gall. Chardon fausse - Bardane.

Hab. In Principatu Fürstenbergensi circa Engen in convallibus altioribus graminosis hu-

SYNGENESIA. POLYG. ÆQUALIS. 365

midis *Krügenthal* rarius. In *Juræ* Helvetiæ vallibus t. *Hallero*.

lor. Julio, Augusto. *Semina* Septembri maturæ. ♀

Radix ramosa, fibrosa, fusca. *Caulis* bicuspidatus et altior, erectus, crassiusculus, angulatus, simpliciusculus, aut ramosus. *Folia* alterna, per caulem decurrentia, *inferiora* infra medium innatifida, quandoque pinnata, *superiora* sessilia, ovata, acuminata, margine inæqualiter acute serrata, longe ciliata, superne viridia, labra, inferne subvillosa, incana. *Flores* caulem ramosque terminantes, 4-5-8 sessiles, in capitulum congesti, purpurei, mediocres. *Calycinae* squamæ imbricatæ, longe subulatæ, glabræ, ultra medium patentireflexæ, atropurpureæ, aut virides. *Semina* fusca, striata. *Pappus* apicillaris, sessilis. *Receptaculum* pilosum.

♂ f. Planta simillima individuis in Helvetiæ alpestribus lectis.

In horto botanico culta, mutationi non succubuit.

Ad rariores omnino Floræ cives pertinet.

†† *Folius sessilibus.*

231. CARDUUS *marianus* foliis amplexicaulibus hastato-pinnatifidis spinosis, calycibus aphyllis: spinis canaliculatis duplicato-spinosis. *Linn. Sp. pl.* 2. p. 1153.

Silybum nervis foliorum albis. Hall. stirp. helv. n. 181.

Carduus albis maculis notatus, vulgaris. Bauh. pin. 181.

Carduus Marianus. Tabern. p. 1078. *Icon.*

366 SYNGENESIA. POLYG. ÆQUALIS

Icon. Curtiz. fl. Lond. t. 148. — Regna bot. — Blackw. t. 79.

Germ. Mariendistel.

Gall. Chardon Marie. Chardon Notre-Dame.

Hab. In vineis, ad pagorum hortos et fimbriatissimè. Prope Ettlingen in vineis.

Flor. Junio, Julio. Semina Augusto matura.

Radix cylindrica, crassa, descendens, undique fibrosa. Caulis tripedalis, bicubitalis, erectus, crassiusculus, fistulosus, sulcatus, inferius villosus-tomentosulus, superne glabriusculus ramosus. Folia radicalia supra terram expansa grandia, speciosa, sinuato-pinnatifida, nitida, margine superne sæpissime venis albis reticulatis picta; Caulina minora, amplexicaulis patentia, suprema basi cauli adpressa, recurva. Flores caulem ramosque terminantes, solitarii speciosi, purpurei. Calycinæ squamæ variegatæ, inferiores rotundæ, spinis ciliatæ; intermedie patentès, acuminatæ, rigidæ, superne canaliculatæ, spina forti flavescente terminatæ, apice basin spinis ciliatæ; superiores et interiores lanceolatæ, inermes, apice purpureæ, marginibus fissis. Semina ovata, subangulata, fusco-nigricantia, nitida. Pappus capillaris, rigidus, sessilis, obliquus. Receptaculum pilosum.

Pharm. CARDUI MARIE Radix; Herba; Semina.

Radix et folia demulcentes et abstergentes virtutes habent. Semina emulsionibus refrigerantibus contra Pleuritidem immiscentur.

Flores apibus melliferi.

Semina aviculis accepta.

1232. *CARDUUS cyanoides* caule ramoso folioso, foliis caulinis alternis sessilibus

SYNGENESIA. POLYG. ÆQUALIS. 367

libus non decurrentibus profunde pinnatifidis, laciniis lineari-lanceolatis integris falcatis, supremis et rameis simplicissimis lineari lanceolatis falcatis integris, omnibus margine reflexis subtus niveis.

Carduus cyanoides β. *polyclonos*. Linn. *Sp. pl.* 2. p. 1152. an? (*folia non decurrentia in individuis palatinis*).

Carduus mollis. Pollich. *fl. pal.* n. 769. (*excluso Linnaei, Bauhini et Clusii Synonymo*).

Acarna cyanoides, *Flor. der Wetterau*. 3. p. 157.

Icon. Sprengel *fl. Halensis*. t. XI.

Per m. Kornblumenartige Distel.

Call. Chardon cyanoide.

Hab. Circa Mannheim in arenosis locis. Inter Ladenburg et Lorsch in sabulosa terra copiose, post Pollichium vidi.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ♀

Radix cylindrica, longa, fibrosa. *Caulis* sesquipedalis, cubitalis, erectus, superne sæpe flexuosus, obsolete angulosus, striatus, raro tomento obsessus, ramosus. *Rami* alterni flexuoso-erecti, foliosi, simpliciusculi. *Folia* radicalia plurima, *caulina* alterna, semiamplexicaulia, sessilia, profunde ad costam mediam fere pinnatifida: *laciniis* alternis, integriusculis lineari-lanceolatis falcatis, margine revolutis; *superiora* rameaque simplicissima, lineari-lanceolata, falcata, omnia superne viridia, subpubescentia, subtus niveo-tomentosa. *Flores* caulem ramosque terminantes, solitarii, erecti, majusculi,

368 SYNGENESIA. POLYG. ÆQUALIS.

purpurei. *Calycinæ squamæ* imbricatæ, lineari lanceolatæ, inermes, reflexiusculæ, patentæ. *Pappus* capillaris per lentem subplumofus tessilis. *Receptaculum* pilosum.

Differt a *Carduo molli* Linnaei cum quo Pollichius eum confundebat, Caule non simplici unifloro, ex flore ad partem caulis infimam non aphyllæ. Foliis radicalibus caulisque infimis multo majoribus, laciniis triplo longioribus falcatis, Calycum squamis non tomentosis, ovato-lanceolatis reflexis, dense imbricatis. Flore minore.

Differt a *Carduo cyanoide*. Willden. Linn. Sp. pl. 3. p. 1655. Caule non unifloro, seu Gmelini *cyanoide*. fl. Sibir. 2. p. 42. t. 15. foliorum segmentis angustioribus longioribus exacte falcatis, margine non denticulatis, Caulibus ramosioribus; a *Carduo polycle* nos. Willd. l. c. Gmel. fl. Sibir. 2. p. 44. f. 16. Caule minus ramoso, ramis simplicibus, unifloris, non multifloris, Floribus duplo majoribus.

Obs. Pollichii plantam etiam vidi in Franconia macilentis pascuis prope Würzburg.

In Horto culta, constantissima permanfit.

CCCLXXII. C N I C U S.



GENESIA. POLYG. ÆQUALIS. 369

t. Hic veniunt etiam Cardui Linn. pappo plumoso instructi f. Cirsium Tournef.

† *Folius decurrentibus.*

3. CNICUS palustris foliis decurrentibus, dentatis, margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Linn. Sp. pl. 2. p. 1151. (Sub. Carduo). Pollich. fl. pal. n. 766.*

irsium caule recto subnudo, alato, foliis semipinnatis, spinosis, calycibus molli-ter spinosis. *Hall. stirp. helv. n. 170.*

arduus palustris. *Bauh. pin. 377. — Prodr. p. 156. — Mapp. fl. Als. p. 54.*

n. *Curtis. fl. Lond.*

m. *Sumpf-Kratzkraut, Sumpf-Distel.*

l. *Cirse des marais.*

. Utrique in pratis humidis paludosis, in sylvaticis et pascuis montosis udis passim frequens.

r. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ♂

Caulis bicubitalis, tricubitalis et altior, eremultangulus, foliis decurrentibus utrinque s, spinosus, pilis albis hirsutus, sæpe viridi purpureo-variegatus, simpliciusculus aut raris. *Rami* alterni, alati, spinosi. *Folia* radi- petiolata, *caulina* alterna, sessilia, lanceo- acuta, decurrentia, pinnatifida, dentata, ata, venosa, margine spinosa, spinis basi urascentibus, apice albidis, superne obscure ia, hirsutula, subtus quammaxime hirsuta, ca; *suprema* angustiora, sæpe recurva *Flo-* caulem ramosque terminantes, congesti,

SYNGENESIA. POLYG. ROTALI

sessiles, purpurascentes. *Calyx* ventricosus, oblongus, sublanuginosus, *squamis* arcte imcatis, ovatis, viridibus, sæpe violaceo-purpurascentibus, lævibus, in mucronem brevem vergentem purpureum terminatis. *Semina* ovata, obsolete angulata, albida, nitida. *Stylus* sessilis, plumosus. *Receptaculum* pilosum. Variat Corollis albis.

Ubi in abundantia crescit, sæpe indicium subje-
tis Turfæ præbet. Maxima sæpe pratorum uliginosis
pestis.

Florēt apibus melliferi.

1234. CNICUS lanceolatus foliis

currentibus pinnatifidis hispidis: laci-
divaricatis, calycibus ovatis spinosis,
lofis, caule piloso. *Linn. Sp. pl.* 2
1149. (*sub Carduo*) — *Pollich. fl.* 1
n. 762.

Cirsium foliis semipinnatis, pinnis bilob-
spinosis, capitulis congestis, conicis,
mentolis. *Hall. stirp. helv. n.* 169.

Carduus lanceolatus latifolius. *Bauh.* 1
385. — *Mapp. fl. p. Als. p.* 53.

Carduus lanceolatus. *Tabernæmont. p.* 10
Icon bona.

Icon. Oeder. fl. Dan. t. 1173.

Germ. Lanzen-Kratzkraut. Sperrdistel.

Gall. Cirse lanceolé.

Hab. Utrunque ad sepes, vias, fossas, et
geres, nec non in ruderatis et viis pa-
rum neglectis passim abunde.

Flor. Julio, Augusto. *Semina* Septembri
tura. ☞

YNGENESIA. POLYG. ÆQUALIS. 371

Radix fusiformi-cylindrica, fusca, fibrosa. *Folia* cubitalis, tripedalis et altior, erectus, glabrus, villosus, ex foliis decurrentibus alatis spinosus, superne albide tomentosus, rarius. *Folia* alterna, quandoque semidecuria, sessilia, pinnatifida, laciniis lateralibus unde bi-trifidis, lineari-lanceolatis, divaricatis, apice spinosis, terminali lineari-lanceolata superne saturate viridibus, asperis, subtus leuotomentosulis. *Flores* caulem ramosque terminantes, erecti, subsolitarii, purpurei, majores. *Calyx* ovatus, squamis ovato-lanceolatis, rigidis, carinatis, late viridibus, apice mucrone pungente munitis, intimis inermibus. *Corollae* calycinae basi praesertim pilis tenuibus obsita. *Receptaculum* pilosum. *Pappus* sessilis, compressus.

Corollis albis.

Flores apibus mel praebent.

†† *Foliis non decurrentibus.*

5. *CNICUS eriophorus* foliis sessilibus bifariam pinnatifidis: laciniis alternis erectis, calycibus globosis, villosis. *Linn. Sp. pl.* 2. p. 1153. — *Pollich. fl. pal. n.* 767. (*sub Carduo.*)

Cardium foliis pinnatis, pinnis bilobis, lanceolatis, spinosis, calycibus tomentosis, sphaericis, spinosis. *Hall. stirp. helv. n.* 168.

Carduus capite rotundo tomentoso. *Bauh. pin.* 382.

Carduus tomentosus, corona fratrum herbariorum. *Lobel. Icon.* 2. p. 9.

n. Jacqu. fl. Austr. 2. t. 171.

372 SYNGENESIA. POLYG. EQU

*Germ. Grofsköpfiges Wollkratzkra
lendistel.*

Gall. Cirse laineux.

Hab. In collibus argillaceo - calcareo
retro Grözingen et Pforzheim.
prope Niefern. In Ditione Basileen
te Vogelberg, Ramstein, Dietisber
serfallen etc. In Principatu Fürste
prope Donaueschingen et alibi a
blicas frequens. In Palatinatu circa
heim ad sepēs; retro Frankenth
gum Edchum, prope Alzey post P
vidi.

Flor. Julio, Augusto, Septembri. Se
gusto, Octobri matura. ♂

Caulis bicubitalis, tricubitalis, er
gulofus, molliter piloso - tomentosus,
Ram? alterni, divisi. *Folia* alterna, fesi
decurrentia, pinnatifida; *laciniis* pro
fidis, longis, divaricatis, fursum
que directis, basi et apice longe spi
perne læte viridibus, villosiusculis,
inferne albide tomentosus, mollibus
caulem ramosque terminantes, soliti
sessiles, magni, purpurascens, rai
di. *Calyces* speciosi, magni, subrotun
mae lanceolato - lineares, apice spino
obtectæ et nidulantes in tomento lax
Pappus sessilis plumosus. *Receptaculum*

Planta omnino perpulchra. Flores apibu
bent.

Amat præsertim calcareo - argilloso, aspi

1236. *CNICUS anglicus* foliis f
semialexicaulibus, lanceolat
dentatis ciliatis, subtus incanis,

NGENESIA. POLYG. ÆQUALIS. 373

nuginosis, caule supra medium aphylla,
lanuginoso subflexuoso unifloro.

anicus pratensis foliis sessilibus lanceolatis,
subdentatis, ciliatis, subtus lanatis, caule
nudiusculo unifloro. *Smith. fl. brit. 2. p.*
854. — Huds. fl. Angl. 2. p. 353. (excluso
Jacquini et Clusii Synonymo).

irsum anglicum Lamarch et Decandolle fl.
franc. 4. p. 118.

arduus heterophyllus. Relhan. fl. Cantabr.
p. 306. n. 591. — Ligtfoot. fl. Scot. 1. p.
456. n. 8.

irsum anglicum radice Hellebori nigri mo-
do fibrosa, folio longo. Bauh. hist. 3. p.
45. Icon.

z. Petiver. herb. brit. t. 22. f. 1. — Lohel.
Icon. 583.

m. Englisches Kratzkraut.

l. Cirse d'Angleterre.

. In pratis montosis humidis apricis Comi-
tatus *Nellenburgensis* inter *Stokach* et *Ziz-*
zenhausen ad dextram nuper mihi obvi-
us. Sine dubio etiam in Principatu superiore
in pratis montosis ad Lacum Bodamicum,
nec non in Principatu Fürstenbergenfi.

r. Junio, Julio. Semina Julio, Augusto ma-
tura. 7

Radix fibrosa, repens, fusca. *Caulis* peda-
bipedalis, erectus, simplex, solitarius, a
ad medium fere foliosus, a medio ad api-
aphyllus, albide laxè lanuginosus, subfle-
us, ultra medium squama sæpe foliacea
alterave præditus. *Folia* radicalia in petiolum
irrentia, lanceolata, versus basin angustata,
gine minutim subsinuato-denticulata, spinu-

374 SYNGENESIA. POLYG. ÆQUALIS.

loso longius ciliata, 4-5 uncias longa, semiunciam ad septem lineas lata; *caulina* alterna, sessilia, semiamplexicaulia, lanceolata, erecta, integriuscula, margine spinuloso ciliata, omnia supra pallide viridia, hispidiuscula, subtus ex albido-glaucolaxe lanuginosa. *Flos* terminalis, unicus, erectus, purpureus, mediocris. *Calyx* ovatus, subtomentosus. *Squamae* imbricatæ ovato-lanceolatæ, acutæ, erectæ, mites, ultra medietatem violaceo-purpurascens. *Pappus* plumosus, sessilis. *Receptaculum* villosum.

Variat. Caulibus demissioribus digitalibus. Foliis profunde sinuato-dentatis. Cæterum in omnibus similis plantæ descriptæ.

Minime confundendus cum *Carduo pannonico* primo intuitu affini, qui a nostro *Cnico* distinguitur Caula tenuiori, laxissime tomentoso, Foliis multo brevioribus, læte viridibus, radicalibus saltem subtus non glauco-albide tomentosis, Flore duplo minore, Pappo capillari, non plumoso.

Distinctus omnino a *Cnico heterophyllo* et *Helenioides*, Caule semper fere brevior, non ad florem folioso. Foliis caulinis non cordato-lanceolatis, sed latioribus, non serratis, serrulatis. Foliis

SYNGENESIA. POLYG. ÆQUALIS. 375

mucronulato spinulosus, subtus tomentosus, infimis laciniatis, caule unifloro.

Cnicus heterophyllus foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque, caule subunifloro, calyce inermi. *Linn. Sp. pl.* 2. p. 1154. (*sub Carduo*).

Cirsium foliis ciliatis, subtus tomentosus, radicalibus petiolatis, ovato-lanceolatis, caulinis lanceolatis aut semipinnatis. *Hall. Stirp. helv.* n. 180.

Cirsium heterophyllum. *Allion. fl. Pedem.* n. 554. t. 34.

non. Oeder. fl. Dan. t. 109.

erm. Verschiedenblättriges Kratzkraut.

all. Cirse variable.

a b. In pratis montosis humidis Principatus superioris cum *Pinguicula vulgari* passim rarius.

lor. Julio, Augusto. *Semina* Septembri maturatione. ☞

Caulis cubitalis, tripedalis, erectus, simplex, angulatus, striatus, laxè incano-lanuginosus, a radice ad apicem fere foliosus, unibrachius. **Folia** radicalia petiolata, lanceolata, cuneata, serrulato-spinulosa; **caulina** alterna, sessilia, late amplexicaulia, cordato-lanceolata, superiora laciniata: *lacinae* 1-2-sæpe 4 ad 5, rectae, subparallelæ, subsinuatae, omnia margine argute spinulosa, superne viridia, inferne ivo tomentosa. **Flos** saltem in speciminibus ossis solitarius, terminalis, erectus, purpureus, majusculus. **Calycinae** squamæ ovato lanceolatae, erectae, mites, glabrae, ex viridi-fusco-virpurascentes. **Pappus** sessilis, plumosus. **Receptaculum** villosum.

376 SYNGENESIA. POLYG. ~~282122~~

Obs. Variationem sic dictam *Helensioide* omnibus ciliatis integris, hucusque in regibus citatis non vidi.

Planta cum *C. Helensioide* in Helvetia lecto horto botanico culta, mutationi non succubuit.

1238. *CNICUS rivularis* foliis in pinnatifidis, laciniis oblongis super integris ferratis inermi spinulosis lobis, caule inermi, floribus cap Murr. Linn. Syst. veg. ed. 14. p. (sub *Carduo*).

Cirsium tricephalodes. Lamarck et D. d. fl. franc. 4. p. 146.

Icon. Jacqu. fl. Austr. 1. t. 91.

Germ. Bachkratzkraut.

Gall. Cîrsc a trois têtes.

Hab. In Margraviatu superiore circa the prope *Müllheim* in pratis riguis, nec prope *Zunzingen* in pratis udis t. V In Principatu superiore in pratis ud Lacum *Bodamicum* passim non infreq In Principatu *Fürstenbergen*si, prope *naueschingen in der Baar* in pratis e cuis humidis, nec non retro *Fürsten* nuper legi.

Flor. Majo, Junio. *Semina* Julio matur

Radix cylindrica, crassiuscula, horis lis, fibris longis aucta, albida. *Caulis* cub tripedalis, solitarius, erectus, inferne su angulatus, supra medium fere foliosus, su teretiusculus, striatus, aphyllus. squama cea una alterave præditus, albide glaucoginosus, simpliciusculus. *Folia* infima in um dentatum attenuata, reliqua caulina ali

SYNGENESIA. POLYG. ÆQUALIS. 377

filia, amplexicaulia, non decurrentia, omnia pinnatifida, plus minusque profunde incisa; aciniac subparallelæ, lanceolatæ, spinuloso-iliatæ, extima major acuminata, omnia saturate viridia, subtus pallidiora, subvillosula. Flores caulem terminantes, sæpe tres, sessiles, caliculus constituentēs, purpurei, medio superiore majore, quandoque 4-5-7 infimis tunc bipedunculatis, perraro unicus tunc major. Calyx ovatus glutinosus. Squamæ imbricatæ, lanceolato-acuminatæ, mites, fusco-purpurascentes. Pappus plumosus, sessilis. Receptaculum pilosum.

Nota. Ad rariores Floræ cives pertinet, a Germaniæ Floristis hucusque omissa planta.

1239. ENICUS tuberosus foliis petiolatis subdecurrentibus subpinnatifidis spinosis, caule inermi, floribus solitariis. *Linn. Sp. pl. 2. p. 1154. — Pollich. fl. pal. n. 768. (sub Carduo).*

Cirsium radicibus bulbosis, foliis semipinnatis, pinnis angulosis, ciliatis, floribus nudis. *Hall. stirp. helv. n. 177.*

Carduus pratensis Asphodeli radice, latifolius. *Bauh. pin. 377. — Moris. hist. 3. Sect. 7. t. 29. f. 27. 28.*

F. Foliis magis dissectis.

Cirsium dissectum. *Lamarck. fl. franc. 2. p. 27.*

Cirsium pratense Asphodeli radice foliis tenuiter incis. *Mapp. fl. Alf. p. 76.*

Jacea quarta. *Tabernæmont. p. 436. Icon.* Caulis ibi nimis ramosus,

378 SYNGENESIA. POLYG. ÆQUAL

Germ. Knollentrageses Kratzkraut.

Gall. Cirse bulbeux.

Hab. Circa Carlsruhe in pratis aquosis Neureuth et Eckenstein frequens, cum oleraceo. In Alsatia Michelfeldae in ad fossas, et prope Neudorf t. Lache prope Argentinam in der Ruprechtsdem Riet inter fluvios Ellam et Rh. In Principatu superiore ad Lacum Bo cum in pratis. In Palatinatu prope derulm in pratis; in montis Donner pascuis gramineis: in pratis ad salinas Dürkheim t. Pollichio.

Flor. Julio, Augusto. Semina Augusto, tembris matura. ¶

Radix e medio trunco multiceps, l plurimis fusiformibus prædita. Caulis cub tripedalis et altior, erectus, anguloso-sul scabriusculus, simpliciusculus, uniflorus flore albide tomentosus, rarius ramo u alterove superne præditus. Folia alterna plexicaulia, basi quandoque subdecurra pinnatifida, pinnis bitrifidis plus mini profundis, margine spinuloso-mucronatis, nia utrinque villosulo-scabriuscula, læte dia. Flores erecti, aut subnutantes pur quandoque ochroleuci, et dilute rosei. subrotundus. Squamæ calycinae lanceol mites, apice sæpe subreflexæ. Pappus se plumosus.

Nota. Variatio Floribus ochroleucis in caule 1-2-3 pedunculatis instructa, minime co genda cum Cnico Erysiethales. Jacq Austr. t. 310. qui a nostra planta differt C crassiore, altiore, Foliis multo majoribus ad mediam costam pinnatifidis, Floribus per nutantibus. Calycum squamis multo vioribus ultra medium incurvis.

SYNGENESIA, POLYG. ÆQUALIS. 379

240. CNICUS *acaulis* subacaulis, uniflorus, foliis lanceolatis pinnatifidis, plicato-undulatis, ciliato-spinosis, squamis calycinis erectis, mucronulatis glabris.

Carduus *acaulis* calyce glabro. *Linn. Sp. pl.* 2. p. 1156. — *Pollich. fl. pal. n.* 770.

Cirsium floribus acaulis. *Hall. stirp. helv. n.* 178.

Cirsium acaulon flore purpureo. *Mapp. fl. Alf. p.* 76.

Chamæleon exiguus Tragi. *Bauh. hist.* 3. p. 62. *Icon.*

Caule palmari, subpedali, folioso unifloro.

Carduus *dubius* *Willden. prodr. fl. Berol. n.* 801. t. 6. f. 11.

on. Jacqu. Icon. rar. — *Oeder. fl. Dan.* t. 1114.

erm. Stengelloses Krätzkraut. Niedere Distel.

ill. Cirse nain,

ib. Utrunque passim. In pascuis graminosis, aridis sabulosis sterilibus. Inter Mühlburg et Dachsland. Prope Kehl, et præsertim in arenosis pascuis Rheno vicinis. Prope Argentinam in der Ruprechtsau et alibi.

lor. Julio, Augusto. Semina Augusto, Septembri matura. ¶

Radix fusiformi-cylindrica, fibris aucta, ca. **Caulis** brevissimus, 1-2-3 uncialis, interm palmaris et longior, uniflorus, aphyllus, et foliosus, piloso-scabriusculus. **Folia** radialis plurima, lanceolata, pinnatifida, subundulata, plicata, utrinque læte viridia, margine mucoso-aculeata, rigidiuscula. **Flos** magnus, purpureus, raro albus. **Calycis** ovati squamæ

imbricatæ, ovatæ, glabræ, mucrone parvulo munitæ, erectæ. *Pappus* sessilis, longus, plumosus. *Receptaculum* pilosum.

Qb f. Planta in sabulosis graminosis aridis apricis macilentis sæpissime subcaulis. In Horto culta semper caulifera, caule semipedali et altiori instructa, quandoque ramosa evadit.

Flores speciosi apibus melliferi.

1241. *CNICUS Lachenalii* foliis amplexicaulibus, pinnatifidis, glabris, laciniiis inciso-bifidis dentatis, margine et apice præsertim longe spinulosis, caule ramoso, calycibus oblongis, bracteatis.

Cnicus rigens foliis oblongo-lanceolatis glabris, margine spinosis pinnatifidis, laciniiis oblique lobatis, calycibus oblongis bracteatis. *Aiton. Hort. Kew. 3. p. 144.*

Cirsium foliis ciliatis semipinnatis, pinnis angulosis spinosis, superioribus amplexicaulibus. *Hall. stirp. helv. n. 176. (excluso Linnaei Synonymo).*

Icon. Lachenal. Act. Helv. IV. t. XVI. p. 294.

Germ. Lachenälisches Kratzkraut.

Gall. Cirse de Mr. Lachenal.

Hab. In agro Basileensi in pascuis macilentis ad *Birsam*, trans fluentum inter pontem majorem et locum *das Schänzlein* dictum, unico loco juxta dumeta non copiose legitur. *Lachenalius cum Cl. Miegie.*

Flor. Julio, Augusto, *Semina* Augusto, Septembri matura. ♀

Radix transversa, nodosa, multiceps, fibras deorsum demittens longas, teretes, albicantes. Ex quovis radicis capite *Caulis* pedalis, cubi-

lalis, erectus, crassiusculus, profunde striatus, hirsutus, ramosus, inferne purpurascens. *Folia* radicalia plurima, pinnatifida, basi angustata, pinnatifida, profunde serrata, superne saturate viridia, raris pilis adspersa, subtus pallidiora, margine spinoso-ciliata; *Caulina* alterna, inferiora radicalibus similia, latiora, sæpe tamen et minus longa; superiora minora profunde laciniata: *laciniis* integris vel semibilobis. *Rami* supremi foliis vestiuntur parvis, lineari-lanceolatis, sublinuatis, integrisve, margine ciliato-spinosis. *Flores* caulem ramosque terminantes, subsolitarii, ex viridi-albicantes, duobus, sæpissime tribus rarius quatuor foliolis linearibus ciliato-spinulosis obvallati. *Calyx* ovali-cylindricus. *Squamæ calycinae* imbricatae, lanceolatae, acutæ, erectæ, glabræ, virides, in spiculam mitem desinentes. *Semina* ovalia, compressa. *Pappus* sessilis, plumosus, longus. *Receptaculum* pilosum.

Cl. Lachenalius et Hallerus confundebant plantam cum Carduo tartarico Linnaei a quo autem quammaxime recedit, ut facile ex Jacquini fl. Austr. t. 90. patet, ubi C. tartarici Icon prostat, ex qua mihi saltem videtur C. tataricum vix specie diversum esse a Cnico oleraceo. Linn.

Minime pro variatione C. oleracei Linn. habenda, a quo differt, Habitu toto, Caulibus demissioribus, ramosis. Foliis longe aliis, magis margine ciliato-spinulosis. Foliolis calyces circumdantibus, multo minoribus, angustioribus, linearibus, paucioribus.

)b f. Planta in horto Botanico Carlsruhano ab amicissimo Lachenalio mihi missa per quindecim annos culta, mutationi non succubuit.

1242. *CNICUS oleraceus* foliis pinnatifidis carinatis nudis, bracteis subcoloratis

SYNGENESIA. POLYG. EQUALIS

integris concavis. *Linn. Sp. pl. 2. p. 11.*
Pollich. fl. pal. n. 771.

Cirsium foliis ciliatis amplexicaulibus, floribus
 comprehendentibus. *Hall. stirp. helv.*
 173.

Cnicus pratensis, Acanthi folio, flore flavo
 cente. *Mapp. fl. Alf. p. 78.*

Carduus pratensis latifolius. *Bauh. pin. 3.*
Feon. Oeder. fl. Dan. t. 86a. — Schkuhr. b.
handb. t. 229.

Germ. Kohlratzkraut. Wiesendistel.

Fr. a ll. Cirse des près. Quenouille des près.

Hab. Utrunque passim in pratis humidis, usque
 ad sylvarum marginalia uda gramina
 gultosa, copiose.

Flor. Julio, Augusto. *Semina* Augusto, Septem-
 brembri matura. 7

Caulis tripedalis, bicubitalis, erectus, striatus,
 glabriusculus, crassiusculus, simpliciusculus
Folia alterna, sessia, amplexicaulia, pinnatifida,
 venosa, glabriuscula, laciniis ovali-lanceolatis,
 serrulatis, margine spinuloso-ciliatis
Flores tres, sæpe breviter pedunculati, pallide
 flavicantes, basi *Bracteis* foliaceis ovalibus, aut
 ovali lanceolatis, pallide luteis, margine ciliato
 spinulosis, obvallati. *Calyx* ovali-cylindricus
 squamis imbricatis, longe lanceolatis, erectis
 glabris, pallide viridibus, apice mucronulatis
Pappus plumosus. *Receptaculum* pilosum.

Variat. Foliis majoribus et minoribus, omnibus
 integriusculis lanceolatis, et ovatis margine denti-
 culatis, ciliato-spinulosis, Calycibus plus minusve,
 quandoque vix bracteatis.

Tota planta sæpe ex viridi pallide flavescent.

Ubi abundanter crescit sæpe indicium Turfæ sub-
 jacentis.

CCCLXXIII. ONOPORDON.

Linn. Gen. pl. ed. Schreb. n. 1256.

Calyx communis subrotundus, ventricosus, imbricatus, magnus, *Squamis* numerosis, spinosis, undique patentibus. *Corolla* composita, tubulosa, uniformis. *Corollulae* hermaphroditae, æquales. *Propria* monopetala, infundibuliformis, tubo tenuissimo, limbo erecto, ventricoso, quinquefido; *laciniis* æqualibus, unica profundius separata. *Calyx* fructiferus leviter connivens. *Semina* ovali-quadrangularia. *Papillae* capillaris, sessilis. *Receptaculum* paleaceum: *alveis* in alveolos coadunatis, truncatis, mucronatis, seminibus brevioribus.

1243. *ONOPORDON Acanthium* calycibus squarrosis: spinis subulatis, foliis ovato-oblongis sinuatis. *Linn. Sp. pl. 2. p. 1158.* — *Pollich. fl. pal. n. 772.*

Onopordon caule alato, foliis ovatis dentatis: dentibus angulosis, aristatis. *Hall. stirp. helv. n. 159.*

Acanos Spina. Scopol. fl. Carn. 2. n. 1013.

Carduus tomentosus, Acanthisfolio, vulgaris. Bauh. pin. 302. — Mapp. fl. Alf. p. 55.

Acanthium vulgare flore albo et purpureo. *Tabernemont. p. 1063. Icones.*

Icon. Curtis. fl. Lond. t. 334. — Oeder. fl. Dan. t. 909. — Schkuhr. bot. handb. t. 230.

Germ. Krebsdistel. Frauendistel. Eselsdistel.

Call. Grand Chardon aux ânes.

INGENESIA. POLYG. ÆQUALI

hab. Utrunque præsertim ad urbium et pag
ruerata, sæpes, vias, passim frequer

Flor. Julio, Augusto, Septembri. *Semina*
tembri, Octobri matura. ♂

Radix fusiformis, fibrosa, extus flavi
intus alba, longa, crassiuscula. *Caulis* tri
lis, tricubitalis et altior, erectus, totus al
alis latis spinosis, ad basin usque ramosus. *l*
alterna, sessilia, decurrentia, ovata, acuta, f
to-dentata, inferiora amplissima, margine
nia spinosa, utrinque lanugine plus minus
cana. *Flores* caulem ramosque termina
erecti, solitarii, magni, purpurei seu albi
lycinæ squamæ numerosæ, spinosæ, prom
tes, basi pilis albis intertextæ. *Spinæ* apice li
Receptaculum cellulis membranaceis tetrag
reticulatum.

Pharm. CARDUI TOMENTOSI f. ONO *DI Herba recens.*

Succus foliorum expressus contra cancrum jam
summo cum effectum abhibebatur hinc Krebsdi
audit.

Nunc iterum a recentioribus, cum primis in C
nomate faciei, narium, et labiorum summo sæpe
fructu coeptus est adhiberi.

Radice decoctum These instar sumptum, sp
cum est in Gonorrhoea incipiente. Scop. l. c.

Flores apibus ceræ mellisque materiam præbe

Discus ejusdem instar Cynaræ edulis. L
Amen. Acad. 6. p. 124.

Oleum ex seminibus pro lampadibus, ponder
elicitur. Durande in Bibl. phys-oecon. A
2. p. 121.

Semina avibus per hyemen sæpe summum solat

CCCLXX

CCCLXXIV. CARLINA.

Linn. Gen. pl. ed. Schreb. n. 1258.

Calyx communis ventricosus, radiatus, imbricatus: *Squamis* numerosis, laxis, acutis, quarum *interiores* in orbem positæ, longissimæ, patentes, nitidæ, coloratæ, radiantes. *Corolla* composita, uniformis, tubulosa: *Corollulæ* hermaphroditæ, æquales. *Propria* monopetala, fundibuliformis: *tubo* tenui; *Limbo* infundibuliformi, quinquesido. *Calyx* fructiferus imbricatus. *Semina* teretiuscula. *Pappus* divisus radios subpatentes, plumosos. *Receptaculum* communium, paleaceo-fetaceum: *paleis* membraceis, basi subcoadunatis in alveolos, multis, radiis subulatis; *setis* paullo longioribus, levatis, paleis intermixtis.

244. *CARLINA acaulis* caule unifloro, flore brevior. *Linn Sp. pl. 2. p. 1160.*

Carlina caule unifloro. *Hall. stirp. helv. n. 183.*

Carlina acaulos magno flore albo. *Bauh. pin. 380. — Mapp. fl. Als. p. 57.*

Chamæleon albus sive *Carlina*. *Tabernæm. p. 1068. Icon.*

2. *Carlina* caulescens magno flore rubente. *Bauh. pin. 380. — Mapp. fl. Als. p. 57.*

Carlina caulescens. *Tabernæm. p. 1068. Icon.*

Leon. Schkuhr. bot. Handb. t. 222. — Blackw. t. 532.

German. Eberwurz. Rosswurz.

Flora Bad. Tom. III.

B b

Gall. Carline a courte tige. Artichaud sauvage.

Ha b. In montosis graminosis apricis, juniperitis siccis passim. In Marggraviatu superiori prope *Badenweiler auf dem Lindengrabe* retro *Sirnitz prope dem Nonnattweyhe* nec non in monte *Blauen, Bölchen et Felberg* cum β . In Alsatia vogelis retro *Baar* prope *Marienkirch*, prope *Kaysersberg* et alibi passim. In Principatu superiore nec non in Principatu Fürstienbergenensi passim non infrequens.

Flor. Julio, Augusto. *Semina* Septembri maturantur. ζ

Radix cylindrica, longa, apice ramosa, rugosa, tuberculata, extus fusca. *Caulis* sere nullus, sæpe uncialis, digitalis, quandoque semipedalis et altior, erectus, subuniflorus. *Folia* radicalia terræ incumbentia, plurima, sessilia, in orbem digesta, lanceolata, pinnatifida: *laciniis* bi-tridentatis, spinula terminatis, glaberrima. *Flos* terminalis, speciosus, maximus. *Calyx* foliola exteriora pinnatifida, margine spinis palmatis obsessa; interiora multo longiora, lineari-lanceolata, alba, radiantia, in sole pulcherrime expansa, alba, nitentia, arida, corollam polypetalam imitantia.

Pharm. CARLINÆ Radix.

Radix amaricans, fragrans, aromatica. olim magicis et veterinariis magis quam hodie medicis nota. Contra pestem depredicata, qua ipsa Carolus magnus pestem ab exercitu suo fugasse, dicitur, unde forsitan, *Carlinæ* nomen ortum. Mathiol in Dioscor. ed. Bauh. p. 491.

Disci carnosissimi florum, calyce corollisque resectis instar capitum *Cynaræ Scolymi* in usus culinarios cedunt, eosque in Pyrenæis cum oleo olivarum commisit inter fercula sapidissima oblatos acceptos que habue-

SYNGENESIA. POLYG. ÆQUALIS. 387

245. *CARLINA vulgaris* caule multifloro, corymboſo, floribus terminalibus, calycibus radio albis. *Linn. Sp. pl.* 2. p. 1161. — *Pollich. fl. pal.* n. 773.

Carlina foliis imis ſemipinnatis, pinnis anguloſis, ſpinolis, ſuperioribus ovato lanceolatis, amplexicaulibus, calycinis herbaceis, ciliatis. *Hall. ſtirp. helv.* n. 182.

Carlina ſylveſtris vulgaris. *Mapp. fl. Alſ.* p. 57. *Heracantha.* *Tabernæmont.* p. 1076. *Icon.*

o n. *Oeder. fl. Dan.* t. 1174, — *Schkuhr. bot. handb.* t. 232.

erm. *Sodkraut.* *Sanddiſtel.* *Sand-Carlina.*

all. *Carlina vulgaire.* *Chardonnette commune.*

ab. Utrinque paſſim ad vias, colles aridos, ericeta, juniperita, aſpera, aprica, frequens.

lor. Auguſto, Septembri. *Semina* Septembri, Octobri matura. ♂

Caulis ſesquipedalis, cubitalis, erectus, res, albide tomentofus, ramoſus. *Rami* almi, erecti. *Folia* alterna, ſeſſilia, amplexiulia, ſinuato-denticulata, margine ſpinofa, perne læte viridia, inferne tomento incano ſeſſa. *Flores* caulem ramoſque terminantes. *rollulae* pallide luteae. *Calycinae* ſquamae anteriores lineares, patentes, radiantes, ſirineae.

CCCLXXV. B I D E N S.

Linn. Gen. pl. ed Schreb. n. 1267.

Calyx communis imbricatus, erectus: *foliis* ſaepe aequalibus, oblongis, canaliculatis

388 SYNGENESIA. POLYG. ~~ÆQUALI~~

concavis. *Corolla* composita uniformis, 1
lofa. *Corollulae* hermaphroditæ, tubulosæ.
pria monopetala infundibuliformis: *limbo* c
quefido, erecto. *Calyx* fructiferus immut
Semina obtusa, angulata, *denticulis* (pa
binis aut pluribus aristatis, rectis, acutis,
trorsum scabro hamosis, superne coronata
ceptaculum paleaceum, planum: *Paleis* c
duis, planiusculis.

Obs. *Corolla* interdum flosculo uno
terove radiato instructa.

1246. *BIDENS tripartita* foliis tri
calycibus subsoliosis, seminibus ere
Linn. Sp. pl. 2. p. 1165. — *Poll. fl.*
n. 774.

Bidens foliis petiolatis trilobatis et quin
lobatis, ferratis, floribus circumval
Hall. stirp. helv. n. 121.

Bidens foliis tripartitis divisis. *Mapp. fl.*
p. 39.

Cannabina aquatica folio tripartitim di
Bauh. pin. 321.

Icon. Curtis. fl. Lond. t. 237.

Ger m. Wasserhanfkrout.

Gall. Bident chanarin.

Hab. Utrinquè ad fossas aquaticas, stagna
ludes, inundata abunde.

Flor. Julio, Augusto, Septembri. *Semina*
tembri, Octobri matura. ☉

Caulis palmaris, pedalis, tripedalis c
tior, erectus, fulcatus, teretiusculus, gl
usculus, sæpissime rubellus, ramosus. F

SYNGENESIA. POLYG. ÆQUALIS. 389

ppositi. *Folia caulina* opposita, petiolata, basi
 connata, tripartita, sæpe etiam quinquepartita,
 sciniis ferratis; *suprema* simplicia, lanceolata,
 serrata, omnia glabriuscula, pilis hinc ciliata.
Floras caulem ramosque terminantes, lutei,
 subnutantes. *Calycina* foliola plura, integrius-
 cula, lanceolata, ciliata, flosculos involucran-
 tia. *Squamæ* calycinæ ovato lanceolatæ, mar-
 gine luteolæ, lineis pluribus nigricantibus,
 parallelis pictæ.

Flavo colore tingit,

1247. *BIDENS cernua* foliis lanceolatis
 amplexicaulibus, floribus cernuis, semi-
 nibus erectis. *Linn. Sp. pl.* 2. p. 1165. —
Pollich. fl. pal. n. 776.

Bidens foliis sessilibus, ferratis, floribus nu-
 tantibus, circumvallatis. *Hall. stirp. helv.*
n. 120.

Bidens folio non dissecto. *Mapp. fl. Alf. p.* 39.

Cannabina aquatica folio non diviso. *Bauh.*
pin. 321.

8. *Bidens minima* foliis lanceolatis, sessilibus,
 floribus seminibusque erectis. *Linn. Sp.*
pl. 2. p. 1165. — *Pollich. fl. pal. n.* 775.
 — *Roth. fl. germ.* 3. p. 301. *Huds. fl.*
angl. p. 310.

Verbesina minima. *Dill. giff.* 167. *app.* 66.
Raj. angl. 3. p. 188. *t.* 7. *f.* 2.

γ. *Flore* aureis in radio ligulis prædito. Co-
 reopsis *Bidens.* *Linn. Sp. pl.* 2. p. 1281.

Chrysanthemum aquaticum, *Bidens* folio
 non diviso, flore aureis petalis undique
 radiato. *Morif. hist.* 3. p. 17. *Sect.* 6.
t. 5. *f.* 26.

390 SYNGENESIA. POLYG. ÆQUALIS.

Herba Cunigundis II. *Tabernæmont. p.* 336.
Icon bona.

Icon. Curtis. fl. Lond. t. 192. — *Oeder. fl. Dan. t.* 341. — *Schkuhr. bot. handb. t.* 235.
— β . *Oeder. fl. Dan. t.* 312. — *Curtis. fl. Lond. t.* 192. n. 8.

Germ. Ueberhängender Zweyzahn.

Gall. Bident penchée

Hab. a. et γ . utrinque in fossis aquaticis, inundatis, paludibus, ad stagna cum praeecedente abunde. β . In paludibus turfosis *auf dem Kaltenbrunn*; in Principatu superiore prope *Constanz* et alibi in turfosis. In *Alsatia* ad stagnum prope *Hagenau*. In Palatinatu prope *Lautern* in paludibus turfosis. *t. Pollichio.*

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ☉

Caulis pedalis, sesquipedalis et altior, erectus, superne striato-sulcatus, hispidulus, saepe rubellus, rubro punctatus, ramosus. *Rami* oppositi, erecti. *Folia* caulina opposita, basi subconnata, ovali lanceolata, ferrata, glabra, patentia. *Flores* caulem ramosque terminantes, solitarii, lutei, saepissime in radio ligulati, cernui, majusculi. *Calycina* foliola lineari-lanceolata, nervosa, ferrulata, reflexa, corolla longiora.

Variatio. β . minima, uncialis, biuncialis, digitalis, uniflora, flore minore, erecto, iustructa.

Flavo colore tingit.

Eadem ac in *Acmella zeylanica* vires in *B. cernua* statuebat *Cl. Nebelius*, et *Blankerhorn* ejus discipulus in peculiari dissertatione.

CCCLXXVI. C A C A L I A.

Linn. Gen. plant. ed. Schreb. n. 1269.

Calyx communis simplex, cylindricus: *squamis* tribus, quatuor ad quinque aequalibus, linearibus, tubum constituentibus. *Corolla* composita, tubulosa: *Corollulae* hermaphroditae numero foliorum calycinorum, uniformes. *Properia* monopetala, infundibuliformis, in tubum sensim attenuata: *limbo* quadrifido, patenti. *Filamenta* quatuor (nec quinque) *Antherae* apice quadridentatae. *Calyx* fructifer immutatus. *Semina* oblonga, glabra. *Pappus* sessilis, capillaris. *Receptaculum* planum, nudum.

1248. *CACALIA alpina* foliis reniformi-cordatis dentatis, utrinque glabriusculis, calycibus subtrifloris.

Cacalia alpina β. Linn. Sp. pl. 2. p. 1171.

Cacalia glabra. Villars. fl. Delph. 3. p. 170.

Cacalia alliariaefolia. Lamarch. dict. 1. p. 532.

Tussilago cacalia. Scopol. fl. Carn. n. 1055.

Cacalia foliis cordatis, serratis, longissime petiolatis, floribus umbellatis. Hall. stirp. helv. n. 137. d.

Cacalia foliis cutaneis acutioribus et glabris.

Bauh. pin. 198. — Moris. hist. 3. p. 94. Sect. 7. t. 12. f. 6.

Cacalia glabro folio. Clus. hist. 2. p. 115.

con. Jacqu. fl. Austr. t. 234.

Germ. Glatte Alpen-Cacalie.

392 SYNGENESIA. POLYG. ÆQUALIS.

Gall. Cacalia des alpes.

Hab. In montosis sylvaticis humidis prope *Reichenbach, auf dem Kaltenbrunnen*, in monte *Kniebis*. In Margraviatu superiore *auf dem Blauen, Stokberg, Belchen*, circa *Sirnitz et Marzell*, in monte *Feldberg* Brisgoviae. In Ditione Basileensi in monte *Vogelberg et Wasserfallen*. In Alsatiae Vogesis in monte *St. Ottiliae*; retro *Rappoltsweyher*, prope *Dusenbach*, in monte *Baldon* et alibi.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura, ☞

Radix ramosa, plerumque praemorfa. *Caulis* cubitalis, tripedalis et altior, erectus, crassiusculus, angulatus, striatus, superne subpubeles, umbellato-paniculatus. *Folia* radicalia longe petiolata, reniformi-subrotunda, sinuato-dentata, subtus venis reticulata, ampla, praemagna. *Caulina* alterna, pauca, minora, radicalibus similia, acutiuscula, omnia utrinque glabriuscula, laete viridia. *Flores* subumbellato-paniculati, copiosi, pallide purpurei quandoque albidi, parvuli. *Calycinac* squamae 3-4 lineares, erectae, obtusae, glabrae, conniventes, ex viridi-purpureae. *Flosculi* in singula flore 3-4 calyce triplo fere longiores. *Filamenta* quatuor. *Antherae* atropurpureae, apice quadridentatae.

Planta elegans in horto botanico ultra viginti annos culta, mutationi non succubuit.

Flores melliferi.

1249. *CACALIA albifrons* herbacea, foliis petiolatis, dentatis, subtus incanis petiolis basi auriculatis, calycibus quadrifloris.

SYNGENESIA. POLYG. ÆQUALIS. 393

Cacalia albifrons. Linn. *Suppl.* 353.

Cacalia Petasites. Lamarck et Decandolle
fl. franc. 4. p. 127.

Cacalia hirsuta foliis reniformi-hastatis duplicato-ferratis, hirsutis, caulinis stipula dilatata amplexicaulibus, calycibus tri vel sexfloris, floribus racemosis. Villars.
fl. Delph. 3. p. 172.

Cacalia tomentosa. Jacqu. *fl. Austr.* 3. p. 20.

Cacalia Alliariae calycibus trifloris, foliis cordatis petiolatis, dentato-angulatis, caulinis subhastatis, vaginis stipularibus cordato amplexicaulibus. Gouan. *Illustr. bot.* p. 65.

Cacalia foliis cordatis, ferratis, longissime petiolatis; floribus umbellatis. Hall.
fl. helv. n. 137. a.

Cacalia foliis crassis hirsutis. Bauh. *pin.* 198.
— Mapp. *fl. Alf.* p. 47.

Cacalia vulgaris incano folio. Clus. *hist.* 2.
p. 115.

on. Jacqu. *fl. Austr.* t. 235.

erm. Wollhaarige *Cacalie*.

all. *Cacalie petasite*.

ab. et *Flor.* in iisdem cum praecedente regionibus frequens. ¶

Affinis praecedenti, cum qua olim Linnaeus Hallerus combinauerunt. Differt autem. Folia subtus tomentosis, incanis, acutioribus, amplexiculis. Petiolis basi saepissime auriculatis. Calyce longiore, corolla duplo brevior. Corollae limbo minus patente. Flosculi in sinu florea 3-4 raro 6, dilute purpurei, aut albi.

Nota. Non confundenda cum *Cacalia tometosa*. Villars. fl. Delph. 3. p. 171. L. mark. Encycl. 1. p. 531. seu *Cacalia leucophylla*. Willden. Linn. Sp. pl. 3. 1736. foliis utrinque denso et candidissimo mento præditis, floribus umbellato-paniculatis, calycibus hirsutis multifloris, sæpe vigintifloris. Hanc frustra hucusque in montosis subalpinis stris quæsi.

CCCLXXVII. EUPATORIUM

Linn. Gen. pl. ed. Schreb. n. 1272.

Calyx communis oblongus, subcylindricus imbricatus: *Squamis* lineari-lanceolatis, erectis, inæqualibus. *Corolla* composita, unilobata, tubulosa. *Corollulae* hermaphroditae, æquales. *Propria* monopetala, infundibuliformis limbo quinquefido, patulo. *Stylus* longissimus erectus, superne bifidus. *Calyx* fructiferus immutatus. *Semina* oblonga. *Pappus* subpluriflorus. *Receptaculum* nudum.

1250. EUPATORIUM *cannabinum*.

calycibus quinquefloris, foliis digitalibus.
Linn. Sp. pl. 2. p. 1173. — Pollich. pal. n. 777.

Eupatorium foliis ovato-lanceolatis, trisectis. Hall. stirp. helv. n. 136.

Eupatorium cannabinum. Bauh. pin. 310.
Mapp. fl. Als. p. 103.

Herba Kunigundis I. Tabernæm. p. 386.

I con. Oeder. fl. Dan. t. 745. — Blackw. t. 1.
Germ. Alpkraut. Kunigundenkraut. Wäldst. Wasserhanfkräut.

Call. Eupatoire chanvrin. Eupatoire d'Avicenne.

Tab. Utrinque in virgultosis sylvaticis, pratensibus humidis, ad pratorum marginalia dumetosa inundata, ad fossas aquaticas cis et trans Rhenum passim abunde.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☞

Radix repens, fibrosa. *Caulis* 3-6 pedalis, erectus, tetragonus, villosus, ramosus. *Rami* oppositi. *Folia* opposita, ternata, quinata; *foliis* ovato-lanceolatis serratis, subvillosis, terminali majori. *Flores* caulem ramosque terminantes, cymoso-umbellati, numerosi, densi, colore carnei, purpurei et albidus. *Calyces* quiniflori.

FORM. EUPATORII Herba; Radix.

Planta amara, subacris. Febres intermittentes, per pedum putrida, infuso Eupatorii debellari, testatur Boerhavius. In Hydrope, Cachexia et jctero commendatur. Planta omnino efficax.

CCCLXXVIII. CHRYSOCOMA.

Linn. Gen. pl. ed. Schreb. n. 1275.

Calyx communis hemisphæricus, imbricatus: *squamis* linearibus acuminatis, extus convexis. *Corolla* composita tubulosa, calyce longior. *Corollulae* hermaphroditæ, tubulosæ, numerosæ, æquales. *Propria* monopetala, infundibuliformis: *limbo* quinquefido, revolutus. *Calyx* fructifer vix mutatus. *Semina* ovato-oblonga compressa. *Pappus* capillaris, sessilis. *Receptaculum* nudum, planum.

396 SYNGENESIA. POLYG. ÆQUALIS.

1251. **CHRYSOcoma** *Linosyris* herbacea, foliis linearibus glabris, calycibus laxis. *Linn. Sp. pl. 2. p. 1178.* — *Ped. fl. pal. n. 778.* — *Hall. stirp. herb. n. 1251.*

Conyza Linariae folio, Mapp. fl. Alf. p. 8.

Linosyris nuperorum. Lobel. hist. 223. Icon. 409.

Linaria folioso capitulo luteo major et minor. Bauh. pin. 313.

Linaria III, aurea. Tabernaemont. p. 120. Icon.

Icon. Schkuhr. bot. handb. t. 238.

Germ. Goldschopf. Golden-Leinkraut.

Gall. Chrysocome linière.

Hab. In collibus graminosis asperis aridis montis Brisgoviae am *Kaiserstuhl*, prope *Roßweil* cum *Seseli Hippomarathro*, et alibi nuper cum *Generosissimo Ittnero* legi. Margraviatu superiore in collibus aridis prope *Efringen* infrequens *t. Vulpio*. *Alsatia* prope *Gemar* in aridis; inter *Wiesleam* et *Colmariam t. Joh. Bauhino*. *Palatinatu* in montibus sylvestribus circa *Stadelsbach*; in vinearum collibus lapideis circa *Dürkheim* cum *Euphrasia lutea*; inter *Obernheim* et *Odernheim* ad vias graminosas post *Pollichium* vidi.

Flor. Augusto, Septembri. *Semina* Octobri matura. 7.

Caulis pedalis, sesquipedalis et altior, erectus, teres, durus, scabriusculus, foliosus, semperne ramosus, dense corymbosus. *Folia* sessilia, sparsa, linearia, integra, angusta, patula, integerrima, scabriuscula, late viridia. *Flora*

SYNGENESIA. POLYG. SUPERF. 397

scorimbofi, caulem terminantes, aurei. *Petali* erecti, sub flore crassiores, polyphylli. *Folia* lineari-setacea, patenti-curvula. *Calyxae* squamae lineari-setaceae, patulae.

Planta pulchra, culta major, digitis trita aromatica.

Flores melliferi.

**SYNGENESIA. POLYG. SUPERFLUA.
CCCLXXIX. TANACETUM.**

Linn. Gen. plant. ed. Schreb. n. 1280.

Calyx communis hemisphaericus, imbricatus: *squamis* acutis, compactis. *Corolla* compacta tubulosa, convexa. *Corollulae* hermaphroditae numerosae, tubulosae in disco. *Feminae* nonnullae in radio. *Propria* hermaphroditica infundibuliformis: *limbo* quinquefido, reflexo. *Femina* trifida, introrsum profundius fissi. *Calyx* fructiferus immutatus. *Semina* oblonga. *Pappus* submarginatus. *Receptaculum* convexum, nudum.

1252. **TANACETUM *vulgare*** foliis bipinnatis, incis, ferratis. *Linn. Sp. pl. 2. p. 1184.* — *Pollich. fl. pal. n. 779.*

• **Tanacetum** foliis pinnatis, pinnis semipinnatis, acute dentatis. *Hall. stirp. helv. n. 132.*

• **Tanacetum *vulgare* luteum.** *Bauh. pin. 132.*
— *Mapp. fl. Alf. p. 299.*

Artemisia tertia Dioscoridis *3. Tabernæm. p. 28. Icon,*

398 SYNGENESIA. POLYG. SUPERF.

Icon. Bulliard. herb. t. 187. — Regnault. bot.
— Oeder. fl. Dan. t. 871. — Blackw. t. 464

Germ. Gemeine Rheinfarren. Wurmkraut.

Gall. Tanaise vulgaire. Herbe aux vers.

Hab. Utrinque ad vias dumetosas, ad agro-
rum marginalia praesertim in Rheni vicini-
abunde passim.

Flor. Julio, Augusto. *Semina* Septembri ma-
tura. ☞.

Radix ramosa, fibrosa, dura. *Caulis* tripe-
dalis, et altior, erectus, angulosus, duriuscu-
lus, glaber, superne ramosus. *Rami* alterni,
erecti, simplices. *Folia* alterna, sessilia, am-
plexicaulia, duplicato-pinnata, dentata, ser-
rata, viridia, punctulata. *Flores* corymbosi,
terminales, fastigiati, aurei, densi, numerosi.
Semina cylindrico-turbinata, angulata, utrius-
que truncata, parva.

Pharm. TANACETI Herba; Flores; Semina.

Planta intense amara, fragrans, incidentes, resolu-
ventes et roborantes vires habet: a vetustissimis tem-

nutatus. *Semina* nuda. *Receptaculum* platum, nudum aut villosum.

3. ARTEMISIA *Abrotanum* foliis ramossissimis fetaceis, caule erecto suffruticoso. *Linn. Sp. pl.* 2. p. 1185.

Abrotanum mas angustifolium majus. Bauh. pin. 137. *Duham. Arbr.* 1. p. 20. t. 4.

Abrotanum mas. Tabern. p. 50. *Icon.*

n. *Blackw. herb. t.* 555.

m. *Stabwurz. Citronenkraut.*

l. *Armoise aurone.*

k. In Gallia, Italia, Hispania regionibus calidioribus. Colitur passim in hortis rusticorum, nec non in vineis *im Bühlerthal* et *am Kaiserstuhl* Brisgovia, ubi quasi spontanea.

r. Julio, Augusto. *Semina* Septembri matura. *Suffrutex.*

Caulis 2-3 pedalis et altior, erectiusculus, ne lignosus, teres, superne herbaceus, striatulus, foliosus, ramosus: *Rami* sparsi, prostrati, foliosi. *Folia* sparsa, alterna, petiolo duplicato-pinnatifida; *laciniis* linearibus, integris, elongatis, utrinque punctatis, subincano-lanuginosa. *Flores* caulem ramumque terminantes, racemosi, nutantes, subadi, pallide lutescentes, parvi. *Calyces* minusve pubescentes.

r m. ABROTANI *Herba.*

Planta fragrans, aromatica, grata, amara, stomachica, emmenagoga. Fidele ab omni temuliculis rusticis vino generoso patrio superfusa, camentum.

400 SYNGENESIA POLYG. SUBERT.

1254. *ARTEMISIA campestris* foliis
multifidis linearibus, caulibus procum-
bentibus virgatis. *Linn. Sp. pl. p.* 1185.
Poll. fl. pal. n. 780.

Artemisia caulibus procumbentibus, foliis
primis sericeis, adultis viridibus, pinnatis:
pinnis multifidis lanceolatis. *Helv. stirp. helv. n.* 131.

Abrotanum campestre. *Bauh. pin.* 136.
Mapp. fl. Alf. p. 3. (cauliculis rubentibus
et albicantibus.)

Abrotanum campestre. *Tabern. p.* 51. *Icon.*
Oeder. fl. Dan. t. 1175. — *Schkyhr. bot.*
handb. t. 240.

Germ. Feldbeyfuss. Feldwermuth.

Gall. Armoise champetre.

Hab. Utrique ad margines agrorum, vias
fabulosas apricas aridas passim frequens.
ca *Carlsruhe* et *Mühlburg* am Lande
ben abunde.

Flor. Julio, Augusto, Septembri. *Semin.*
Septembri, Octobri matura. 7

Radix ramosa, lignosa, dura, extus flav-
cans, intus albida. *Caules* ex radice plures, di-
bitales, tripedales et altiores, basi procum-
bentes, adscendentes, demum erecti, duri, ter-
etes, glabri, sæpe fusco-rubelli, quandoque
virides, ramosi. *Rami* copiosi, alterni, erecti.
Folia alterna, pinnato-multifida, pinnulis
neari-lanceolatis, angustis, 3-5-partitis,
niora sericea, adulta glabra. *Flores* caulem
terminali, terminantes, fusco-purpurascen-
tes, parvi, subnutantes racemosi. *Racemi* seci-
Calyces glabri. *Squamæ* virides, margine
cantes

ntes aut fusco-purpurascens, splendens, abrae, quandoque subpubescentes. *Receptaculum nudum.*

In regionibus sabulosis aridis arenam figit.

255. *ARTEMISIA pontica* foliis multipartitis subtus tomentosis, floribus subrotundis nutantibus; receptaculo nudo. *Linn. Sp. pl. 2. p. 1187. — Pollich. fl. pal. n. 781.*

Artemisia foliis duplicato-pinnatis, pinnulis parallelis tomentosis. *Hall. stirp. helv. n. 129.*

Abinthium ponticum tenuifolium incanum. *Bauh. pin. 138.*

Abinthium Ponticum Galeni. Tabernæmont. p. 17. Icon.

Jon. Jacqu. fl. Austr. t. 99. — Blackw. t. 527, et m. Römischer Beyfuß.

all. Armoise du Pont,

a b. In Palatinatu retro *Alzey* inter *Alwig* et *Niederulm* ad vias gramineas juxta agros cum *Althæa hirsuta* et *Inula germanica. t. Pollichio* et *Tabernaemontano.*

lor. Augusto, Septembri. *Semina* Septembri, Octobri matura. ♀

Radix ramosa, fibrosa, repens, extus flavas, intus albida. *Caules* ex una radice plures, bitales, tripedales, erecti, incani, foliosi, mosi. *Rami* alterni, sparsi, erecti, tenues, brusculi, axillares. *Folia caulina* alterna, bipinnata: *laciniis* linearibus, angustis, acutis, cinque glauco-tomentosa, *superiora* et *rametiora* pinnata; *summa* simplicia linearia,

acuta, integra. Flores caulem ramosque terminantes, racemosi, secundi, parvi, subrotundi, nutantes, tomentoso-incani, disco lutei. Receptaculum nudum. Semina nuda.

Pharm. ABSINTHII PONTICI Herba.

Planta grate aromatica, amara, tonica, stomachica.

Nota. Ad rariores Floræ cives pertinet.

1256. ARTEMISIA *Absointhium* foliis compositis multifidis, floribus subglobosis pendulis: receptaculo villoso. *Linn. Sp. pl. 2. p. 1188.*

Absointhium foliis incanis, pinnatis, pinnis primis bilobatis, pinnulis incisiss lanceolatis, spicis heteromallis, floribus pendulis. *Hall. stirp. helv. n. 124.*

Absointhium Ponticum seu *Romanum* officinarum, seu *Dioscoridis*. *Bauh. pin. 138.*
— *Mapp. fl. Alf. p. 3.*

Absointhium montanum f. *romanum*. *Taberna. p. 2. Icon.*

Icon. Regnault. bot. Blackw. t. 17.

Germ. Wermuth.

Gall. Absinthe.

Hab. Circa *Carlsruhe* in ruderatis passim, am *Landgraben* versus *Mühlburg* sine dubio ex hortis elapsa. In *Brisgovia* am *Kaiserstuhl* in muris arcis *Limburg*, nec non in rupibus prope *Burgheim* et *Altbreisach*. In Ditione *Basileensi* in muris arcis *Ramstein*. In Principatu *Fürstenbergensi* in rupibus et muris arcis *Heiligenstein* frequens, in rupibus *Hohen-Engen* prope *Engen* et alibi in summis arcium muris nuper vidi.

SYNGENESIA. POLYG. SUPERF. 403

lor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. 7

Caulis 2-3 pedalis et altior, erectus, strio-angulatus, villosus, albidus, ramosus. *Racemi* sparsi, alterni, erecti, simpliciusculi, striati. *Folia* alterna, petiolata, pinnatifida: *pinnis* alternis pinnatifidis, cuneiformibus: *laciniis* linearibus, obtusiusculis, mollia, villosa, punctata, *superiora* sessilia, cuneiformia, trilobata. *suprema* integra, lineari lanceolata. *Flores* nulem ramosque terminantes, nutantes, sublobosi, lutei, racemoli. *Racemi* compositi, obliqi, secundi, virgati. *Bractee* infra pedicellos lineari-subulatae. *Receptaculum* villosum. *Pharm.* ABSINTHII Herba.

Planta intense amara, aromatica, antipudredinosa, emica, stomachica, antacida, resolvens, anthelmintica.

Cerevisiae acidae immissa, mox acescentiam tollit. *inn.* fl. Suec. p. 286.

Flores apibus melliferi.

257. *ARTEMISIA vulgaris* foliis pinnatifidis planis, incisis, subtus tomentosis, racemis simplicibus, floribus ovatis, radio quinquefloro. *Linn. Sp. pl.* 2. p. 1189. — *Pollich. fl. pal.* n. 782.

Artemisia foliis pinnatis inferne tomentosis, pinnis acute dentatis, spica paniculata, erecta. *Hall. stirp. helv.* n. 130.

Artemisia vulgaris major, caule et flore purpurascens. *Bauh. pin.* 137. — *Mapp. fl. Als.* p. 28. et caule ex viridi albicante. *ib.*

Artemisia I. rubra, et II. alba. *Tabernæm.* p. 27. *Icon.*

494 SYNGENESIA, POLYG. SUPP.

Icon. Bulliard. herb. t. 350. — Oeder. fl. Dau. t. 1176. — Blackw. t. 431.

Germ. Gemeiner Beyfuss.

Gall. Armoise vulgaire.

Hab. Utrinque in ruderatis pagorum et urbium ad muros, sepes, ripas fluviorum passim abunde.

Flor. Augusto, Septembri, Octobri. *Semin.* Octobri, Novembri matura. 3

Radix ramoso-fibrosa, repens. *Caulis* 3-4 pedalis, erectus, striato-angulatus, glaber sæpe ex rubello-viridis, ramosus. *Rami* alterni simpliciusculi, virgati, striati, pubescentes, erecto-patentes. *Folia* alterna, sessilia, super lineata, viridia, glabra, subtus tomentosa, albide cinerea, pinnatifida: *laciniis* linearilanceolatis, *lobis* pinnatifidis, suboppositis, lanceolatis, acutis: *laciniis* lineari-lanceolatis acutiusculis, subdentatis, *inferioribus* integris *infinis* lunulatis amplexicaulibus. *Flores* ramoso paniculati, caulem ramosque terminantes, erecti, ovati, albido-cinerei, disco flavi. *Florum Radius* sæpe 6-7-8 florus. *Receptaculum* nudum.

Pharm. ARTEMISIAE RUBRAE f. ALBAE Herba

Planta praecedente minus amara.

Flores apibus mel praebent.

Ex Artemisia per combustionem multum Potassae redundat.

1258. *ARTEMISIA Dracunculus* foliis lanceolatis glabris integerrimis. *Linn. Sp. pl. 2. p. 1189.*

Abrotanum lini folio acriori et odorata. *Tournef. inst. 459.*

SYNGENESIA. POLYG. SUPERF. 405

Dracunculus hortensis. Bauh. pin. 98.

Dracunculus hortulanus. Tabern. p. 855. Icon. 20 n. Blackw. t. 116. — Regnault. bot. crm., Dragun. Kaifersallat.

all. Esdragon.

ab. In Sibiria. Tataria. Colitur in hortis et passim in vineis.

lor. Augusto, Septembri. Semina Octobri matura. 7.

Radix ramoso-fibrosa, repens. *Caules* ex radice plures, 2-3 pedales, erecti, superne bangulati, glabri, læte virides, ramosi. *Rami* terni, erecti. *Folia* alterna, sessilia, lanceolata, integra, acuta, utrinque glabra, læte viridia, *ramca* minora, angustiora, patentia. *Flores* caulem ramosque terminantes, racemosomiculati, parvi, gemini, ovati, nutantes, lco-ferruginei. *Calyx* viridis, subpubescens. *reptaculum* nudum.

h arm. DRACUNCULI Herba.

Planta grate aromatica, suaveolens, fragrans, vica, stomachica, incitans appetitum provocans Folia cata et recentia adhibentur in culinis ad cibi conditum gratissimum. Persis ab antiquissimis retro temporibus tanquam quotidianum ciborum condimentum tiffima. Acetum odore et sapore gratissimum.

Ad embamata et acetaria pæsertim recipitur.

. CCCLXXXI. GNAPHALIUM.

mus ex Filaginis Linn. speciebus pluribus et Gnaphalio constructum.

Calyx communis imbricatus, rotundatus: *lamis* marginalibus scariosis, coloratis. *Colla* composita: *Corollulae* hermaphroditæ,

206 STINGENESIA. POLYGA. SUPPER

tubulosae, interdum femineis apetalis mixtae. *Propria* hermaphroditis infundibuliformis, limbo quinquefido, reflexo. *Femineis* (6 quae) corolla nulla. *Stamina* Hermaphroditis *Filamenta* quinque, capillaria, brevissima. *Anthera* cylindracea. *Pistillum* Hermaphroditis *Germen* ovatum. *Stylus* filiformis, longitudine flaminum. *Stigma* bifidum. *Femineis* *Germen* ovatum. *Stylus* filiformis longitudine hermaphroditis. *Stigma* bifidum, reflexum. *Calyx* fructiferus persistens, nitidus. *Semina* hermaphroditis oblonga parva, coronata *Pappo* capillari s. plumoso. *Femineis* similia hermaphroditis. *Receptaculum* nudum.

Nota. Datur species floribus dioicis.

GNAPHALIA FILAGINNOIDEA plurimas Flagellis species Linneanas continent, isthic transferenda. Est *Calyx* communis teres s. pentagonus, imbricatus. Squamis ovato-lanceolatis, exterioribus acutis, tomentosis, interioribus nitidis, coloratis, acuminatis. Corolla composita: Corollulis hermaphroditis, tubulosis, paucis in centro disci. Femineis tubulosis numerosis in reliqua parte disci: Femineis aliis paucis, fere apetalis, inter squamas calycis exteriores. *Propria* Hermaphroditis infundibuliformis: limbo quadrifido, patente. Femineis in disco, infundibuliformis, tubo tenui basi tumido; limbo quadrifido, erecto. Femineis inter squamas calycis, vix conspicuas, tubo tenuissimo, limbo bifido, acuto. *Stamina* hermaphroditis. *Filamenta* quatuor, brevissima. *Anthera* tubulosa. *Pistillum* Hermaphroditis *Germen* exiguum, abortiens. *Stylus* capillaris, longitudine limbi. *Stigmata* duo, erecto-patula. Feminis in disco: *Germen* oblongum. *Stylus* capillaris, limbo longior. *Stigmata* duo, patentia. Feminis intra calycem: *Germen* oblongum. *Stylus* capillaris, limbo longior. *Stigmata* duo, longa, patentia. *Calyx* fructiferus in-

~~pinnatis~~, Semina hermaphroditis sterilia, pappo coronata. Feminis in disco oblonga, pappo coronata. Pappus brevis simplex. Feminis intra calycem, oblonga, nuda, Receptaculum nudum.

† *Chrysocoma*.

259. *GNAPHALIUM arenarium* herbaceum foliis lanceolatis: inferioribus obtusis, caule simplicissimo, corymbo composito. *Linn. Sp. pl. 2. p. 1195.* — *Pollich. fl. pal. n. 783.*

Elychrysum arenarium. *Lamarck et Decandolle. fl. franc. 4. p. 132.*

Elychrysum f. *Stochas citrina latifolia.* *Bauh. pin. 264.*

Stoechas citrina germanica, latiore folio. *Bauh. hist. 3. p. 153.*

Amaranthus luteus L. *Tabern. p. 778.*

Icon. Oeder. fl. Dan. t. 641 — *Blackw. t. 524.*

Germ. Gelbes-Sand-Ruhrkraut. Immerschön. Gelbe Rheinblume.

Gall. Elychryse des sables.

Hab. Utrique in regionibus apricis graminosis fabulosis sterilibus cis et transrhenanis passim frequens. Prope *Carlsruhe*, *Mühlburg auf dem Hardthof* et alibi in fabulosis asperis abunde.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☞

Caulis palmaris, pedalis et altior, erectus, teres, albide tomentosus, simplex. *Folia* alterna, sessilia; semiamplexicaulia, lineari-lanceolata, obtusa, integra, utrinque tomentosa, mollia, cinereo-albida. *Flores* caulem terminantes,

408 SYNGENESIA. POLYG. SUPERF.

corymbosi. *Corymbi* pedunculati, subglobosi, densiusculi, multiflori, speciosi, aurei, aut aurantiaci. *Calycinae* squamae ovatae, obtusae, splendentes, glabrae, aureae aut aurantiae, persistentes.

Obs. Prostant individua flosculis omnibus mere hermaphroditis 6-7 fidis, instructa. Schkuhr. bot. handb. 3. p. 87.

Nota. Mappus citat *Gnaphalium Stoechas* Linn. *Elichrysum* seu *Stoechas citrina* angustifolia. Bauh. pin. 264. in Alsatiae aridis prope Hagenau auf der Hart, et in pascuis frutibus aridis ad Rhenum, ubi autem non crescit. Copiose ibi vidi *G. arenarium* nostrum. Hic *G. Stoechas* in regionibus calidioribus Galliae Italiae et Hispaniae obveniens, fruticosum, omnino in Flora delendum.

Pharm. *STOECHADIS CITRINAE* Flores.

Egregium in graminosis sabulosis saepe decus.

1260. *GNAPHALIUM luteo - album*
herbaceum, foliis semiamplexicaulis, ensiformibus, repandis obtusis utrinque pubescentibus, floribus conglomeratis. Linn. *Sp. pl.* 2. p. 1196. — Pollich. *fl. pal.* n. 784.

Filago foliis mollissimis, tomentosis, obtusis umbella convexa, electrina. Hall. *fl. helv.* n. 147.

Elichrysum sylvestre latifolium, capitulis conglomeratis. Bauh. *pin.* 264.

Icon. Plucken. Alm. 171. t. 31. f. 16.

Ger m. Gelbweisses Ruhrkraut.

Gall. Perlier a fleurs jaune et blanc..

Hab. Circa Carlsruhe in cultis arenosis; im Beurtheimer Wald, am Landgraben et

SYNGENESIA. POLYG. SUPERF. 409

alibi non infrequens. In Alsatia prope Argentinam in agris fabulosis in der *Ruprechtsau* et alibi vidi. In Palatinatu rarius circa *Lautern* in pascuis arenosis ficcis; in vinearum collibus ficcis circa *Dürkheim*; inter *Dürkheim* et *dem Holz-hof* cum *G. arena-rio* copiose t. *Pollichio*.

lor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ☉

Caulis pedalis, sesquipedalis et altior, erectus, teres, albide tomentosus, mollis, plus minusve ramosus. *Rami* alterni, erecti, simpliciter. *Folia* alterna, sessilia, semiamplexicaulia, linearia, obtusa, superiora acutiuscula, subundulata, margine reflexa, integra, utrinque citreo-albide-tomentosa, mollia. *Flores* caulinum ramosque terminantes, corymbosi, globoso-fasciculati, parvi. *Flosculi* feminei albidi, hermaphroditi, pallide lutei. *Calycinae* squamæ lato-lanceolatae, obtusiusculæ, argenteæ, pallesque lutescentes, splendentes. *Pappus* cellularis.

†† *Argyrocoma* f. *calyce albo*.

261. GNAPHALIUM dioicum sarmentis procumbentibus, caule simplicissimo, corymbo simplici terminali, floribus dioicis. *Linn. Sp. pl.* 2. p. 1199. — *Pollich. fl. pal.* n. 785.

Antennaria dioica. *Gaertn. Sem. et fruct. pl.* 2. p. 410. t. 167. f. 3. *Wetterauische flora.* 3. p. 193.

Filago flagellis reptans sexu distincta, flosculis omnibus androgynis. *Hall. stirp. helv.* n. 157.

440 SYNGENESIA. POLYGAEEAE

Elichrysum montanum flore rotundi
purpureo-candido-suave rubent
gato-longiore folio et flore pu
et albo. *Tournef. inst. Mapp. fl.*
96. 97.

Gnaphalium montanum rotundiore
giore folio. *Bauh. pin.* 263.

Gnaphalium montanum album-pur
et suave rubens. *Tabernaemont.*
Icones.

Icon. Bulliard. herb. t. 325. — *Ragna*
Germ. Katzenpfötgen. Weisse und roth
öhrlein. Engelsblümlein. (Belchen
nostr.)

Gall. Perliere dioique. Pitechatier.

Herb. Utrique in pascuis sterilioribus,
nolis aridis apricis, ericetis, ju
passim frequens. Abunde in pascu
tosis altioribus in montibus *Belchen*,
et *Feldberg*.

Flor. Majo, Junio, Julio. *Semina* Jul
gusto matura. 7

Radix longe lateque repens. *Cauli*
lis, palmaris et altior, erectus, teres,
tomentosus, mollis, simplex. *Folia* r
numerosa, in petiolum angustata, obve
ceolata, obtusa, integra; *caulina* alter
filia, linearia, omnia subtus albide ton
Flores caulem terminantes, 3-4-7-9
lato-corymbosi, medio reliquis sæpe
albi, incarnati, purpurei, suave rubei
variegati. *Calycinæ* squamæ imbricatæ,
latæ, obtusiusculæ, tomentosulæ, exterio
riores, interiores longiores, apice pu
aut albæ, splendentes, patenti reflex

osculi androgyni atque femineis mixti. *Pap-*
as capillaris apice penicillatus. *Receptaculum*
robiculatum marginibus denticulatis scabrum.

harm. GNAPHALII f. PILOSELLÆ f. PEDIS
CATI Flores.

Planta subadstringens hodie medicinæ fere exul.

In pascuis montis Belchen pulcherrima, et abun-
 tus, a monticulis Marggraviatus superioris ibi sub no-
 mine vernaculo Belchen-Röslein, nota.

262. GNAPHALIUM *margaritaceum*

herbaceum, foliis lineari-lanceolatis,
 alternis, caule superne ramoso corym-
 bis fastigiatis. *Linn. Sp. pl. 2. p. 1198.*

Elichrysum margaritaceum. Lamarck et De-
cand. fl. franc. 4. p. 131.

Filago foliis lanceolatis, viridibus, subtus
 tomentosis, floribus umbellatis. *Hall.*
stirp. helv. n. 146.

Gnaphalium americanum. Clus. hist. 1. p. 327.

r m. Perlen-Ruhrkraut.

11. Elychryse perlé.

b. In america boreali, Kamtschatka, in mon-
 tibus Angliæ, Galliæ, Helvetiæ? Colitur
 passim in hortis rusticorum et aliorum præ-
 fertim in Marggraviatu superiore.

r. Julio, Augusto, Septembri. *Semina* Sep-
 tembri, Octobri matura. *z*

Radix repens. *Caulis* pedalis, sesquipeda-
 erectus, totus albus, tomentosus, superne
 natus. *Folia* alterna, sessilia, lineari-lanceo-
 la, acuta, integra, subtus tomentosa. *Flores*
 aulem ramosque terminantes, umbellato co-
 mpositi, albi, numerosi, subglobosi.

Filago foliis linearibus, alis spiciteri
flor. helv. n. 148.

- α. *Gnaphalium rectum* caule erecto fi
foliis lineari-lanceolatis subtus
floribus terminalibus et axillarib
libus spicatis. *Willden. Linn. Sp.*
1885. — *Smith. fl. brit. 2. p.*
Wetterauische flor. 3. p. 191.

Gnaphalium sylvaticum. Scopol. fl. (
1046. t. 56.

Gnaphalium spicatum. Mapp. fl. Al

Gnaphalium rectum. Bauh. hist. 3. p

Gnaphalium anglicum vel Belgicum
Icon. 482.

Gnaphalium III. majus. Tabernæm.
Icon bona.

- β. *Gnaphalium norvegicum* caule h
simplicissimo, foliis lanceolatis
acutis trinerviis, subtus lanugine
ca florum terminali, foliosa. *Jac*

n. *α. Oeder. fl. Dan. t. 1229. — Schkuhr. bot. handb. t. 243. (Icon optima.) — β. Oeder. fl. Dan. t. 254.*

m. *Wald-Ruhrkraut.*

l. *Perlier des bois.*

i. *α. Utrinque in sylvis fabulosis ficcioribus, passim frequens. β. in sylvis fabulosis humidiusculis passim cum α. non infrequens.*

r. *Augusto, Septembri. Semina Septembri, Octobri matura. ♂*

Variatio α. Caulis pedalis, sesquipedalis, tus, strictus, foliosus, simplex, tomentosus. Folia alterna, sessilia, lineari-lanceola, acutiuscula, integra, superne viridia, subtomentosa, incana, erecta. Flores ex axillis eorum subpedunculati, inferiores subsolitarii, priores sensim aggregati, in spiculas axillares les, fusco-spadiceas. Calycinae squamæ imatæ, ovatæ, obtusæ, infimæ tomentosae, pæ glabrae, margine pellucidæ, striis fusco-icantibus percussæ, interiores exterioribus giores. Flosculi 3-4-5 hermaphroditi, cenz, externi feminei, plurimi. Pappus oculo ato plumosus.

l. Prostant individua a parte caulis inferiore ad apicem usque ex alis foliorum floriferi; alia a medio caule floriferi, alia in summo tantum floriferi cum rudimentis florum in axillis foliorum saltem superiorum. Folia saepe angusta, acuta, quandoque superne latiuscula, et obtusa, plus minusve utrinque, aut saltem subtus sericeo-tomentosa. Hinc Variatio β. floribus superne in spicam foliosam densiorem congestis. vix ob variationes intermedias, specie ab α. distincta.

ota. Individua variationis β. in Lapponia ab amico quodam lecta, simillima sunt illis, quæ in regionibus nostratibus, nec non in Helvetia, Gallia et Hispania vidi.

Gnaphalium longifolium humile ramis
capitulis nigris. *Raj. hist.* 295.

α. Gnaphalium nudum floribus nudis.
fl. germ., p. 292.

**β. Gnaphalium tomentosum floribus
tosis.** *Hoffm. l. c.*

Icon. Oeder. fl. Dan. t. 859.

Germ. Morastiges Ruhrkraut.

Gall. Perlier des marais.

Hab. α. et β. Utrunque in fossis exsicca
midis inundatis passim abunde cur
Busonio.

Flor. Julio, Augusto. **Semina** Augusti
tembri matura. ☉

Radix fibrosa. **Caules** ex una radi
plures, digitales, palmares, erectiuscu
tes, albide tomentosi, molles, ramosi
alterni, diffusi. **Folia** alterna, sessilia,
lanceolata, integra, obtusiuscula, utrin
minusve tomentolula. **Flores** caulem ra
terminantes, subsessiles, 5 - 7 - 13 fasc

165. GNAPHALIUM germanicum
 caule herbaceo erecto dichotomo, foliis
 lineari-lanceolatis tomentosis, floribus
 globofo-fasciculatis lateralibus termina-
 libusque tomentosis.

Gnaphalium germanicum. Smith. fl. brit. 2.
p. 874. — Willden. Linn. Sp. pl. 3. p.
1894. — Lamarck et Decandolle. fl. franc.
4. p. 135.

Filago germanica panicula dichotoma, flo-
 ribus rotundatis axillaribus hirsutis, fo-
 liis acutis. *Linn. Sp. pl. 2. p. 1311. —*
Poll. fl. pal. n. 830.

Filago caule dichotomo, corymbo sphaeri-
 co, in angulo sessili. *Hall. stirp. helv. n.*
153.

Filago seu Impia. *Dodon. pempt. 66. Mapp.*
fl. Alf. p. 105.

Gnaphalium germanicum. Bauh. hist. 3. p.
158.

Gnaphalium II. vulgare. *Tabernæmont. p. 780.*
Icon.

n. Oeder. fl. Dan. t. 997. — Schkuhr. bot.
handb. t. 266.

erm. Teutsches Ruhrkraut. T. Fadenkraut.
all. Perlier d'Allemagne.

ab. Utrique passim in arvis et agris inter sti-
pulas post messem et pascuis sterilibus are-
notis frequens.

for. Junio Julio, Augusto. Semina Augusto,
Septembri matura. ☉

Radix ramoso-fibrosa. *Caulis* palmaris,
 pedalis, sesquipedalis, erectus, teres, albide

416 SYNGENESIA. POLYG. SUPERF.

tomentosus, aut incanus, ramosus. *Rami* pariter alterni, erecti, longiusculi, dichotomi, foliis lanuginosi. *Folia* alterna, sessilia, lineari-lanceolata, acutiuscula, integra, utrinque albidis lanuginosa, mollia. *Flores* ad caulis et ramorum dichotomiam sessiles, terminalesque densi conglomerati, in capitula rotunda ex albidis lutea, tomentosa, multiflora. *Calycinae* squamae apice terminatae in aristam brevem fuscam. Inter calycis squamas flosculi feminei pappo desituti. *Flosculi* in disco feminei plures, pauci hermaphroditi, omnes feminibus perfectis, pappo plumoso praediti.

In sabulosis arenam figit.

1266. *GNAPHALIUM gallicum* caule herbaceo erecto, subdichotomo, foliis linearibus revolutis acutis, floribus subulatis confertis axillaribus. *Smith. fl. brit.* 2. p. 874. — *Willden. Linn. Sp. pl.* 3. p. 1895. — *Lamarck et Decand. fl. franc.* 4. p. 136.

SYNGENESIA. POLYG. SUPERF. 417

Tab. In arvis macilentis locisque sabulosis elevatis passim. Circa *Carlsruhe* retro *Mühlburg* prope *Grünwinkel* in arenosis asperis cum *Sagina erecta*. In Margraviatu superiore in arenosis prope *Zienken*. In sylva prope *Haltingen*. Prope Argentinam circa *Osswald* in agris arenosis. In Palatinatu circa *Lautern* raro t. *Pollichio*.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caulis digitalis, palmaris, erectus, incanus, subtomentosus, dichotomus, exsuccus. *Folia* alterna, sessilia, linearia, angusta, acuta, sessiliformia, apice subreflexa, incana, vix tomento manifesto praedita. *Flores* ex dichotomia sessiles, conglomerati, densi, conferti coeco-subulati ex viridi canescentes. *Flosculi* in praecedente et sequentibus speciebus.

267. GNAPHALIUM arvense floribus conicis lateralibus, caule herbaceo paniculato. *Linn. Sp. pl.* 2. p. 1312. — *Poll. fl. pal.* n. 833. (sub *Filagine*.)

Gnaphalium arvense *Lamarck et Decand. fl. franc.* 4. p. 136. — *Willden. Linn. Sp. pl.* 3. p. 1897.

Filago caule recto ramosissimo, foliis mollibus tomentosis, florum acervis sessilibus in spicas coalescentibus. *Hall. stirp. helv.* n. 154.

Filago altera. *Dodon. Pempt.* 67. — *Mapp. fl. Als.* p. 105.

Gnaphalium majus angusto oblongo folio. *Bauh. pin.* 263.

Caulis palmaris, pedalis, erectu
tomentosus, ramosus. *Rami* copiosi,
terni, erecto-patentes, foliosi, lanug
bi, superne divisi. *Folia* sparsa, alterna
femiamplexicaulia, lineari-lanceolata
cula, integra, utrinque alba, toment
res caulem ramosque terminantes, si
conglomerati, tomento albo connexi

Distinguitur a reliquis speciebus, Ca
fere, ad apicem ramofo, Foliis magis l
tomentosis, albis, Floribus maxime tom

1268. GNAPHALIUM *montanu*

herbaceo stricto subdichotomo
neari lanceolatis adpressiusculi
mentosis, capitulis florum axilla
minalibusque paucifloris, coni

Gnaphalium montanum. Lamarck e
fl. franc. 4. p. 136. — *Willd.*
Sp. pl. 3. p. 1896.

Filago montana caule erecto, divi
bus conicis lateralibus axillarib
Sp. pl. 3. p. 1311. — *Dellich fl.*

- l.** *Gnaphalium minimum* caule erecto ramoso, foliis lanceolatis acutis, planis, floribus conicis subconfertis lateralibus terminalibusque. *Smith. fl. brit. 2. p. 873. — Willden. Linn. Sp. pl. 3. p. 1896.*

cr m. Berg-Ruhrkraut.

all. Perlier des montagnes.

- g b.** Utrunque passim in collibus apricis, macilentis sabulosis cum β . circa *Carlsruhe* versus *Mühlburg am Landgraben* in sabulosis, *auf dem Hardthof* et alibi.

lor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caulis digitalis, palmaris, pedalis, erectus, setus, teres, gracilis, incano-subtomentosus, **folius**, superne ramoso-dichotomus. **Rami** angati, tenues, foliosi, tomentosi. **Folia** sparsa, ternata, sessilia, lineari-lanceolata, acuta, infra, cauli subadpressa, incano-tomentosula, **ramulae** sublatiora, breviora, magis tomentosa. **Flures** in dichotomia, axillares, terminalesque, 5 conglomerati, subsessiles, conico-pyramidales, incani, tomentosuli.

varietas β . pusilla, 2-3 uncialis, erecta, in omnibus partibus simillima, tantum magis tomentosa, depauperata. Folia etiam cauli adpressa.

Nota. Synonyma veterum ad hanc plantam et variationem β . omnino difficillime eruenda.

- bc.** Variatio nostra β . quae a Cl. Willdenowio in Angliæ arenosis citatur, in omnibus partibus simillima, individuis prope Londinum lectis, ut in individuis ab acutissimo Schweyckerto mihi communicatis perspexi.

CCCLXXXII. CARPESIIUM

Linn. Gen. pl. ed. Schreb. n. 1284.

Calyx communis imbricatus; *foliolis* exterioribus majoribus, patulis, reflexis; *interioribus* æqualibus, arctis, brevioribus. *Corolla* complicita, æqualis. *Corollulae* hermaphroditæ in disci *Femineae* in ambitu. *Propria* hermaphroditi infundibuliformis: *limbo* quinquefido, patulis *Femineis* tubulosa, quinquefida, connivens. *Calyx* fructiferus immutatus. *Semina* obovata, nuda. *Receptaculum* nudum.

1269. *CARPESIIUM cernuum* floribus terminalibus. *Linn. Sp. pl. 2. p. 1203.* — *Hall. stirp. helv. n. 134.* — *Scopol. fl. Carn. n. 1030.*

Balsamita conyzæ folio, flore cernuo. *Vaill. act. 336.*

Chrysanthemum conyzoides cernuum foliis

SYNGENESIA. POLYG. SUPERF. 421

tern Hardt Alsatiae t. C. Bauhino et Lachenalio.

lor. Julio, Augusto. Semina Septembri maturatione. ¶

Radix obliqua, capillata, fusca. *Caulis* sesipedalis, cubitalis, erectus, teres, striatus, hirsutus, superne ramosus. *Folia* alterna, elliptico-lanceolata, rariter obsolete ferrulata, utrinque hirsuta. *Flores* caulem ramosque terminantes, solitarii, cernui, pallide et fordide lutei, pappuli. *Calycina* foliola exteriora, foliis radicis similia, longa, patentia, demum reflexa, inaequalia. *Receptaculum* nudum, planum, punctulatum. *Semina* pappo destituta, ovali-linearia, striata, fusca, ad apicem glanduloso-unguicula.

Nota. Ad rariores omnino Florae nostrae cives pertinet, hucusque in collectionibus Florae Germaniae omissa,

CCCLXXXIII. C O N Y Z A.

Linn. Gen. plant. ed. Schreb. n. 1286.

Calyx communis imbricatus, subrotundus, parrosus: *squamis* acutis, exterioribus patentiusculis. *Corolla* composita, tubulosa: *Corollulae* hermaphroditae, numerosae, tubulosae, in disco. *Femineae* numerosae similes in ambitu. *Propria* hermaphroditis infundibuliformis: limbo quinquesido, patulo. *Femineis* infundibuliformis; limbo trifido. *Calyx* fructiferus connatus. *Semina* oblonga. *Pappus* capillaris. *Receptaculum* nudum, planum.

*Conyza major vulgaris. Bauh. pin.
Mapp. fl. Alf. p. 79.*

*Conyza major. Tabern. p. 1243. Ico
Icon. Bulliard. herb. t. 342. — Oeder.
t. 622.*

Germ. Dürrewurz. Flöhkraut.

Gall. Conise vulgaire.

Hab. Utrunque passim in collibus asperis
fertim calcareis, dumetosis, apricis
non ad viarum sepes. In monte *Tham*
am Eisenhafen abunde.

Flor. Augusto, Septembri. *Semina*
bri, Octobri matura. ♂

Caulis cubitalis, tripedalis et altior
rectus, angulosus, villosus, superne ramosus
cymosus. *Folia* alterna, sessilia, ovato-lanceolata,
minutim serrulata, villosa, fulva. *Stemmatea* lanceolata,
angustiora. *Flores* numerosae terminales, luteae. *Calyx* ovali-cylindricus
mae calycinae apice scariosæ, fusconigrae.

SYNGENESIA. POLYG. SUPERF. 423

CCCLXXXIV. ERIGERON.

Linn. Gen. pl. ed. Schreb. n. 1287.

Calyx communis oblongus, cylindraceus, imbricatus: *squamis* subulatis, erectis, gradatim longioribus, subæqualibus. *Corolla* composita radiata: *Corollulæ* hermaphroditæ, tubulæ in disco. *Femineæ* ligulatae in radio. *Prothecia* hermaphroditi infundibuliformis: *limbo* quinquefido. *Feminea* ligulata, linearis, subulata, erecta, sæpius integerrima. *Calyx* fructiferus connivens. *Semina* oblonga, parva. *Pappus* longus, capillaris. *Receptaculum* nudum, planum.

271. *ERIGERON canadense* caule floribusque paniculatis hirtis, foliis lanceolatis ciliatis. *Linn. Syst. veg.* 754. — *Poll. fl. pal.* n. 789. — *Hall. stirp. helv.* n. 84.

Conyza annua acris alba elatior, linariæ foliis. *Morif. hist.* 3. p. 115. *Sect.* 7. t. 20. fig. 29.

Virga aurea virginiana hirsuta annua di flor pallido. *Zanoni nov. Icon.* 78. — *Mapp. fl. Alf.* p. 328.

Conyzella. *Dillen. cat. Giff. app.* 142.

Ger. m. Canadaisches Berufskraut.

Hall. Vergerette de Canada.

Tab. Utrunque in ruderatis, muris, sylvis caeduis, ad carbonariorum areas, in agris et arvis passim abunde.

SYNGENESIA POLYG. STERN.

Flor. Julio, Augusto. **Semina** Septembri, Octobri matura. ☉

Caulis cubitalis, tripedalis et altior, erectus, sulcatus, hirtus, ramosus, aut simpliciusculus. **Folia** radicalia ovato-lanceolata, sessilia sparsa, sessilia, patentia, lineari-lanceolata, integra vel rariter denticulata, villosa. **Fl.** raceminales, racemoso-paniculati, albi parvi. **Flosculi** radii albi, disci pallide lutei.

Planta ex America in Europam migravit: in qua quammaxime propagavit.

1272. ERIGERON *acre* pedunculis albis unifloris. *Linn. Sp. pl. 2. p. 1311.*
Poll. fl. pul. n. 490.

Erigeron caule alterne ramoso, petioliis unifloris semiflosculosis, pappum æquantibus. *Hall. stirp. helv. n. 85.*

After arvensis coeruleus acris. *Mapp. fl. Al. p. 12.*

Conyza coerulea acris. *Bauh. pin. 265.*

Conyza coerulea. *Tabern. p. 1244. Icon.*

Icon. Curtis. fl. Lond. — Schkuhr. bot. hand. t. 241.

Germ. Scharfes Beruskraut.

Gall. Vergerette acre.

Hab. Utrunque in sabulosis aridis apricis, collibus et muris passim non infrequens.

Flor. Julio, Augusto, Septembri. **Semina** Septembri, Octobri matura. ☉

Radix fibrosa. **Caulis** pedalis, sesquipedalis, erectus, rigidiusculus, striatus, sæpe purpurescens, foliosus, ramosus, aut simpliciusculus.

SYNGENESIA. POLYG. SUPERF. 425.

3. *Folia* alterna, sessilia, inferiora obtuse obovata, basi angustata, superiora lanceolata, angustiora, sæpe subtortuosa, reflexa, *ramæ* linearia, erecta, omnia hirsuta. *Flores* pedunculati. *Pedunculi* alterni, erecti, uniflori. *Flosculi* radii purpurei, erecti, disci flavescentes. *Calycinae* squamae subulatae, erectæ, purpureæ, hirsutæ.

Flores nonquam explicati.

Planta acriuscula olim in magicas numerabatur.

Flores melliferi.

CCCLXXXV. TUSSILAGO.

Linn. Gen. pl. ed. Schreb. n. 1289.

Calyx communis cylindraceus: squamis lanceolato-linearibus, aequalibus, discum æquantibus, submembranaceis. *Corolla composita*, varia. *Corollulae* in aliis hermaphroditæ, tubulosæ omnes, vel tantum in disco. *Femineæ* aliis nullæ, aliis ligulatæ. *Propria* hermaphroditæ fundibuliformis: limbo quinquefido vel quarfido, acuto, reflexo, calyce longiore. *Femineæ* nulla aut ligulata, angustissima, calyce longior, integra. *Calyx* fructiferus vix mutatus. *Semina* oblonga, compressa. *Pappus* capillaris, stipitatus. *Receptaculum* nudum.

273. *TUSSILAGO alpina* scapo subnudo unifloro, foliis cordato-orbiculatis crenatis utrinque virentibus. *Murr. Linn. Syst. veg. ed. 14. p. 755. — Hoff. Synops. 456.*

SYNGENESIA POLYD. 5012372

Petasites scapo unifloro, foliis reniformibus
Hall. Stirp. helv. n. 143.

Tussilago alpina rotundifolia glabra. *Bam-
 pin. 197.*

Tussilago alpina s. glabro folio. *Clus. hist. 1
 p. 113.*

Tacon. Jacqu. fl. Austr. t. 246.

Germ. Glatter Alpen-Hustattig.

Gall. Tussilage des alpes.

Hab. In Principatus superioris vicinia
Bregenz et in toto tractu alpium ex col-
 lectorum *am Höhenkasten et Hö-*
sentis nuper vidi. Nec non in sylva
Jurac.

Flor. Junio, Julio. *Semina* Augusto matura.

Radix reptans, stolonifera. *Scapus* ses-
 pedalis, pedalis, solitarius, erectus, teres, lu-
 ruginosus in medio circiter squama foliacea pra-
 editus, fusco-purpurascens. *Folia* radicalia po-
 tiolata, cordata, subrotunda, crenulata, cras-
 sifuscula, supra glabra, subtus virentia et villosa.
Flos terminalis, solitarius, erectus, discoides.
Squamae calycinae circiter quatuordecim, acu-
 tae, fusco-purpureae. *Flosculi* feminei fertiles
 in radio, apetalis; reliqui hermaphroditi tubo-
 losi, fertiles, limbo quinquesido purpureo in-
 structi. *Semina* striata. *Pappus* capillaris, ses-
 lis. *Receptaculum* nudum, punctatum, planum.

Amat subalpina calcarea, nec summa alpina ju-
 respuit.

1274. *TUSSILAGO discolor* scapo uni-
 floro subnudo, foliis cordato-orbiculatis
 crenatis, supra lucidis, subtus tomentosis

SYNGENESIA. POLYG. SUPERF. 457

incanis. *Murr. Linn. Syst. veg. ed. 14. p. 755. — Host. Synopf. 456.*

Tussilago alpina var. β . *Linn. Sp. pl. 2. p. 1214.*

Tussilago alpina rotundifolia canescens. Bauh. pin. 197.

Tussilago alpina prima canescente folio. Clus. hist. 2. p. 113.

con. Jacqu. fl. Austr. t. 247.

erm. Zweifarbiger Alpen-Huflattig.

Hall. Tussilage discolor.

Herb. et Flor. in iisdem cum praecedente regionibus. 7

Maxime affinis omnino praecedenti, a qua differt, *Foliis* superne lucidis, viridibus, glaberrimis, subtus semper denso tomento brevissimo incanis.

Obs. Planta cum praecedenti specie in horto botanico Carlsruhano culta, constantissima permanfit, hinc non immerito pro peculiari specie habenda.

1275. TUSSILAGO *Farfara* scapo imbricato unifloro, foliis subcordatis angulatis denticulatis, (flore radiato.) *Linn. Sp. pl. 2. p. 1214. — Pollich. fl. pal. n. 791.*

Petalites scapo unifloro, flore radiato. *Hall. stirp. helv. n. 143.*

Tussilago vulgaris. Bauh. pin. 179. — Mapp. fl. Alf. p. 315.

con. Curtis. fl. Lond. — Bulliard. herb. t. 329. — Regnault. bot. — Blackw. t. 201.

erm. Huflattig. Rosshuf. Rosshüflein.

Hall. Tussilage. Le Pas d'Afne.

408 SYNGENESIA. POLYG. SUECICÆ.

Hab. Utrique in agris argillofis humidis, a viarum excavatarum marginalia, nec no in vineis calcareo argillofis paffim frequen

Flor. Febuario, Martio, Aprili. *Semina* Maj matura. 7

Radix cylindrica, prælonga, longe latequ fub terra reptans, albida, ex una parte folia ex altera flores emittens. *Scapi* digitales, palmares, et altiores, erecti, teretes, ftriati, to mentofi, squamis foliaceis lanceolatis adpreffi præditi. *Folia* radicalia, petiolata, cordat fubrotunda, anguloso-dentata, fuperno viridi fubtus albide tomentofa. *Flos* folitarius, terminalis, radiatus, luteus. *Calyx* cylindricus squamis oblongis acutis, alternis anguftioribus. *Corollulæ* in radio femineæ, ligulatæ, in difc hermaphroditæ tubulofæ: *limbo* quinquefid acuto, reflexo. *Semina* oblonga. *Pappus* fefilis capillaris.

Pharm. TUSSILAGINIS f. FARFARÆ Flores Folia; Radix.

Flores in morbis pituitofis pectoris, tuffi laudantur. Foliorum recentium fuccus, feu decoctum fatum tum contra Scrofulas et ulcera fcrofulofa interdum efficax, Cullen treatife of the mat. med. p. 46.

Ubi in abundantia crefcit femper argillofum aquæ folum indicat, unde aquilegis fignum. Plin. Fig. argillam ad rupes præruptas fluviorum, Linn. Suec. p. 289.

Flores apibus mel præbent.

1276. TUSSILAGO *nivea* fcapo thyrfoides floribus nudis, pedicellis fubramofis, foliis hafate cordatis finuatis, dentati fubtus albiffime-tomentofis. Hoppe. T fchenb. 1803. p. 48.

SYNGENESIA. POLYG. SUPERF. 429

Tussilago paradoxa thyrsis subovato, flosculis femineis nudis multis, hermaphroditis ternis, antheris liberis, foliis triangulari-cordatis denticulatis. *Retz. obs. 2. p. 24. t. 3.*

Tussilago frigida *Hofst. Synops. 457. — Suter. fl. Helv. p. 180.*

Petasites floribus spicatis erectis, foliis calycinis lanceolatis, flosculis paucissimis androgynis. Hall. stirp. helv. n. 141.

Bern. Weissfilziger Huflattig.

Cell. Tussilage blanc de neige.

Hab. In episcopatu (olim) Basileensi circa *Belletai* in locis aquosis humidis t. *Lachenalis* nec non in monte *Jura*.

Flor. Aprili, Majo. *Semina* Majo, Junio matura. 7

Radix horizontalis, repens, cylindrica, albida, fibras teretes longas emittens. *Scapi* palmares et altiores, erectusculi, striato-canaliculati, villosuli, ex viridi-albidi, squamis foliaceis ovali-lanceolatis, præditi. *Folia* radicalia petiolata, hastato-cordata, subtriangularia, sinuato-dentata, subtus tomentosa, nivea. *Flores* scapum terminantes, in thyrsum oblongum pauciflorum abeuntes. *Pedunculi* erecti uniflori, rarius biflori, basi bracteati. *Squamæ* calycinæ lineari-lanceolatae, virides, fusco-villosæ. *Flosculi* feminei nudi, multi, hermaphroditi pauci, terni, omnes tubulosi, calyce longiores.

Nota. Omnino diversa a *C Tussilagine frigida*. *Linn.* ob defectum flosculorum radii ligulætorum.

INGENESIA, POLYG. SUPERF.

ta ex loco citato ab amicissimo Lachenalio ante octodecim jam annos sub T. frigida, in horto culta, mutationi non succubuit.

1277. TUSSILAGO *alba* scapo thyrsoideo, floribus nudis, pedicellis ramosis, foliis orbiculato-cordatis sinuatis dentatis, subtus albido-tomentosis. *Hoppe Taschenb.* 1803. p. 45.

Tussilago alba thyrso fastigiato; flosculis femineis nudis paucis. *Linn Sp. pl.* 2. p. 1214.

Petasites spicis laxis umbellatis, flosculis plerisque androgynis. *Hall. stirp. helv. n.* 134.

Petasites flore albo. *Camerar. epit.* 593.

Icon. Oeder. fl. Dan. t. 524.

Germ. Weisser Huflattig.

Gall. Tussilage blanchatre.

Hab. In ditione Basileensi prope *Muttent* in *Engenthal. t. Lachenalio*; nec non in convallibus *Iuræ* helvetiæ.

Flor. Aprili, Majo. *Semina* Junio matura. 7

Primo intuitu affinis sequenti speciei a qua differt, *Foliis* multo minoribus albide villosis, *Floribus* in thyrsum laxiorem, umbellam fere mentientem digestis, albis, *Pedunculis* longioribus, magis ramosis diffitis, *Bracteis* in basi pedunculorum longioribus, angustioribus, *Calycinis* squamis longioribus acutioribus, *Flosculis* majoribus albis, aut dilute carneis, plerisque androgynis, cum paucis in ambitu femineis.

1278. TUSSILAGO *Petasites* thyrso ovato flosculis femineis nudis paucis. *Linn. Syst. veg. ed.* 14. p. 756. — *Pollich. fl. pal.* n. 7 92.

SYNGENESIA. POLYG. SUPERF. 431

Petasites floribus dense spicatis, flosculis androgynis. *Hall. stirp. helv. n.* 138.

Petasites major et vulgaris. *Bauh. pin.* 197. — *Mapp. fl. Als. p.* 238.

Petasites. *Tabernaemont. p.* 1127. *Icon.*

con. Curtis. fl. Lond. t. 134. — *Regnault bot.* — *Oeder. fl. Dan. t.* 842. — *Bulliard. herb. t.* 391. — *Blackw. t.* 222.

erm. Großblättriger Huflattig. Pestilenzwurz.

zell. Tussilage Petasites.

Hab. Utrinque in pratis humidis, riguis montosis aliisque, praesertim ad rivos et rivulos passim abunde.

Flor. Martio, Aprili. *Semina* Majo, Junio matura. ♂

Radix cylindrica, horizontalis, repens, firmas longas emittens, rubello-albida, magna. **Scapus** spithameus et longior, erectus, teres, villosulus, ex viridi-rubellus vel albidus, simplex, squamis foliaceis lanceolatis praeditus. **Folia** radicalia petiolata, cordata, subrotunda, venosa, inaequaliter sinuato-dentata, dentibususco rubris subcartilagineis, inferne subtomentosa, in planta deflorata amplissima. **Florum** Thyrus ovalis, demum oblongus, subconicus, leniusculus. **Pedunculi** uniflori, basi. **Bracteis** lanceolatis, longitudine pedunculi, suffulti. **Calycinae** squamae lanceolatae, glabrae, apice subincurvae. **Corollulae** pallide purpureae, saepe omnes hermaphroditae, tubulosae, in radio quaeloque 2-3 femineae.

Pharm. *PETASITIDIS Radix.*

Radix acris, aromatica, fragrans alexipharmaca, utredini resistens, hodie fere in medicina neglecta, nunquam efficax.

SYNGENESIA. POLYGA. RUBRA.

Contus bubonibus in peste impongetur.

Flores mellifera.

Ob nimiam saepe abundantiam pratorum pestis.

1279. TUSSILAGO *hybrida* thyris d
longo, flosculis femineis natis plurim
Lin. Sp. pl. 2. p. 1215.

Petalsites floribus spicatis pendulis, petio
multifloris, calycibus aequalibus, flos
lis paucissimis androgynis. *Hall. fl
helv. n. 140.*

Petalsites major, floribus pedunculis lon
infidentibus. *Dill. hort. Eltham. 309
230. f. 297.*

Icon. Schkuhr. bot. Handb. t. 242.

*Germ. Bastard-Huflattig. Bastard-Pestwe
Gall. Petasite-batard. Tussilage-batard.*

Hab. In pratis et ad rivulos passim. In Margg
superiori prope *Schweighof, Oberweiler
Niederweiler*, et *Mullheim am Mühlenbach*
et alibi. Ad *Birjam* et *Wiesam* in prat
frequens.

Flor. Martio, Aprili, Majo. *Semina* Majo,
Junio matura. ♀

A praecedenti specie facile distinguitur
Scapis altioribus, tenuioribus, *Squamis* in scape
foliaceis linearibus, angustioribus, *Thyrso* lon
giori, densiori. *Pedunculis* tenuioribus, copio
sioribus, *Bracteis* angustioribus, lineari-setaceis
Floribus minoribus, *Flosculis* femineis pluribus
paucissimis androgynis.

Planta in horto botanico cum praecedenti speci
cultata mutationi non succubuit.

Radix fragrans ut praecedens.

CCCLXXXVI.

CCCLXXVI. SENE C I O.

Linn. Gen. plant. ed. Schreb. n. 1290.

Calyx communis calyculatus, conicus, truncatus: *squamis* subulatis plurimis, in cylindrum perne contractum parallelis, contiguis, æqualibus, paucioribus basin imbricatim tegentibus, micibus emortuis (sphacelatis.) *Corolla* composita, calyce altior: *Corollulae* hermaphrodite, tubulosæ, numerosæ, in disco. *Femineae* subulatae, in radio, si quæ adsint. *Propria* hermaphroditis infundibuliformis: *Limbo* reflexo, quinquefido. *Femineis* (si quæ) oblonga, obsolete tridentata. *Calyx* fructiferus conico-concavens. *Semina* ovata. *Pappus* capillaris, longus. *Receptaculum* nudum, planum.

† *Floribus flosculosis.*

80. *SENECIO vulgaris* corollis nudis, foliis pinnato-linuatis amplexicaulibus, floribus sparsis. *Linn. Sp. pl. 2. p. 1216.* — *Pollich. fl. pal. n. 793.* — *Hall. stirp. helv. n. 58.*

Senecio minor vulgaris. Bauh. pin. 131. — *Mapp. fl. Alf. p. 285.*

Senecio I. II. Tabernæmont. p. 462. Icon. con. Curtis. fl. Lond. — *Bulliard. herb. t. 197.* — *Regnault. bot.* — *Blackw. t. 132.*

erm. Gemeines Kreuzkraut. Gelbes Vogelkraut.

Hall. Senecion commune.

Hab. Utrique in hortis, agris, ruderatis, vineis et muris, frequens.

Flor. Per totum fere annum, in hyemem.

Caulis palmaris, pedalis et altior, erectus subangulosus, ramosus. *Folia* alterna, sessilibus amplexicaulia, pinnatifida, sinuato-denticulata glabra, saturate viridia. *Flores* caulem ramisque terminantes, lutei, subcylindrici, non radiati, pedunculati. *Pedunculi* striati, uniflori. *Calycinae* squamae apice nigrae. *Corollulae* omnes tubulosae. *Semina* striata, fusca.

Pharm. *SENECIONIS Herba.*

Hodie in medicina exoleta.

Herba et seminibus delectantur Passeres canori.

Succus expressus in Anglia equis, a vermibus infestatis ingeritur. Raj. hist. pl. 1. p. 290.

Flores melliferi.

†† *Floribus radiatis: Radio revoluto.*

1281. *SENECIO viscosus* corollis revolutis, foliis pinnatifidis viscidis, squamis calycinis laxis longitudine perianthii. *Linn. Sp. pl.* 2. p. 1217. — *Poll. fl. pal.* n. 794.

Senecio foliis viscidis pinnatis, pinnis semipinnatis angulosis, calycibus hirsutis. *Hall. stirp. helv.* n. 60.

Senecio hirsutus viscidus, graveolens. *Dillen. hort. Eltham.* 347.

Senecio incanus pinguis. *Bauh. pin.* 131.

Jacobaea pannonica I. *Clus. hist.* 2. p. 21. — *Mapp. fl. Alf.* p. 152.

SYNGENESIA. POLYG. SUPERF. 236

Senecio tomentosa III. *Tabern. p.* 463. *Icon. rudis.*

non. Dillen. l. c. t. 258. *f.* 336.

erm. Klebriges Kreuzkraut.

all. Seneçon visqueux.

ab. Utrisque in sylvis cæduis et ad carbonariorum areas passim abunde, nec non in ruderatis, hinc inde.

or. Julio, Augusto. Semina Augusto, Septembri matura. ☉

Caulis cubitalis, tripedalis, erectus, teres, iatus, viscido-pubescent, ramosus. *Rami* terni, subdichotomi. *Folia* alterna, sessilia, amplexicaulia, pinnatifida, segmentis obtusis, læte viridia, utrinque villosa viscida. *ores* caulem ramosque terminantes, radiati, illide lutei, pedunculati. *Pedunculi* erecti, illori, bracteolati, villosa-viscidi. *Flosculi* triiligulati. *Ligulae* primum patentes, demum revolutæ. *Calyces* laxi, villosa-glandulosi, villi. *Squamæ* calycinæ 15-20, apice nigricantes.

282. SENECIO *sylvaticus* corollis revolutis, foliis pinnatifidis, denticulatis, caule corymbose erecto. *Linn. Sp. pl.* 2. p. 1217. — *Poll. fl. pal. n.* 795.

Senecio foliis pinnatis, pinnis alterne majoribus semipinnatis, angulosis, calycibus lævibus, radiis brevissimis revolutis. *Hall. stirp. helv. n.* 59.

Senecio minor latiore folio s. montanus. *Bauh. pin.* 131.

non. Oeder. fl. Dan. t. 869. — *Dill. hort. Eltham. t.* 258. *f.* 337.

Germ. Wald-Kreuzkraut.

Gall. Seneçon des forêts.

Hab. In sylvis cæduis apricis passim frequenter nec non in arvis macilentis apricis sabulosis et ruderatis.

Flor. Junio Julio, Augusto. *Semina* Augusto Septembri matura. ☉

Diffinguitur a præcedente specie affinis *Caule* plerumque graciliore, villosa, non viscido, *Foliis* non viscidis, *Floribus* copiosis minoribus. *Calycinis* squamis non viscidis numero paucioribus 12-13, rarius 15, minoribus.

Flores melliferi.

Arborum cotyledones contra folis radios et ventorum injurias cum præcedente specie defendit.

††† *Floribus* radiatis: radio patenti, foliis pinnatifidis.

1283. *SENECIO erucifolius* corollis radiantibus, foliis pinnatifidis dentatis subhirsutis, caule erecto. Linn. Sp. pl. 1. 1.

quens. Inter *Durlach* et *Aue* ad fossas aqua-
ticas viam versus.

lor. Julio, Augusto. *Semina* Augusto, Sep-
tembri matura. ¶

Distinguitur a *S. Jacobaea* cum qua *Hälle-*
s conjunxit, *Caule* altiore 3-4 pedali, magis
ro, hirtō, *Foliorum* pinnis gracilioribus,
btus hirsutis, *Floribus* pallide sulphureis, *Ca-*
ribus basi tomentoso-villosis: *squamulis* apice
bellis aut fuscis,

Nota. Icon Curtisii plantam nostram egregie re-
præsentat, descriptio autem ejusdem et Polli-
chii omnino magis ad *S. tenuifolium* Jac-
quini quadrat.

284. *SENECIO tenuifolius* corollis ra-
diantibus, foliis omnibus erectis incanis
hirsutis pinnatis: pinnis linearī-subulatis,
pedunculis corymbosis, caule erecto sim-
pliciusculo incano tomentoso.

Senecio tenuifolius corollis radiantibus, fo-
liis omnibus pinnatis, pinnis linearibus,
pedunculis corymbosis, caule erecto.
Murr. Linn. Syst. veg. ed. 14. p. 758.

on. *Jacqu. fl. Austr. 3. t. 278.*

erm. *Dünnblättriges - Kreuzkraut.*

all. *Senecion à feuilles menues.*

ab. In montosis apricis, præsertim in colli-
bus calcareo-argillaceis dumetosis asperis,
ad vineas montis *Thurnberg*, prope *Grö-*
zingen, *Berghausen*, circa *Bruchsal* et ali-
bi non infrequens.

lor. Augusto, Septembri, Octobri. *Semina*
Octobri, Novembri matura. ¶

SYNGENESIA. POLYG. 30822.

Distinguitur a præcedente specis, *Habitu* frictiori, *Caule* simpliciore, tenuiore, densius folioso, albide tomentoso, aut villosa, in capulis frictioribus, erectis, fere cauli adpressius profundissime pinnatifidis, fere pinnatis, pinnulis exacte linearibus, margine revolutis, in griseis, multo semper angustioribus, hemioribus superne hirtis, late viridibus, subtus villosis incanis, *Floribus* corymbosis, subminoribus, *lycum* squamis minoribus, viridibus, apice alteratis.

Nota. Jacquiniana Icon, plantam nostram me repræsentat, omnino diversam a præcedente cum qua sæpissime confunditur. Hæc amat collina, sicca aspera, illa sequitur loca densius humida, marginalia ad aquarum fossas.

1285. **SENECIO** *Jacobaea* corollis radiatis, laciniatis, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. *Linn. Sp. pl.* p. 1219. — *Poll. fl. Dan. n.* 797.

Senecio foliis lineatis pinnatis, pinnis dentatis, extrema maxima. *Hall. stirp. helv.* n. 62.

Jacobæa vulgaris laciniata. *Bauh. pin.* 13. *Mapp. fl. Alf.* p. 151.

Herba *Jacobaea* I. *Tabern. p.* 465. *Icon.*

β. *Jacobaea* vulgaris laciniata, flore discoidi. *Vaill. act.* 1720. p. 383.

Senecio *Jacobæa* β. flore discoide, absque radio. *Poll. l. c.*

Icon. Oeder. fl. Dan. t. 944.

Germ. Jacobskraut.

Gall. Senecion Jacobée. Herbe de St. Jacques.

Hab. Utrunque in pratis udis, sepibus, agri muris passim frequens.

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. 7

Caulis cubitalis, tripedalis, erectus, angustus, striatus, glabriusculus, ramosus. *Folia* alterna, amplexicaulia, pinnatifida, laciniata, aequaliter dentata, obtusa, lacinia extrema minore, glabra, late viridia. *Flores* terminales, symphylae, radiati, lutei. *Calycinae* squamae pice nigrae.

Variationem β . in Belgiae dunis vulgarem, hucusque apud nos non vidi, et unicum saltem exemplar in herbis circa Lautern vidit Pollichius. l. c.

Pharm. JACOBÆAE Herba.

Hodie fere in medicina exoleta, amariuscula, adstringens, olim in dysenteria maxime commendata.

Flores melliferi.

286. *SENECIO aquaticus* radio patente, flosculis ellipticis, foliis lyratis ferratis, inferioribus obovatis integris, feminibus glabris. *Smith. fl. brit.* 2. p. 885. — *Willd. Linn. Sp. pl.* 3. p. 1997. — *Lamarck et Decand. fl. franc.* 4. p. 163.

Senecio aquaticus corollis radiantibus, foliis dentatis, radicalibus ovatis, caulinis pinnatifidis: lacinia extrema majore. *Hudsf. fl. angl.* 2. p. 366.

Jacobaea latifolia. *Bauh. hist.* 2. p. 1057.

Jacobaea tertia latifolia. *Clus. hist.* 2. p. 43. *Icon.*

non. *Oeder. fl. Dan.* t. 784.

erm. *Wasser-Jacobskraut. Wasser-Kreutzkraut.*

all. *Senecion aquatique.*

440 SYNGENESIA. POLYG. SUPERT.

Hab. Cum præcedente præsertim in pratis missioribus aquosis cis et transrhenanis. In Principatu superiore in pratis udis ad L. cum Bodamicum frequens.

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. 7

Affinis omnino præcedenti specie, a qua differe videtur, *Foliis* radicalibus ovatis, tripliciusculis, integriusculis, superioribus simpliciusculis; *laciniis* extima majore, *Calyce* hemisphærico, *Seminibus* glabris.

Nota. Ob variationes inter hanc et Jacobaeam intermedias, rectius forsan pro variatione S. Jacobaeae aestimanda.

Flores apibus melliferis.

††† *Floribus* radiatis: radio patente, *Foliis* indivisis.

1287. *SENECIO paludosus* corollis radiantibus, foliis ensiformibus acute serratis subtus subvillosis, caule stricto. *Lin. Sp. pl.* 2. p. 1220. — *Poll. fl. pal.* n. 798.

Senecio foliis linearibus, serratis, floribus umbellatis. *Hall. stirp. helv.* n. 66.

Jacobaea palustris altissima, foliis serratis. *Tournef. — Mapp. fl. Als.* p. 152.

Conyza palustris ferratifolia. *Bauh. pin.* 166.

Consolida palustris III. *Tabernæmont. p.* 946. *Icon.*

Icon. *Oeder. fl. Dan.* t. 385.

Germ. Sumpf-Kreutzkraut.

Gall. *Senecion des marais.*

b. Utrunque in pratis udis, fossis inter salices. Prope *Dachsland*, *Knietingen*, *Eckenstein*, *Rufsheim* et alibi passim non infrequens. In Principatu superiore in pascuis udis ad lacum Bodamicum prope *Roschach* et *Arbon* abunde.

r. Julio, Augusto. *Semina* Septembri maturata. 7.

Caulis 2-3-5 pedalis, erectus, teres, fulvus, simpliciusculus, glabriusculus aut villosus.

Folia alterna, sessilia, erecta, ensiformi-lanceolata, acuminata, argute serrata, superne viridia, glabra, subtus incana, aut albidomentosa. *Flores* caulem terminantes, umbellati-corymbosi, radiati, lutei, majusculi. *Pediculi* ramosi, subtus foliolo lanceolato, involucro, suffulti. *Calyx* cylindricus, *squamis* liri-lanceolatis glabris, tomentosulisque, apicibus.

Individa ad lacum Bodamicum lecta, foliis subtus tomentoso-niveis instructa sunt, alia in Rheni vicinia lecta, subtus fere glabra.

nota. Non confundenda cum *Cineraria palustri* Linn. foliis lato-lanceolatis, profunde dentato-sinuatis, non ensiformi-anguste-lanceolatis, argutissime serrulatis.

88. *SENECIO nemorensis* corollis radiantibus octonis, foliis lanceolatis biferratis subtus villosis, caule ramoso. *Linn. Sp. pl. 2. p. 1221.* — *Pollich. fl. pal. n. 799.*

Senecio foliis ovato-lanceolatis, ferratis subtus hirsutis, floribus umbellatis. *Hall, stirp. helv. n. 64.*

Iacobæa nemorensis latiore rigidiore et hirsuto folio. *Rupp. fl. Jen. 142.*

caulis, erectus, subangulosus, villo re-
 fus, hinc scabriusculus, superne ramof-
 radicalia in caespitem congesta, ovata
 in petiolum superne concaviusculum c-
 tia, margine denticulata; *caulina* alter-
 tuor quinque inferiora quoque denticu-
 ceolata, semiamplexicaulia, summa
 ma, supra villosiuscula ex albicante viri-
 tus alba et tomentosa. *Flores* ad cauli-
 tatem quasi in umbellam irregularem.
Pedunculi subpubescentes, singuli fo-
 ceolato interstincti. *Calyx* cylindricus,
 parallelis linearibus villosiusculis, apice
Corollae radiatae ubique flavæ; *Ligu-*
 multo longiores, tredecim, duodecim
 tridentatæ. *Semina* pappo sessili corol-
Pollichius.

Nota. Pollichii plantam non vidi.

Obs. Senecio nemorensis Jacquini f
 t 184. vix est nemorensis Linnae

Folia in Jacquini planta, quam ex Auf-
 oculos habeo, utrinque glabriuscula, margi-
 serrata, sed simplicissime, et aequaliter cren-
 culata. cartilaginea. Flores in individuo u

SYNGENESIA. POLYG. SUPERF. 443

Pollichii planta omnino diversissima a Jacquiniana ut facile ex ejus descriptione patet. Forsan novae species in loco natali adhuc inquirenda.

189. SENECIO sarracenicus corollis radiantibus, floribus corymbosis foliis lanceolatis ferratis glabriusculis. *Linn. Sp. pl. 2. p. 1221. — Pollich. fl. pal. n. 800.*

Senecio foliis lanceolatis, ferratis, glabris, floribus umbellatis. *Hall. stirp. helv. n. 65.*

Virga aurea latifolia, ferrata. *Bauh. pin. 268. Mapp. fl. Als. p. 327.*

Virga aurea five Solidago Sarraceniça, latifolia ferrata. *Bauh. hist. 2. p. 1063. omnino!*

non. Jacqu. fl. Austr. t. 186.

erm. Heidnisch-Kreutzkraut.

all. Seneçon Sarracin.

ab. Utrunque in sylvis montosis umbrosis humidis alperis passim abunde. In montosis sylvis retro *Ettlingen*, prope *Frauenalb*, *Herrnalb*, *auf dem Kaltenbrunn*; in sylvis prope *Oppenau*, *Griesbach*, *Petersthal*, frequens, et in toto tractu *Sylvae nigrae* passim abunde. In *Alsatiae* *Vogesis* frequens.

lor. Julio, Augusto. *Semina* Augusto, Septembri matura. **7**

Radix repens. *Caulis* 4-5 pedalis, erectus, sulcatus, fulcatus, glaber, foliosus, simpliculus, superne corymboso-ramosus. *Folia* versa, alterna, infima petiolata, superiora subsilia, oblonga, lanceolata, lata, acuta, vesa, utrinque glabra, superne saturate viridia, consplendentia, subtus pallide viridia, margine

longa, striata, fulca. *Pappus* capillaris.

Variat interdum in uno eodemque individuo margine simpliciter et biferrato-dentatis.

Pharm. CONSOLIDAE SARRACENICAE.

Planta adstringens, vulneraria non cum *S. Virga aurea* confundenda.

Flores apibus melliferi.

1290. *SENECIO Fuchsi* corollis
tibus, floribus corymbosis, foliis
lanceolatis acutis, argute dentatis,
utrinque glabris, petiolatis.

Senecio ovatus corollis radiantibus
ovato-lanceolatis argute ferratis
que glabris subsessilibus, *Willden*
Sp. pl. 3. p. 2004. an?

Jacobaea ovata. *Flora der Wettera*
212. an? videtur.

Jacobaea alpina foliis longioribus
Mapp. fl. Alf. p. 151.

Solidago Sarracenicæ, *Fuchs. hist.*

b. Utrisque in nemoribus ficioribus, ad sylvarum marginalia passim frequens. Circa *Carlsruhe im Phasanengarten, im Beurtheimer Wald*; prope *Mühlburg im Lutherischen Wäldlein* et alibi in sylvis Rheno vicinis frequens.

Julio, Augusto. *Semina* Septembri matuta. 7

Caulis tripedalis et altior, erectus, angustis, purpureo-rubellus, subflexuosus, foliis, superne corymbo-ramosus. *Folia* caulisparsa, alterna, breviter petiolata, ovaliceolata, utrinque glabra, laete viridia, firmiuscula, margine utrinque infra apicemque duplicato aut simpliciter minute denticulato-ferrata, in ferraturis apice cartilagineis chirsuta, tres uncias vix longa, novem lineas unciam vix lata. *Flores* terminales corymbosi. *Corymbi* erecti. *Pedunculi* aut ramuli alni superne bifidi, nudi, pedicelli uniflori, acteis fetaceis, angustissimis, erectis, praeteri. *Calycinae* squamae 9-10, lineares, striatae, apice nigrae, glabriusculae. *Bractae* sublyce, 5-6, fetaceae, erectae, calyce breviores. *Corollulae* radii 5-7 ligulati. *Ligulae* lineares, obtusae, quatuor lineis longitudinalibus rufae, luteae. *Corollulae* disci infundibuliformes, luteae.

Differt omnino a praecedente specie, Caule deflori, purpureo-rubello, Foliis omnibus breviter petiolatis ovali-lanceolatis, firmioribus, multo brevioribus et angustioribus, Corymbo florum minus usque, Loco natali sicco, aprico.

Nota. Synonyma veterum allegata, omnino ad hanc plantam nec ad *S. Sarracenicum* nostrum spectant. Icon Fuchsi quoad formam et habitum plantae egregia.

446 SYNGENESIA. POLYG. SUPERF.

Obs. Hanc plantam etiam in Gallia, Helvetia et Franconia vidi, ubi cum *S. nemorensi* et *S. racenico* saepe confunditur.


Diversa omnino a *S. nemorensi* Jacqu. Auftr. t. 184.

Planta in horto botanico culta, mutationi non subiecit. Flores non suaveolentes.

CCCLXXXVII. A S T E R.

Linn. Gen. pl. ed. Schreb. n. 1291.

Calyx communis imbricatus: squamis interioribus apice prominulis, inferioribus patulis. Corolla composita, radiata. Corollulae hermaphroditæ numerosæ in disco. Femineae ligulatae in radio. Propria hermaphroditi infundibuliformis: limbo quinquefido patulo. Femineae ligulata, lanceolata, tridentata. Calyx fructifer vix mutatus. Semina oblonga, ovata. Pappus capillaris. Receptaculum nudum, planiusculum.



non. Blackw. herb. t. 109.

erm. Blaues Sternkraut.

all. Aster Amellus. Oeil de Christ.

ab. Utrunque in collibus apricis graminosis asperis. In monte Thurnberg, prope Grözingen, Weingarten, copiose. In Brisgovia ad montem Kaiserstuhl in collibus vulcanicis prope Rothweil et alibi frequens.

lor. Augusto, Septembri. Semina Octobri matura. 7.

Caulis pedalis, sesquipedalis, erectus, teres, minutus, foliosus. Folia alterna, inferiora pedunculata, ovali-lanceolata, obtusiuscula, integerrima; superiora sessilia, integra, omnia subserrata, hirsuta, scabriuscula. Flores caulem terminantes, corymbosi. Calycinae squamae breviter, patentes, apice rubellae. Corollulae disciformes. Radii ligulae numerosae, patentes majusculae, pulcherrime coeruleo-violaceae, rarius albae.

erm. ASTERIS ATTICI Herba.

Amariuscule, subaromatica, adstringens, hodie in officina exoleta.

Flores apibus mel praebent.

Egregium in collibus graminosis decus.

Est etiam flos in pratis cui nomen Amello. Virg.

nota. Plures Asteris species americanae in parte sylvatica amoenissima am Steinbacher Baad confitae sunt, inter quas praecipue Aster cordifolius sese multiplicavit. Aster chinensis cum multifariis variationibus pulcherrimis passim in hortis colitur.

92. *ASTER annuus* foliis ovatis: inferioribus crenatis, caule corymboso, pedunculis nudis, calycibus hemisphaericis.

Linn. Sp. pl. 2. p. 1229. — *Lamar Decand. fl. franc.* 4. p. 146. — *Roiger. germ.* 2. p. 357.

Aster annuus ramosus albus latifolius densis. *Morif. hist.* 3. p. 122.

Bellis ramosa umbellifera. *Cornuti cana.* t. 194.

Icon. Oeder. fl. Dan. t. 486.

Germ. Einjähriges weisses Sternkraut.

Gall. Aster blanc annuelle.

Hab. In pratis et sylvaticis umbrosis graminibus et transrhenanis passim non infrequenter. In pratis udis ad Rhenum circa *Dachsniedern, Knielingen, Schrök, Linkenheim* et cum Allio anguloso frequens. Prope *Edelsheim* in nemorosis. In Marggraviam superiore prope *Zienken*. In Brisgovia *Kaiserstuhl* nuper vidi cum *Chlora pinnatifida* *Linn.*

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ☉

Caulis sesquipedalis, cubitalis et alius erectus, sulcatus, striatus, hirtus, læte viridis superne divifus. *Folia* alterna, inferiora palmata, ovata, sinuato-crenata; superiora sessilia, ovato-lanceolata, integrifolia, ciliolataque subcrenulata. *Flores* terminales, compositae, quasi in umbellam congestae, erectae. *Corollae* disci luteae; *Radiales* ligulati, albi, patuli. *Bellidis* florem imitantes. *Calyx* hemisphaericus.

Planta elegans saepe in Januarium usque apud nos florens.

Flores apibus melliferi.

CCCLXXXVIII. SOLIDAGO.

Linn. Gen. pl. ed. Schreb. n. 1292.

Calyx communis oblongus, imbricatus: *nis* oblongis, angustis, acuminatis, rectis, inventibus. *Corolla* composita, rariata. *Illulae* hermaphroditæ tubulosæ, plurimæasco. *Femineae* ligulatae in radio, paucae. *ria* hermaphroditi infundibuliformis: *Lim* inquefido, patulo. *Feminea* ligulata, lan-
ta, tridentata. *Calyx* fructiferus vix muta-
Semina obovata, oblonga. *Pappus* capiti
Receptaculum nudum, planiusculum.

3. *SOLIDAGO Virgaurea* caule sub-
flexuoso angulato, racemis paniculatis
erectis confertis. *Linn. Sp. pl. 2. p. 1235.*
— *Pollich. fl. pal. n. 802.*

olidago caule angulato hirsuto, foliis ovato
lanceolatis dentatis, racemis multifloris
congestis. *Hall. stirp. helv. n. 69.*

irga aurea vulgaris latifolia. *Bauh. hist. 2.*
p. 1062, — Mapp. fl. Alf. p. 328.

irga aurea ferratis foliis. *Tabernæm. p. 1260.*
Icon.

1. *Oeder. fl. Dan. t. 663. — Regnault. bot.*
Blackw. herb. t. 169.

m. Goldruthe. Heidnisch - Wundkraut.
Goldwundkraut.

. Verge d'or commune.

. Utrique in dumetis elevatis, sylvis, et
collibus dumetosis asperis. Amat sylvarum
marginalia, præcipitia dumetosa virgultosa.

450 SYNGENESIA. POLYG. SUPERF.

Flor. Augusto, Septembri, Octobri. *Semina* Octobri, Novembri matura. 7

Radix ramoso-fibrosa, extus fusca, intus albida. *Caulis* cubitalis, tripedalis et altior, erectus, sulcatus, glabriusculus. *Folia* alternis inferiora petiolata, ovato-lanceolata, reliqua sessilia, sub lanceolata, omnia serrata, anteriores versus saepissime recurviuscula, duriuscula, glabriuscula, quandoque utrinque hirsutula. *Flora* aurei, paniculato-racemosi, axillares. *Racemi* terni, erecti, densi, multiflori.

Variat. Caulibus plus minusve ramosis.

Pharm. VIRGAE AUREAE Herba.

Planta amara, adstringens, vulneraria, ad ulcera mundanda commendata.

Solidaginis seu Consolidae Saracenicae herba officinis pharmaceuticis hinc inde subministrat.

Flores melliferi.

Nota. Solidago canadensis alicujusque species virginianae passim in hortis coluntur.

CCCLXXXIX. CINERARIA.

Linn. Gen. plant. ed. Schreb. n. 1294.

Calyx communis simplex, polyphyllus, foliolis aequalibus. *Corolla* composita, radiata. *Corollulae* hermaphroditae, aequales, numerosae, in disco. *Femineae* ligulae, numerosae, litorum calycis in radio. *Propria* hermaphroditi infundibuliformis: *Limbo* quinquefido, erecto. *Feminea* ligulata, lanceolata, apice denticulata. *Calyx* fructiferus immutatus. *Semina* linearis quadrangula. *Pappus* capillaris, copiosus. *Receptaculum* nudum, planiusculum.

94. *CINERARIA palustris* floribus corymbosis, foliis lato-lanceolatis dentato-finuatis, caule villoso. *Linn Sp. pl.* 2. p. 1243. — *Lamarck et Decand. fl. franc.* 4. p. 169.

Jacobæa aquatica elatior foliis magis dissectis. *Morif. hist.* 3. p. 110. *sect.* 7. *t.* 19. *f.* 24.

Zonysa aquatica laciniata. *Bauh. pin.* 266.

Zonysa Helenitis foliis laciniatis. *Lobel. Icon.* 347. *bona.*

n. *Oeder. fl. Dan.* *t.* 573.

m. *Sumpf-Aschenkraut.*

l. *Cinéraire des marais.*

b. In ditione Basileensi prope *Michelfelden* in pascuis aquosis in *der Hardt*, ubi legit *Zeyherus.*

•. Junio; Julio. *Semina* Augusto matura. *♂*

Radix fibrosa. *Caulis* pedalis, sesquipedalis us, crassiusculus, striatus, villosus, a basi ad foliosus, simplex. *Folia* radicalia in petiolecurrentia, lato-lanceolata, profunde inciso-nuato-ferrata, fere pinnatifida, novem circuncias longa, ad tres fere uncias lata, vel villosiuscula, læte viridia; *caulina* sparsa, na, sessilia, semiamplexicaulia, ovali-lanceolata, erecta, grosse et inæqualiter ferrata, uncias longa, septem lineas ad unciam læte viridia, subvillosa. *Flores* caulem inantes, densissime corymbosi, pallide lutei. *anuli* villosi-lanuginosi, incani. *Calycinae* lanceolatae, acuminatae, villosae. *Corollæ* radii ligulatae. *Ligulae* patentes, pallide lustridentatae; *Disci* infundibuliformes, cohes. *Semina* decem-angulata, angulis al-

452 SYNGENESIA. POLYG. SUPER

ternis eminentioribus. *Pappus* per lente
mosus.

Obs. Villi prælongi in pedunculis et calycib
phani, articulati.

Nota. Ad rarissimos florae cives pertinet; h
in Helvetia non visa.

Lobellii Icon plantam nostram optime e

1295. *CINERARIA cordifolia* fol
tiolatis cordatis duplicato-dentatis
fis subtus subvillofis incanis, pedi
incrassatis albide tomentosis un
caule angulato.

Cineraria cordifolia. Linn. *Suppl.* 375.
den. Linn. *Sp. pl.* 3. p. 2083. — *La*
et Decand. fl. franc. 4. p. 171.

Cineraria cordifolia foliis cordatis dup
dentatis, subtus incanis, floribus
bolis, pedunculis foliosis. *Gouan*
p. 69.

Senecio foliis cordato-lanceolatis f
Hall. stirp. helv. n. 63.

Jacobæa alpina foliis subrotundis f
Bauh. pin. 131. *Prodr.* p. 69. *Icon l*

Icon. Jacqu. *fl. Austr.* 2. t. 176.

Germ. Herzblättriges Aschenkraut.

Gall. Cineraire a févilles en coeur.

Hab. In Principatu superiore ad lacum
micum prope *Argon* et *Roschach*
cum *Senecione* paludoso frequens
non prope *St. Gallen*.

Flor. Junio, Julio. *Semina* Augusto ma

Radix fibrosa. *Caulis* pedalis, sesqui
lis, erectus, angulatus, striatus, crassu

aber, aut villoso-incanus, foliosus, simplicifolius. *Folia* caulina alterna, petiolata, cordata, ovata, venosa, superne laete viridia, subtus villoso-incana, quandoque glabriuscula, medio incano destituta, margine grosse duplicato-crenata, circiter tres uncias longa, binas uncias lata, plana. *Petioli* superne canaliculati, albide tomentosuli, basi sæpe nudi, quandoque canaliculati: *auriculis* oppositis ovali lanceolatis, parvis. *Flores* caulem terminantes, subcorymbosi, majusculi, lutei. *Pedunculi* circiter 2, alterni, erecti, incrassati, albide tomentosi, *bracteis* longis setaceis suffulti. *Corollulae* ligulatae, patentes; *Disci* infundibuliformes, concolores.

296. *CINERARIA lanceolata* foliis radicalibus ovalibus, caulinis sessilibus linearibus lanceolatis omnibus integris, tomentosis, caule simplici, floribus umbellatis.

Cineraria campestris floribus corymbosis, foliis radicalibus petiolatis ovatis tomentosis, caulinis sessilibus linearibus. *Retz. prodr. fl. Scand. ed. 2. n. 1027.* — *Hofst. Synops. 464.* — *Willden. Linn. Sp. pl. 3. p. 2081.* — *Lamarck et Decand. fl. franc. 4. p. 169.*

Cineraria integrifolia β. *Villars. fl. Delph. 3. p. 225.*

Cineraria integrifolia pratensis. *Jacqu. fl. Austr. p. 48.*

Cineraria lanceolata. *Lamarck. fl. franc. p. 125.*

Jacobæa montana, lanuginosa, angustifolia, non laciniata. *Bauh. pin. 131.* *Mapp. fl. Alf. p. 151.*

Jacobæa pannonica secunda. Clus. hist. p. 22.

Icon. Jacqu. fl. Austr. 2. t. 180.

Germ. Lanzetblättriges Aschenkraut.

Gall. Cinéraire des champs.

Hab. In Sponhemia prope *Winningen* ad Mofellam in prato montoso sylvatico udo non infrequens cum sequente specie vidi an 1786. In Alsatia versus *Hochfeld im hohen Wald. t. Mappo.*

Flor. Majo, Junio, *Semina* Julio, Augusto maturata. ♂

Radix fibrosa. *Caulis* sesquipedalis, vix altior, erectus, striatus, sublanuginosus, simplex. *Folia* radicalia in petiolum brevem decurrentia, ovalia, obtusiuscula, integra, *caulina* alternata, sessilia, lineari-lanceolata, plus minusve tomentosa, integerrima. *Flores* caulem terminantes, umbellati, lutei, majusculi. *Pedunculi* seu radii 5-7, breviusculi, crassiusculi, tomentosi, albidii, basi squamula suffulti, uniflori. *Corollulae* radii ligulatae, patentes, integrae, luteae; *Disci* infundibuliformes concolores. *Pappus* per lentem plumosus.

Distinguitur a sequenti affini, Caule demissiore, Foliis radicalibus caulinisque infimis non spathulatis, dentatis, Pedunculis brevioribus.

Obs. Hanc speciem nunquam in campestribus sed in pratis subalpinis in Helvetia vidi, hinc nomen mutavi.

Nota. Ad rariores Florae cives pertinet.

1297. *CINERARIA spathulaefolia* foliis radicalibus caulinisque infimis in petiolum membranaceum decurrentibus, spathulatis, obtusis, denticato-crenatis, superioribus sessilibus lanceolatis, inte-

griusculis, tomentosulis, caule simpliciter
floribus umbellatis.

Cineraria integrifolia floribus umbellatis,
foliis denticulatis lanceolatis, radicalibus
spathulatis. *Murr. Syst. veg. ed. 14. p. 764.*
— *Hoff. Synops. 464.* — *Willden. Linn.*
Sp. pl. 3. p. 2082. — *Lamarck et Decand.*
fl. franc. 4. p. 170.

Cineraria integrifolia alpina. *Jacqu. fl. Austr.*
p. 48.

con. *Jacqu. fl. Austr. t. 179.*

erm. Spatelblättriges Aschenkraut.

a ll. Cinaire a sévilles d'une Spathule.

a b. In Sponhemia prope *Winningen* ad Mo-
sellam in prato sylvatico udo non infrequens.

er. Majo, Junio. Semina Julio matura. ♂

Distinguitur a præcedente specie, *Caule*
tiore 2-3 pedali. Foliis radicalibus caulinis in-
mis in petiolum longum, membranaceum
beuntibus, exacte spathulatis, obtusioribus,
margine crenato-denticulatis, *Floribus* subma-
ioribus, *Pedunculis* longioribus, medio saepe
revissimo.

b c. Individua loco citato lecta, helveticis subalpi-
nis simillima, nisi quod minus illa lanuginosa.

Nota. Ad ratiore Florae cives omnino pertinet.

Haec species omnino folia non habet integra,
ne triviale nomen mutandum existimavi.

CCCXC. I N U L A.

Linn. Gen. pl. ed Schreb. n. 1295.

Calyx communis imbricatus: foliolis laxis,
stulis: exterioribus majoribus, aequalis lon-
tudinis. Corolla composita, lata. Corollulae

456 SYNGENESIA. POLYG. SUPERF.

hermaphroditæ aequales, numerosissimæ in disc
Femineæ ligulatæ, numerosæ, confertæ, in
 dio. *Propria* hermaphroditi infundibuliform
limbo quinquefido erectiusculo. *Femineæ* lig
 lata, linearis, integerrima. *Antheræ* inferne
 setas decem desinentes. *Calyx* fructiferus imm
 tatus. *Semina* linearia. *Pappus* capillaris lon
 gitudine seminis. *Receptaculum* nudum, planum

1298. INULA *Helenium* foliis amplex
 caulibus, ovatis, rugosis subtus toment
 tosis, calycum squamis ovatis. *Linn. S*
pl. 2. p. 1236.

Aster foliis ovato-lanceolatis ferratis subt
 tomentosis, calycinis ovato-lanceolatis
 maximis. *Hall. stirp. helv. n. 72.*

Aster *Helenium*. *Scopol. fl. Carn. ed 2.*
 1078.

Aster omnium maximus, *Helenium* dictum
Mapp. fl. Alf. p. 33.

Helenium vulgare. *Bauh. pin. 267.*

Helenium. *Tabernæmont. p. 953. Icon.*

Icon. Oeder. fl. Dan. t. 728. — Regnault, bot.
— Blackw. herb. t. 473.

Germ. Alant. Alantwurz. Oltwurz.

Gall. l'Aunée. Inule Hélélière.

Hab. Ad pagorum et urbium passim sepes
 rudrata infrequens. Prope *Durrbach* in
dem Staufenberg nuper copiose nascentes
 vidi. In Alsatia circa Argentinam; prope
Zabern, inter *Niederbronn* versus *Rhenus*
 copiose in agris t. *Mappo.*

lor. Julio, Augusto. *Semina* Septembri mat-
tura. ¶

Radix ramosa, majuscula, albida, fibris
formibus albidis obfessa. *Caulis* 3-4-6 peda-
lus, erectus, crassiusculus, validus, angulosus,
futuræ scaber, ramosus. *Folia* radicalia petio-
la, magna, ovata, obtusa; *caulina* alterna,
plexicaulia, ovato-lanceolata, integra, cras-
sacula, scabra, subtus tomentosa. *Flores* ter-
minales, magni, lutei. *Calycinae* squamæ solia-
re, ovali-lanceolatæ, præmagnæ, asperæ.

herm. ENULÆ CAMPANÆ *f. HELENII* *Radix.*

Radix amara, aromatica, acris, viribus roboran-
s, resolventibus, tonicis.

herm. In ovium scabie efficax.

herm. Passim etiam in rusticorum hortis colitur. Dome-
stica medicina: *Radix* vino generoso patrio infusa, mel-
addito, in raucedine cum tusci juncta.

Flores apibus mel præbent.

99. *INULA Britannica* foliis amplexi-
caulibus lanceolatis distinctis serratis sub-
tus villosis, caule ramoso villoso erecto.
Linn. Sp. pl. 2. p. 1237. — *Roth. fl. germ.*
2. p. 325.

Aster palustris luteus, folio lanuginoso lon-
giori. *Mapp. fl. Als.* p. 33.

Conyza affinis. *Bauh. pin.* 265.

Conyza palustris repens, *Britannica* dicta.
Morif. hist. 3. p. 113. — *Sect.* 7. t. 19. f. 8.

Britannica *Conyzoides* quibusdam, *Aster*
Pannonicus. *Clus. Bauh. hist.* 2. p. 1047.
non. Oeder. fl. Dan. t. 413.

Germ. Sumpf-Alant.

Gall. Inule britanique.

Hab. In regionibus cis et transrhenanis indatis ad fossas aquaticas passim frequenter. Circa *Dachsland an der Federbach*, prope *Knielingen*, *Eckenstein* et alibi in udis gminosis dumetosis. Prope Argentinam Romanum versus passim.

Flor. Julio, Augusto. *Semina* Septembri maturantur. ¶

Radix repens. *Caulis* sesquipedalis, bi-
pedalis et altior, erectus, teres, villosus, super
ramosus. *Rami* alterni, subcorymbosi. *Folia*
alternata, sessilia, amplexicaulia, lanceola
acuminata, margine obsolete rariter et reme-
denticulata, quandoque integruscula, plani-
subtus villosulo-incana, superne læte viridia
bruscula. *Flores* caulem ramosque simpli-
terminantes, solitarii, majusculi, lutei. *Fili-
culi* radii ligulati. *Ligulae* numerosae, lineari-
graciles, longæ. *Calycinae* squamae lanceola
infimæ laxæ, longiores.

Variat. Caulibus plus minus ramosis, raro simplicibus
unifloris.

Obs. Planta in horto culta, multo procerior, si-
duplo longioribus, magis acuminatis integris-
dita est.

Nota. In Britannia non crescit: Britannicæ
men a veteribus inditum.

Flores melliferi.

Nota. *Inula odora* et *I. oculus Christi*
unquam in Germania, certe non in Suevi-
tractu nostro crescunt. *I. odora* Linn. hypo-
nostras sub dio non patitur in calidioribus
Galliæ et Hispaniæ obvia.

1300. INULA dysenterica foliis amplexicaulibus, cordato-oblongis subtomentosis, caule villosa paniculato, squamis calycinis setaceis. *Linn. Sp. pl. 2. p. 1237.*
— *Poll. fl. pal. n. 803.*

Aster foliis amplexicaulibus undulatis subtus tomentosis. *Hall. stirp. helv. n. 79.*

Aster dysentericus. *Scop. fl. Carn. ed. 2. n. 1079.*

Aster pratensis autumnalis, conyzæ folio. *Mapp. fl. Als. p. 32.*

Conyza media *Asteris* flore luteo, vel tertia *Dioscoridis.* *Bauh. pin. 265.*

Conyza I. *Tabernæm. p. 1243. Icon.*

Icon. Curtis. fl. Lond. t. 164. — Oeder. fl. Dan. t. 410. — Bulliard. herb. t. 299.

Term. Ruhr-Alant. Ruhrkraut.

Hall. Inule dysenterique.

Tab. Utrique in locis udis ad fossas in palcuis cum *Mentha Pulegio*, nec non ad stagna Rheno vicina copiose.

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. *¶*

Radix repens, cylindrica, fibrosa, albida. *caulis* pedalis, cubitalis, erectus, teres, firmus, lanuginosus, ramosus, sæpe ramosissimus. *folia* sparsa, alterna, sessilia, amplexicaulia, cordato-oblonga, acutiuscula, margine subseriata, superne subhirsuta, saturate viridia, sub-
s albide tomentosa, et rugosula. *Flores* caum ramosque terminantes, solitarii, erecti, *teji*, majusculi. *Flosculi* radii ligulati, numerosi, patentes, tridentati, semiunciam longi. *calycinæ squamæ* laxiusculæ, lineari-setacæ, *ice* subreflexiusculæ, villosæ.

iosa, undulata, hirsutula. *Flôres* caulem ramosque terminantes, erecti, parvi, numerosi, emisphaerici, lutei. *Ligulae* radii brevissimae, numerosae, tridentatae. *Calycinae* squamae lineares, inaequales, erectae, tomentosae.

Amat loca hyeme inundata in quibus copiosissima.

Arenam in sabulosis humidis figit.

Planta digitis trita, foetidiuscula, pulices fugare citur.

302. INULA *salicina* foliis sessilibus lanceolatis recurvis serrato-scabris, floribus inferioribus altioribus, ramis subangulatis glabris. *Linn. Sp. pl.* 2. p. 1238. — *Pollich. fl. pal.* n. 805.

Aster foliis glabris ciliatis ovatis, venosis. *Hall. stirp. helv.* n. 76.

Aster montanus luteus salicis glabro folio. *Bauh. pin.* 266. — *Mapp. fl. Alf. p.* 33.

Bulbonium luteum l. *Tabernæm.p.* 716. *Icon. Habitus optima.*

erm. Weidenblättriger - Alant.

all. Inule fauliere.

lab. In pratis virgultosis inter *Dachsland* et *Knielingen*. In pratis montosis apricis in Marggraviatu superiore prope *Sirnitz*, et auf der *Hexmatte* retro *Lippurg*. In dittonis Basileensis graminosis asperis montis *Muteti* et *Dornach*. In Alsatia retro *Barr* in pratis montosis, et retro *Dusenbach*; in Alsatia inferiore prope *Lampertsloch*, non inter *Colmar* et *Bollweiler t. Joh. Bauhino*. In Palatinatu inter *Dürkheim*, *Lambsheim*, *Frankenthal* et dem *Holzhof. t. Pollichio*. In Sponhemia ad *Mosellam* in pascuis montosis.

462 SYNGENESIA. POLYG. SUPERT.

Flor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ♂

Caulis pedalis, sesquipedalis, erectus, falcato-angulatus, glaber, superne ramosus. *Folia* alterna, sessilia, semiamplexicaulia, lineari-lanceolata, acuta, recurviuscula, margine tenuissime serrulata, aspera, utrinque glabra, venulosa, lacte viridia, lineas quinque ad semunciam lata. *Flores* caulem ramosque terminantes, erecti, solitarii, majusculi, lutei. *Receptaculi* Ligulae numerosae, lineares, longae. *Calycis* squamae ovato-lanceolatae, inaequales, apice recurviusculae, glabrae.

Variat. Caulibus simplicibus unifloris.

Nota. Non confundenda cum *Inula ensifolia* Linn. quae ab hacce distinguitur, Caule simplici subunifloro, Foliis sparsis, numerosioribus, lineari-lanceolatis, longe acuminatis, nervosis, integris margine asperis, longe angustioribus, 2 ad 3 lineas latis, erectis, non recurvis. Haec in Austria inferiore, vix autem in Suevia aut reliqua Germania crescit. Conf. Jacqu. fl. Austr. t. 162. ubi optime depicta est.

1303. *INULA hirta* foliis sessilibus lanceolatis recurvatis subserrato-scabris, floribus inferioribus altioribus, caule teretiusculo subpiloso. *Linn. Sp. pl.* 3. p. 1239. — *Poll. fl. pal.* n. 806.

Aster foliis ovatis venosis, scabris hirsutis. *Hall. stirp. helv.* n. 75.?

Aster luteus, hirsuto salicis folio. *Bauh. pin.*

Aster tertius pannonicus Clusii luteus folio hirsuto salicis. *Bauh. hist.* 2. p. 1047.

Aster flore luteo II. *Tabernam. p.* 717. *Icon. Icon. Jacqu. fl. Austr.* t. 358.

erm. *Rauhhariger Alant.*

all. *Inule heriffée.*

ab. In Palatinatu inter *Oppenheim*, *Worms* et *Gernsheim* in pascuis hinc inde. *t. Pollichio.*

or. Julio, Augusto. *Semina* Augusto, Septembri matura. *?*

Differt a præcedente specie, *Caule* non pro sed pubescenti, molli, *Foliis* lanceolatis, non glabris sed piloso-hirsutis, *argine* subintegris aut rariter obtuse crenulatis.

304. *INULA germanica* foliis sessilibus lanceolatis recurvis, scabris, floribus subs fasciculatis. *Linn. Sp. pl. 2. p. 1239. — Poll. fl. pal. n. 807.*

Aster foliis lanceolatis amplexicaulius oris reflexis, ramis multifloris, calycibus oblongis laxis. *Gmel. fl. Sibir. 2. p. 181. t. 78. f. 1.*

Aster thuringiacus altissimus latifolius montanus, flore luteo parvo. *Hall. jen. 181.*

Conyzae affinis germanica. *Bauh. pin. 266. Moris. hist. 3. sect. 7. t. 19. f. 26.*

on. *Jacqu. fl. Austr. t. 134.*

erm. *Teutscher Alant.*

all. *Inule d'Allemagne.*

ab. In Palatinatu inter *Odernheim* et *Oppenheim* ad vias gramineas cum *Chrysocomate Lynosyri*. In viis ac praecipitiis gramineis retro *Alzey*, ad viam versus *Niederulm* cum *Althæa hirsuta* et *Artemisia pontica*. *t. Pollichio.*

464 SYNGENESIA, POLYQ. 5072

Flor. Julio, Augusto. *Semina* Septem-
tura. ☞

Radix repens, cylindrica, dura. Ci-
dalis, sesquipedalis, cubitalis, erectus
simpliciusculus, pubescens, scabriusculi-
mitate ramosus. *Folia* sparsa, alterna,
semiamplexicaulia, oblongo-lanceolata,
cula mucrone terminata, integra aut
sime rariter denticulata, hirsutula, aspi-
nosa, margine scabra. *Flores* terminali-
bellato-fasciculati, dense congesti, parvi
Radix ligulae numerosae, 2-3 dentatae
cinnae squamae inaequales, lanceolatae,
apice reflexae, villosiusculae.

1305. *INULA montana* foliis lan-
hirsutis integerrimis, caule uniflo-
lyce brevi imbricato. *Linn. Sp.*
1241. — *Pollich. fl. pal. n.* 808.

After foliis ellipticis integerrimis tom-
caule unifloro. *Hall. stirp. helv.*

After *Atticus* luteus, montanus villosus
no flore. *Bauh. pin.* 267.

After montanus luteus mas et foemi-
bernæmont. p. 717. *Icon.*

Germ. Berg-Alant.

Gall. Inule des montaignes.

Ha b. In Palatinatu in vinearum collibus
lapidosis circa *Dürkheim* et *Calstac*
Adoni vernali, *Stipa pennata* et *Cu-*
Otite. t. Pollichio.

Flor. Majo, Junio. *Semina* Junio, Jul-
tura. ☞

Caulis pedalis, sesquipedalis, erectus,
pilosus, scaber, subangulosus, su-

aplicifoliusculus subuniflorus. Folia alterna, sessilia, subobliqua, erecto-patentia, lanceolata, tussifuscula, integerrima, pallide viridia, ubique scabra, pilosa. *Flos* terminalis luteus, speciosus pollicem et ultra in radio habet. *Calyx* communis, segmentis lanceolatis, ciliatis, pilosis, imbricatis, erecto-patentibus. *Sic Pollichius.*

Nota. Rarioribus omnino florae civibus adscribenda quam frustra hucusque in terris Badensibus et confinibus quaesivi.

CCCXCI. A R N I C A.

Linn. Gen. pl. ed. Schreb. n. 1296.

Calyx communis imbricatus, radio corollae prior: *foliolis* lanceolatis, longitudine calycis, erectis. *Corolla composita* radiata. *Corollae* hermaphroditae in disco numerosissimae. *Corollae* in radio. *Propria* hermaphroditi tubulosa, erecta, quinquefida, aequalis. *Femina* lanceolata, longissima, tridentata, patens. *Calyx* fructiferus immutatus. *Semina* oblonga. *Stylus* capillaris, pubescens, longus. *Receptaculum* nudum aut tomentosulum.

606. *ARNICA montana* foliis ovatis integris: caulinis geminis oppositis. *Linn. Sp. pl. 2. p. 1245. — Poll. fl. pal. n. 809.*

Arnica foliis conjugatis ovatis integerrimis. *Hall. stirp. helv. n. 90.*

Doronicum plantaginis folio alterum. *Bauh. pin. 185. — Mapp. fl. Alf. p. 94.*

Chalta alpina. *Tabern. p. 714. Icon mala.*

lera Bad. Tom. II.

G g

Icon. Regnault. bot. — Oeder. fl. Dan. t. 1.
— Blackw. t. 595. — Schumbr. bot. han
t. 248.

Germ. Bergwolverley. Fallkraut.

Gall. Arnique montanière. Plantain des m
tagnes.

Hab. In pratis pascuisque montosis præsert
altioribus, auf dem Kaltenbrunn, auf
Herrnwiese; in convallibus Bühlerthal p
sim frequens. In Marggraviatu superio
in montis Kandel, Belchen et Feldba
pascuis nec non prope Gersbach. In Be
goviae montibus prope Fürtwangen, G
tebach et alibi in pascuis montosis pass
abunde. In Alsatiae Vogels in summis
cuis montis Ballon frequens; in mo
Bloutberg retro Marienkirch; in mo
Hochfeld, prope Schönenburg copie
prope Oberhütt im Urbisthal, nec
in monte Passberg prope Buchweiler.
Palatinatu circa Lautern retro der Es
ferth, Hochspeyer in sylvarum pascuis ge
mineis passim; prope Espelsteg; in pasc
turfosis inter Hirschhausen, Speßbach
Ramstein; ad patibulum prope Landst
in montibus sylvaticis prope Steinba
t. Pollichio.

Flor. Majo, Junio. *Semina* Julio, Augu
matura. 7

Radix fibrosa, fusco-lutescens, intus albid
Caulis pedalis, sesquipedalis, erectus, simple
teretiufculus, striatus, subpubescens, super
subtriflorus. *Folia* radicalia ovato-lanceolat
obtusiuscula, integra, trinervia, laete viridi
subtus pallidiora, crassiuscula, subvillosa,
petiolum decurrentia; *caulina* duo, opposit

SYNGENESIA. POLYG. SUPERF. 467

ternave, sessilia, lanceolata, subminora, *summa* ad pedunculorum basin connata, multo inora, acuta. *Flores* caulem terminantes, saepe tres, subnutantes speciosi, aurei, pedunculati. *Pedunculi* erecti, hirsuti, *Bractea* una terave alterna, foliacea, erecta, sessili, lineari, integra, hirsuta praediti. *Flosculi* radii longe pulati, lineares, obtusi, striati, lutei, apice dentati, 14-15 lineas longi, binas lineas. *Calycinae* squamae lanceolatae, erectae; *involuculae*, apice fusco rubellae. *Receptaculum* hac specie non nudum, sed tomentosum.

Pharm. ARNICÆ Radix; Herba; Flores.

Planta acris, amara, odorata, emetica. errhina, diaphoretica, efficacissima saepe in Paralyfi.

Putant quidam sanguinem per eam resolvi grumoso, unde Panaceam lapforum vocant. Ab antiquis temporibus, jam in casu ab alto, ad singulas et contusiones, maxime commendata.

In contusionibus quas bovillum genus sibi mutuo ext, praedicatur. Schreb. Samml. 1. p. 226.

Rustici Smolandiae folia in pulvere nasali assument, unde sternutant, uti ab Helleboro, eadem fumo hauriunt, ceu tabacum. Linn. fl. Suec. p. 295.

Nota. Flores Arnicae in officinis pharmaceuticis saepe alterati cum alienis floribus luteis flosculosis, e. gr. m Calendula officinali, Doronico Pardaliche, Authemide tinctoria, Inula dysenterica et alia specie.

CCCXCII. DORONICUM.

Linn. Gen. plant. ed. Schreb. n. 1297.

Calyx communis foliolis lanceolato-subulatis, aequalibus, erectis, duplicis seriei. *Colla composita.* *Corollulae* hermaphroditae

Gg 2

SYNGENESIA. POLYG. SUPE

lae, numerosae in disco. *Feminae*
i radio. *Propria* hermaphroditis
mis: *Limbo* quinquefido, patulo.
ulata, lanceolata, tridentata. *Ca*
leviter connivens. *Semina* obovat
fulcata. *Pappus* hermaphroditis
emineis nullus. *Receptaculum*

1.

Bellidiastro pappus etiam in

1307. *DORONICUM pardalia*
foliis cordatis obtusis denticulatis
calibus petiolatis; caulinis amp
libus. *Linn. Sp. pl.* 2. p. 1247.

Doronicum foliis cordatis obtusis, im
petiolatis, superioribus amplexic
Hall. stirp. helv. n. 88.

Doronicum maximum foliis caulem
xantibus. *Bauh. pin.* 185. — *Cam*
823.

Doronicum. I. *Tabernaemont. p.* 71

Icon. Jacqu. fl. Austr. t. 350. — *Schku*
Handb. t. 249.

Germ. Gemeine Gemswurz.

Gall. Doronic mont-aux panthères.

Hab. In sylvaticis montosis. Ní fallor
Principatu Fürstenbergensi. In parte
rosa horti Anglici aulici *Carlsruhani*
quasi spontanea.

Flo r. Junio, Julio, Augusto. *Semina* A
Septembri matura. 7

Radix cylindrica, articulata. repens, flora, barbata, alba. Caulis 3-5 pedalis, us, striatus, simplex. Folia radicalia caulinferiora petiolata, basi auriculata, cordata, lita, villosa, mollia, venosa; superiora a, amplexicaulia, acutiora. Flores terminatus alterve longe pedunculatus, erectus, s majusculus. Calycinae squamae ciliatae, æ, acutae, corolla duplo breviores. Radia li ligulati, lineati, tridentati, numerosi. na hermaphroditis decemfiriata, hirsuta, , pappo piloso coronata. Feminis femina s hirsuta, pappo destituta. Receptaculum cum subvillosum.

r m. DORONICI Radix.

Radix aromatica. recens subdulcescens, non vena-
Hodie in medicina exoleta.

Flores melliferi.

8. *DORONICUM plantagineum*

foliis ovatis acutis subdentatis, ramis alternis. Linn. Sp. pl. 2. p. 1247. — Lamarck et Decand. fl. franc. 4. p. 174.

Doronicum plantaginis folio. Bauh. pin. 184.

Doronicum minus officinarum. Dalechamp. hist. 1202. f. 2.

m. Wegwartblättrige Gemswurz,

l. Doronic a feuilles de plantain.

b. In Vogefis prope Bourmont et Remiremont. t. Buchoz.

ta. Plantam hucusque frustra in terris Badensibus et confiniis quaesivi.

9. *DORONICUM Bellidiastrum*

scapo nudo simplicissimo unifloro. Linn. Sp. pl. 2. p. 1247.

SYNGENESIA. POLYG. SUPERF.

Arnica Bellidiastrum. Villars. fl. Delph. p. 212. — Lamarck et Decand. fl. fr. 4. p. 176. Willden. Linn. Sp. pl. 3. p. 210.

Arnica caule nudo unifloro, foliis ovato lanceolatis, serratis. Hall. stirp. helv. n. 9.

After Bellidiastrum. Scopol. fl. Carn. n. 104.

Bellidiastrum alpinum foliis brevioribus hirsutis, caule palmari, flore albo. Michx. gen. 32. t. 29.

Bellis media. Tabernæmont. p. 708. Icon. Clus. hist. 2. p. 44. — Camer. Epit. 65.

Icon. Jacqu. fl. Austr. t. 400.

Germ. Masliebenartige Gemswurz.

Gall. Doronique paquerette.

Hab. In ditione Basileensi in monte Farnspurg Wallenburg, Ramstein, Wasserfalla Langenbruck, Digten, Schaffmatt et alibi non infrequens. In Principatu superiori prope Constanz, Gottlieben, Stein et alibi in umbrosis graminosis abunde nuper videtur. In Principatu Fürstenbergensi prope Donaueschingen in der Baar, prope Fürstenberg et alibi frequens.

Flor. Aprili, Majo, Junio, Semina Junio, Julio matura. 7

Radix obliqua, cylindrica, longe fibrosa, fusco nigricans. Scapus palmaris, spithameus, sesquipedalis, erectus, simplicissimus, teres, hirsutus, aphyllus, uniflorus. Folia radicalia plura, in petiolum decurrentia, obovata vel lanceolata, dentato-serrata, obtusa, utrinque villosa, laete viridia, asperiuscula. Flos Bellidis majusculus. Calycinae squamae lineari-lanceolatae, erectae, subæquales, villosulae, virides. Flosculi foeminei in radio ligulati, 2.

SYNGENESIA. POLYG. SUPERF. 471

mutati, quandoque integri, patentés, calyce
iplo longiores, nivei vel rubelli. *Discus* luteus.
Stamina radii et disci pappo subpiloso coronati.
Receptaculum nudum, convexum.

Nota. Bellis sylvestris media, caule carens quam
Mapp. fl. Alf. p. 37. in pratis et sylvis retro
Barr citat, non huc, sed ad variationem majore
rem Bellidis perennis pertinet. Regionem
prope Barr saepe lustravi, neque vero in his,
aut ad jacentibus regionibus unquam, Doronicum
Bellidiastrum vidi.

b. f. Planta in horto botanico culta, mutationi non
succubuit.

CCCXIII. BELLIS.

Linn. Gen. pl. ed. Schreb. n. 1300.

Calyx communis hemisphaericus, erectus:
lobis decem ad viginti, duplici serie positis,
oblongatis, æqualibus. *Corolla* composita radiata:
Corollulae hermaphroditæ tubulosæ, numero
se in disco. *Femineae* ligulatae plures in
radio. *Propria* hermaphroditi infundibuliformis,
quinquefida. *Feminea* ligulata, lanceolata, vix
identata. *Calyx* fructiferus immutatus. *Semina*
obovata, compressa. *Pappus* nullus. *Receptaculum*
nudum, conicum.

310. BELLIS perennis scapo nudo. *Linn.*

Sp. pl. 2. p. 1248. — Pollich. fl. pal. n. 810. — Hall. stirp. helv. n. 91.

Bellis sylvestris minor. *Bauh. pin. 261. — Mapp. fl. Alf. p. 37.*

Bellis I — IV. *Tabern. p. 707. 708. Icones.*

Pharm. BELLIDIS MINORIS Herba ; Flo

Multae in hortis floribus rubris, carneis, impletis, variationes.

Flores apibus melliferi.

CCCXCIV. CHRYSANTHEMU

Linn. Gen. pl. ed. Schreb. n. 1307

Calyx communis hemisphaericus, intus : squamis acute incumbentibus : interius per gradus majoribus, margine scariosis i terminatis squama scariosa. Corolla com radiata : Corollulae hermaphroditae tubi numerosae in disco. Femineae duodeni in radio. Propria hermaphroditis infundibuliformis, quinquefida, patula, longitudine aequa. Femineis ligulata, oblonga, tridentata. Calyx fructiferus immutatus. Semina oblonga, perispermata destituta. Receptaculum nudum, punctatum, convexum.

Linn. Sp. pl. 2. p. 1251. — Pollich. fl. pal. n. 812.

Matricaria foliis radicalibus petiolatis, ovatis, crenatis, caulinis amplexicaulibus dentatis. Hall. stirp. helv. n. 98.

Leucanthemum vulgare. Mapp. fl. Als. p. 168.

Bellis sylvestris caule folioso major. Bauh. pin. 261.

Bellium majus. Tabern. p. 710. Icon.

n. Bulliard. herb. t. 211. — Oeder. fl. Dan. t. 994. — Blackw. herb. t. 42.

m. Grofse Maslieben. Grofse Gänseblumen.

1. Leucantheme vulgaire. Grand Marguerite.

Utrunque in pratis, et cultis.

Junio, Julio. Semina Augusto matura. 7

Caulis pedalis, sesquipedalis et altior, erectus, simpliciusculus, quandoque ramosus, iusculus, interdum villosus. Folia radicalinaque inferiora petiolata, oblonga, plusve profunde dentata, superiora sessilia, amplexicaulia, rariter ferrata, glabra aut a. Flos terminalis magnus. Radii ligulæ dæ, patentes. Discus convexus, luteus.

at præsertim in montosis calcareis, Caulibus longioribus, crassioribus, Foliis obtuse crenatis. Floribus majoribus.

m. BELLIDIS MAJORIS Herba; Flores.

lores melliferi.

2. CHRYSANTHEMUM *inodorum*

foliis pinnatis multifidis, caule ramoso diffuso. Linn. Sp. pl. 2. p. 1253. — Poll. fl. pal. n. 813.

obsoletis. *Linn. fl. Succ. p.*
Schreb. Spic. p. 11.

Icon. Oeder. fl. Dan. t. 696. — Schk
handb. t. 253.

Germ. Geruchlose Wucherblume.

Gall. Leucantheme inodorante.

Hab. Circa Carlsruhe et in alibi in
ad vias, in agris hortis et ad
marginalia frequens.

Flor. Junio, Julio, Augusto. Semi
tembri matura. ☉

Radix fibrosa. Caulis pedalis, ses
lis, erectiusculus, sæpissime diffusus
ramosus. Rami alterni diffusi. Folia
sessilia, pinnata, multifida, glabra,
dia. Flores caulem ramosque termina
litarii, mediocres. Radii ligulae albæ
ciam longæ, striatæ, patentes, albæ
tatae. Discus luteus. Calycis hemisphæ
mae lanceolatae, obtusæ, margine
naceae, fuscae. Seminum corona integ
ra. Legumina. Pericarpium. Semina

: conico, Radii ligulis triplo majoribus patentibus,
: deflexis, odore fere nullo.

Flores sæpe cum floribus Chamomillae confun-
tur.

13. *CHRYSANthemum corymbo-
sum* foliis pinnatis inciso-ferratis, caule
multifloro. *Linn. Sp. pl. 2. p. 1251. —*
Poll. fl. pal. n. 811.

Pyrethrum corymbosum. Scop. fl. Carn. n.
1043. Willden. Linn. Sp. pl. 3. p. 2155.

Pyrethrum foliis pinnatis, pinnis semipinna-
tis acute dentatis, floribus umbellatis.
Hall. stirp. helv. n. 95.

Matricaria Tanacetifolio flore minore, semine
umbilicato. *Mapp. fl. Alf. p. 189.*

Tanacetum inodorum flore minore. *Bauh.*
pin. 132.

Tanacetum Leucanthemum. Tabernæmont.
p. 379. Icon.

. *Matricaria Tanacetifolio*, flore majore, se-
mine umbilicato. *Mapp. fl. Alf. p. 189.*

Tanacetum inodorum flore majore. *Bauh.*
pin. 132.

o n. *Jacqu. fl. Austr. t. 379.*

: r m. *Doltentraubige Wucherblume.*

ll. *Leucantheme en corymbe.*

ib. In Brisgovia am Kaiserstuhl in rupibus
et ad arcem antiquam *Limburg*, circa
Sponck, *Burgheim* et *Altbreisach*. In Marg-
raviatu superiore in monte *Schloßberg*
prope *Badenweiler*; in montosis lapidosis
asperis retro *Oberweiler* et supra *Vöges-*
heim; in monte prope *Crenzach*, *Farns-*

spurg, Wallenburg, Ramslein, Landbruck, Digten, Schaffmatt et m. Wallen etc. cum β. In Alsatia ad arces tiquas passim. In Palatinatu in sylvis r tolis circa Steinbach abunde; in m Platten et Reifberg t. Pollichio.

Flor. Julio, Augusto. Semina Septembri tura. 7.

Radix subcylindrica, obliqua, subram fibrosa, fusco-nigricans. Caulis cubitalis, pedalis, erectus, sulcato-angulatus, fir glabriusculus, inferne simplex, superne ram corymbosus. Folia alterna, sessilia, pinn foliolis oblongis, acutis, inciso-serratis, l mis confluentibus, subtus villosiuscula. Fl corymbosi, terminales, numerosi, umbel planam imitantes, majusculi. Flosculi rad gulati, albi; disci lutei. Calycinae squamae gine fusco-nigricantes, villosulae. Semina pa destituta, membrana denticulata, coronata

Planta inodora, degustata insipida, demum uscula.

Flores melliferi.

†† *Chrysanthema.*

1314. CHRYSANTHEMUM *segeti*
foliis amplexicaulibus superne lacin
inferne dentato-serratis. *Linn. Sp. p.*
p. 1254. — Pollich. fl. pal. n. 814.

Chrysanthemum folio minus secto, gla
Bauh. hist. 3. p. 105. — Mapp. fl.
p. 71.

Bellis lutea foliis profunde incisus m
Bauh. pin. 262.

Chrysanthemum arvense. Tabernaemont. p. 366. Icon.

Icon. Curtis. fl. Lond. — Bulliard. herb. t. 339. — Oeder. fl. Dan. t. 995.

Berm. Gelbe Wuttherblume.

Pall. Chrysanthème des blés.

Hab. Inter segetes passim. Retro Grözingen cum Adonide aestivali rarius. Prope Argentinam inter segetes prope Ofswald et Illkirch cum Caulalide grandiflora. In Palatinatu circa Lautern in agris; circa Niedermisau et Jettenbach etc. t. Pollichio. In Sponhemia inter frumenta passim non infrequens.

Flor. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Radix subfusiformis tenuis, perpendicularis, subfibrosa albida. *Caulis* pedalis, sesquipedalis et altior, erectus, teretiusculus, glaber, iridi-glaucus, ramosus. *Rami* alterni, brevifculi. *Folia* alterna, sessilia, semiamplexicauda, erecto-patula, oblonga, aut oblongo-subvata, basi sagittata, sinuato dentata, mollia, trinque glauca. *Flores* caulem ramosque terminantes, erecti, solitarii, flavi, majusculi. *Carcinae* squamae ovatae, interiores majores, imae margine membranaceae, lacerae, subfusae, scariosae. *Corollulae* radii 16-18 ligulae, blongae, truncatae, emarginatae, subtridentatae, neis duabus exaratae, luteae. *Discus* luteus. *Semina* oblonga, utrinque truncata, sulcata, obangulata, subincurva, pallide fusca, pappo aestitata. *Receptaculum* nudum, punctatum, convexum.

Flores apibus melliferi.

co hemisphaerico. *Femineae* in radio
Propria hermaphroditi infundibuliformi
quefida, patens. *Feminea* oblonga, tri
Calyx fructiferus immutatus. *Semina* c
Pappus nullus. *Receptaculum* nudum
vexum.

1315. *MATRICARIA Parthenium*
compositis planis: foliolis ovatis
pedunculis ramosis. *Linn. Sp.*
1255.

Pyrethrum foliis petiolatis compositi
foliolis ovatis incisis, pedunculis
corymbosis, caule erecto. *Smith.*
2. p. 900. — *Willden. Linn. Sp.*
p. 2155.

Matricaria foliis pinnatis, pinnis serr
tis, laciniis obtusis, floribus um
Hall. stirp. helv. n. 100.

Matricaria vulgaris f. *fativa*. *Bauh. p.*
— *Mapp. J. l. Alf. p.* 189.

SYNGENESIA. POLYG. SUPERF. 479

n. Bulliard. herb. t. 203. — Regnault. bot.
— Oeder. fl. Dan. t. 674. — Blackw. t. 192.
r m. Mutterkraut. Metram.

ll. Matricaire odorante. Matricaire.

b. Utrunque ad pagorum et urbium muros,
sepes et rudrata, nec non in hortis. β .
passim in hortis colitur.

o r. Julio, Augusto, Septembri. *Semina* Sep-
tembri, Octobri matura. γ

Radix cylindrica fibrosa. *Caulis* cubitalis
altior, erectus, angulatus, firmitus, glaber,
noscus. *Rami* alterni, erecti, subpubescentes,
divisi, corymbosi. *Folia* alterna, petiolata,
dentata, glabriuscula, punctis minutis adper-
sa, pinnatifida; *pinnis* suboppositis ovato-ob-
longis, inciso-serratis. *Flores* terminales, co-
mpositi, erecti, pedunculati, inferne foliosi,
diocres. *Bractea* lineares, angustae ad basin
dunculorum, et in ipsis pedunculis. *Calyci-*
e squamae lineares, convexo-carinatae, ci-
reae, margine membranaceae, subciliatae,
interiore scariosae, adpressae. *Flosculi* radii ligulati,
5; disci lutei. *Semina* pappo destituta. *Re-*
ceptaculum nudum, convexum.

aria t. Flosculis omnibus tubulosis, radio nullo; et
flosculis disci ligulatis.

arm. MATRICARIAE Herba; Flores.

Odor plantae peculiaris, graviusculus. Sapor ama-
nus, aromaticus. Resolventes, tonicas, emmenago-
gogas vires habet. Nimis omnino hodie in medicina ne-
cessaria. Ab Anglis succus plantae in febribus intermit-
tentibus adhibetur, hinc illis nomen *Feverfew*, etiam
infuso aquoso exhibent.

316. MATRICARIA *Chamomilla* re-
ceptaculis conicis, radiis patentibus, (de-

mum reflexis) feminibus nudis, f
calycinis margine aequalibus. *Li*
pl. 2. *p.* 1256. — *Poll. fl. pal.* n. 8

Matricaria foliis planis capillaribus,
cato pinnatis, pinnis lanceolatis
trifidisque. *Hall stirp. helv.* n. 101.

Chamaemelum vulgare, *Leucanthem*
oscoridis. *Bauh. pin.* 135. — *Ma*
Alf. *p.* 68.

Chamaemelum I. *Tabernaemont.* *p.* 1
Icon. Curtis. fl. Lond. t. 331. — *Blachw.*
Germ. Chamillen. Aechte Chamillen.
Gall. Chamomille.

Hab. Utrunque in agris, cultis, vineis,
ratis et alibi passim abunde.

Flor. Junio, Julio, Augusto, Septemb
mina Septembri, Octobri matura.

Flosculi radii ligulati albi, patentes,
deflexi. *Discus* conicus luteus. *Receptac*
conicum, nudum, qua nota facile ab *Anth*
distinguitur.

***Pharm.* CHAMOMILLÆ VULGARIS Fl**
Herba.

Nota. Plantæ quamvis tritissimæ, vulgatissim
officinis pharmaceuticis sæpe ob botanices
rantiam *Anthemis arvensis* et *Cotula*,
Chrysanthemum inodorum substituuntur.

Obs. Forſan Linnaeus sub *Matricaria* ſua veol
radiis reflexis, noſtram Chamomillam in mac
arenosis apricis obvenientem, ſemper minorer
diis magis reflexis, floribus magis odoratis
veolentibus comprehendebat.

Vidi in Caſtilia, præſertim Madriti plantulam n
ramoſam diffuſam fragrantiffimam, quam maxime ſu
le

ntem, floribus triplo minoribus, praeditam quam Bonici madritenses pro vera *Matricaria suaveolente* bent, ex qua infusum Theiforme omnium gratissimum et amicissimum Ludovicum Née, (botanicum aculeum, cujus inventa ex Cavanillidis Iconibus sum nota sunt) degustavi. Haec plantula, sane efficacissima, in Madriti officinis pharmaceutricis, sub *CHAMOMILLA SUAVEOLENTE* nota. Plantula annua ex nine in horto Carlsruhano culta, mutationi non succubuit, ibi etiam fragrantissima; pro dolor periit, nec nina peperit. Haec omnino vix in Suecia et Germania ullibi conspicua! Flosculis radii ligulatis brevissimis, saepe deficientibus instructa. Receptaculum conicum, nudum. Odor omnino Chamomillae sed fragrantior, magis suaveolens.

CCCXCVI. ANTHEMIS.

Linn. Gen. plant. ed. Schreb. n. 1312.

Calyx communis hemisphaericus: squamis paribus, subaequalibus. *Corolla* composita radiata: *Corollulae* hermaphroditae tubulosae numerosae in disco convexo. *Femineae* plures quam inque in radio. *Propria* hermaphroditi infunduliformis, quinquedentata, erecta. *Femineae* ulata, lanceolata, interdum tridentata. *Calyx* setiferus immutatus. *Semina* oblonga. *Pappus* nullus seu marginatus. *Receptaculum* paraceum, convexum, s. conicum.

† *Radio discolore: albo.*

§17. *ANTHEMIS arvensis* receptaculis conicis, paleis setaceis, seminibus coronato-marginatis. *Linn. Sp. pl. 2. p. 1261.* *Pollich. fl. pal. n. 816.*

Chamæmelum foliis glabris duplicato-
tis, nervo foliaceo, pinnulis lanceo-
feminibus lævibus. *Hall. stirp. helv.*

Chamæmelum inodorum. Bauh. pin.
Mapp. fl. Alf. p. 68.

Chamæmelum inodorum sive *Cotula*
foetida. *Bauh. hist. 3. p. 120.*

Cotula. I. *Tabernæmont. p. 70. Icon.*
Is oñ. Oeder. fl. Dan. t. 1178.

Germ. Unächte Ackerchamomille.

Gall. Anthemide des Champs.

Hab. Utrunque in agris, arvis, ruderat-
neis passim frequens.

Flor. Junio Julio, Augustio. *Semina* Au-
gustio Septembri matura. ☉

Radix fibrosa. *Caulis* pedalis, sesqu-
lia, adscendens, sæpe diffusus, a basi ran-
teretiufculus, striatus, villosiusculus. *Ra-*
terni. Folia alterna, sessilia, bipinnata,
fida, latiuscula, utrinque villosula, gl-
viridia. *Flores* caulem ramosque termin-
solitarii, mediocres, *Radio* ligulati albi,
patentes; *Disco* lutei. *Calycinæ* squam-
pressæ, obtusæ, pubescentes, margine
branceæ, albidæ. *Receptaculum* paleis la-
latis, acuminatis, splendentibus, obsessum
mina pappo destituta, coronato-marginata

Odore multo debiliore prædita, quam
species.

Flores melliferi. ♀

1318. *ANTHEMIS Cotula* recept-
conicis: paleis fetaceis, feminibus
Linn. Sp. pl. 2. p. 1261. Pollia
pal. n. 817.

SYNGENESIA, POLYG. SUPERF. 483

Chamæmelum foliis glabris duplicato-pin-
natis, nervo foliaceo, pinnulis lanceo-
latis, seminibus lævibus. *Hall. stirp. helv.*
n. 104.

Chamæmelum foetidum. *Bauh. pin.* 135. —
Mapp. fl. Als. p. 69.

Chamæmelum foetidum f. *Cotula foetida.*
Bauh. hist. 3. p. 120.

on. Curtis. fl. Lond. t. 329. — *Oeder. fl.*
Dan. t. 1179. — *Blackw. t.* 63. — *Schkuhr.*
bot. hand. t. 254.

erm. Hundchamomille. Stinkende Chamille.

all. Anthemide puante. Camomille puante.

ab. Utrunque in agris, cultis, ruderatis pas-
sim frequens.

for Julio, Augusto. Semina Septembri ma-
tura. ☉

Distinguitur a præcedente specie, *Caule* ma-
jore erecto, minus ramoso, *Calycum* squamis
minus marginatis, *Paleis* in receptaculo, seta-
is minoribus, angustioribus, *Odore* peculiari,
etido.

harm. COTULÆ FOETIDÆ Herba; Flores.

Nota. *Anthemis nobilis.* Linn. et A. Pyre-
thrum, nec in terris Badensibus nec facile us-
quam in Germania sponte crescunt.

†† *Flore concolore f. luteo.*

319. ANTHEMIS tinctoria foliis bipin-
natis serratis, subtus tomentosis, caule
corymbofo. *Linn. Sp. pl.* 2. p. 1263. —
Poll. fl. pal. n. 818.

Hh 2

Chamæmelum foliis pinnatis hirsutis simplicibus lanceolatis. *Hal helv. n. 105.*

Buhtalmum Tanaceti minoris foliis. *pin. 134. — Mapp. fl. Als. p. 45*
Icon. Oeder. fl. Dan. t. 741. — Blackw
Germ. Färberchamille. Gillblume.
Gall. Camomille teinturiere.

Hab. In collibus lapidosis asperis et antiquas passim. Retro *Durlach* in rarius. In *Marggraviatu* superiore ac prope *Badenweiler*, et auf dem *Cren Horn*. In *Brisgovia* prope *Altbreisach*, *Rothweil*, *Limburg* et alibi a *Jersuhl*. In *Alsatia* prope *Othmar*, *Rhenum* ad arces antiquas prope *Rweyher*, *Bebelen*, et alibi in vineis *Kaisersberg*; in arvis siccis in *Ried Markelsheim*. In *Palatinatu* prope *heim* in montibus aridis lapidosis, in arcis circa *Oppenheim* et alibi. *Cirtern* in arvis et collibus sterilibus *lichio*. In *Sponhemiae* collibus passim infrequens.

Flor. Junio, Julio, Augusto. *Semina* A Septembri matura. 7

Radix ramoso-fibrosa, albida. *Ca* quipedalis, bipedalis, erectus, teres, usculus, subincanus, ramosus. *Rami* divisi. *Folia* alterna, bipinnata, denticulata. *suprema* pinnata, summa sæpe integra omnia pubescente-scabriuscula, ex viridibus cana. *Flores* caulem ramosque terminantes solitarii, erecti, majusculi, lutei. *Calycis* squamae arcte incumbentes, interioribus lanceolatis, obtusis, marginibus subciliatis, villosiusculis. *Radii* f

SYNGNESIA, POLYG. SUPERF. 485

lati, patentes, obtusi, striis binis longitudinalibus exarati, tridentati. *Discus* luteus. *Seta* subcurvata, striata, subrotunda, utrinque appendiculata.

Flores luteo colore lanam tingunt.

CCCXCVII. ACHILLEA.

Linn. Gen. pl. ed. Schreb. n. 1313.

Calyx communis ovatus, imbricatus: *Squar-*
ovatis; acutis, conniventibus. *Corolla com-*
ta radiata. *Corollulae* hermaphroditae, tu-
sae in disco. *Femineae* ligulatae quinque
ecem in radio. *Propria* hermaphroditi infun-
diformis, quinquefida, patula. *Feminea* li-
ta, obcordata, patens, trifida: lacinula
media minore. *Calyx* fructiferus vix muta-

Semina ovata, flocco instructa. *Pappus*
us. *Receptaculum* paleaceum, elevatum:
is lanceolatis, longitudine flosculorum.

20. *ACHILLEA Ptarmica* foliis lan-
ceolatis acuminatis argute serratis. *Linn.*
Sp. pl. 2. p. 1266. — *Pollich. fl. pal.*
n. 819.

Achillea foliis linearibus lanceolatis acutis-
sime serratis. *Hall. stirp. helv. n. 117.*

Ptarmica vulgaris, folio longo serrato, flore
albo. *Bauh. hist. 3. p. 147.* — *Mapp. fl.*
Alf. p. 253.

Anacetum album acutum. *Tabernaemont.*
p. 1164. Icon.

ionas et paludes paucim frequens.

Flor. Julio, Augusto, Septembri,
Septembri, Octobri matura, 7

Radix horizontalis, repens, subger-
alba, fibrillosa. *Caulis* pedalis, sesqui-
et altior, erectus, inferne teres, gla-
perne subangulatus, ramosus, subvillo-
lia sparsa, alterna, sessilia, amplexica-
nearia, acuta, margine argute serrulata,
subtus trinervia, saturate viridia. *Flores*
ramosque terminantes, corymbosi, nivi-
cteae lineares in pedunculis. *Calycinae* o-
vato-lanceolatae, erectae, subcarinatae,
subciliatae, subcariosae, villosae. *Flosculi*
ligulati albi. *Ligulae* patentes, bifidae,
apice obtusae tridentatae. *Flosculi* discini
albi. *Semina* nuda utrinque subulata,
apice truncata. *Receptaculum* paleaceo-
leis membranaceis, lineari-lanceolatis
sis, vix longitudine florum.

Variatio β . floribus impletis pulcherrimis passu-
tis obvia.

SYNGENESIA. POLYG. SUPERF. 487

Pecoribus gratum pabulum. Schreb. Samml. 3.
58.

Flores apibus mel praebeant.

121. ACHILLEA *Millefolium* foliis bipinnatis nudis: laciniis linearibus dentatis, caulibus fulcatis. Linn. Sp. pl. 2. p. 1267. — Poll. fl. pal. n. 820.

Achillea foliis pinnatis, pinnis longe æqualibus pinnatis, pinnulis trifidis et quinquefidis. Hall. stirp. helv. n. 107.

Millefolium vulgare album. Bauh. pin. 140.
— Mapp. fl. Alf. p. 198.

Millefolium terrestre II. Tabernæm. p. 373.
Icon.

Millefolium purpureum majus. Bauh. pin. 140. — Mapp. fl. Alf. p. 198.

Millefolium purpureum III. Tabernæmont. p. 374. Icon.

n. Curtis. fl. Lond. — Oeder. fl. Dan. t. 737. — Blackw. t. 18.

r m. Schafgarben. Garbenkraut.

11. Millefeuille.

b. Utrique in aggeribus, pratis et pascuis, ad viarum marginalia passim frequens.

or. Junio, Julio, Augusto, Septembri. Semina Septembri, Octobri matura. 7

Radix repens, filiformi-cylindrica, fibrilis, cinereo-albida. Caulis pedalis sesquipedalis et altior, erectus, subangulatus, inferne priusculus, superne fulcatus, sublanuginosus, ramosus. Folia alterna, radicalia petiolata, linearia sessilia, subamplexicaulia, bipinnata, rarius glabra, subtus villosula; pinnis oppositis,

circiter duodecim. *Receptaculum* palea
Pharm. MILLEFOLII Herba; Summit

Planta odorata, amaricans, subaromatica
cula, tonicas, resolventes et vulnerarias vires
Flores melliferi.

1322. *ACHILLEA nobilis* foliis bip
inferioribus nudis planis, super
obtusis tomentosis, corymbis c
confertissimis. *Linn. Sp. pl. 2. p. 1*
Poll. fl. pal. n. 821.

Achillea foliis tomentosis duplicato p
pinnis remotis, pinnulis semip
Hall. stirp. helv. n. 109.

Millefolium nobile Tragi. *Mapp. fl.*
197.

Tanacetum minus album odore can
Bauh. pin. 132.

Icon. Schkuhr. bot. handb. t. 255.

Germ. Edle Schafgarben.

Call. Achillea nobilis

SYNGENESIA. POLYG. SUPERF. 409

excavatis praeruptis, ad vinearum marginalia *am Kaiserstuhl* abunde. In Marggr. superiori in collibus aridis circa *Blansingen*, *Efringen* et alibi. In Alsatia in arvis inter *Osten* et *Siegelheim* et passim alibi. In Palatinatu circa *Dürkheim* in collibus et viis siccis abunde. In Sponhemiae collibus aridis.

Fr. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ζ

Distinguitur a praecedente specie, *Caule* niffiore, frictiore, duriore, magis angulato, *ato*, incano, pubescente, *Foliis* durioribus, *rioribus*, incanis, *Corymbo* florum densiffimo congesto, *Calycum* squamulis minoribus, *gine* lutescentibus, *Corollulis* radii non nigrae aut carneis, sed semper sordide albido-lide sulphureis, minoribus; *disci* ex albido-eis.

a r m. MILLEFOLII NOBILIS *Herba*; *Summitates*.

Ob odorem et saporem fragrantiores! magis in phoram incidentem, in remediis omni jure praecedi speciei anteponi meretur. Planta efficax, nimis in icina neglecta.

CCCXCVIII. G A L I N S O G A.

Ruiz et Pav. fl. Peruv. et Chil. prodr. 110.

Calyx communis hemisphaericus imbricatus squamulis erectis concavis. *Corolla* composita radiata. *Radii* subquinque feminei, fertuorum corolla ligulata, obovata, truncatrifida. *Disci* flosculi plures, hermaphroditi, indibuliformes, *limbo* quinquedentato, ere-

SYNGENESKA: POLEGE: SUNNER:

etc. Calyx fructiferus immutatus. Semina tu
 1323. Coronata. Pappo sessili, brevi, pale
 sed, paleis ciliatis. Receptaculum paleaceum
 paleis distinctibus apice mucronatis, sapis
 Radix fibrosa. Caulis sesquipedalis, bip

1323. *GALINSOGA parviflora* ca
 herbaceo, foliis oppositis ovato-lance
 latis subserratis, pedunculis elongat
 uni-bifloris. Cavan. Icon. 3. p. 41.

Galinsoga parviflora foliis ovatis triplin
 viis serratis. Willden. Linn. Sp. pl. 2258.

Wiborgia Acnella. Roth. Catalect. 1. p. 112.

Bidens mercurialis folio, flore radiato.
 vill. peruv. 1. p. 46. t. 32.

Icon. Cavanill. l. c. t. 281.

Germ. Kleinblumige Galinsoga.

Gall. Galinsoga à petites fleurs.

Hab. In Peru. Ex horto botanico Carlsruh
 in agrum prope dem Holzmagazin mig
 tione facta, nunc per tres annos quasi s
 tanea.

Flor. Junio, Julio. Semina Augusto matura.

Radix fibrosa. Caulis sesquipedalis, bip
 dalis, erectus, glaber, ramosus. Rami opp
 siti, erecti, divisi. Folia opposita, breviter
 tiolata, subconnata, ovato-lanceolata, sub
 rata, trinervia, glabra, late viridia. Flor
 axillares, terminalesque, parvi. Pedunculi, ere
 cti, elongati uni aut saepe biflori. Calycina
 squamae ovatae duplici serie, glabrae, quinqu

SYNGENESIA, POLYG. SUPERF. 497

exteriores carinatae, apice scariosae, interiores laniusculae coloratae. *Flosculi Radii* quinque gulati, parvi, albi; *Disci* lutei.

Hujus plantae semina accepi Madriti a Dno PALAU VERDERA professore Botanices dignissimo, amantissimo, cui seni generoso multa, gratissima mente deo. Planta in horto botanico sese quovis anno sponte effeminat, forsan brevi tempore in cultis Europae ingena.

Flores apibus melliferi.

CCCXCIX, BUPHTHALMUM.

Linn Gen. plant. ed. Schreb. n. 1321.

Calyx communis imbricatus. *Corolla communis* radiata: *Corollulae* hermaphroditae numerosae, discum planum constituentes. *Feminae* in radio. *Propria* hermaphroditi, infundibuliformis, limbo quinquefido, patulo. *Femina* ligulata, longior, patens, tridentata. *Calyx* fructiferus immutatus. *Semina* oblonga, margine inciso multifido coronata. *Receptaculum* paleaceum, convexum.

324. BUPHTHALMUM *salicifolium*
foliis alternis lanceolatis, subserratis villosis, calycibus nudis, caule herbaceo,
Linn. Sp. pl. 2. p. 1275,

Bupthalmum foliis imis petiolatis, ovatis, superioribus amplexicaulibus lanceolatis, hirsutis, rariter dentatis. *Hall. stirp. helv. n. 118.*

Aster montanus luteus foliis oblongis. *Mapp, fl. Alf. p. 33.*

Ha b. In ditione Basileensi in montosis, *Wasserfallen*, *Dietisberg*, *Farnspurg*, *Homburg* et alibi t. *Lho.* In Principatu superiore prope *Oeningen*, nec non ad Castellum *hentwyl*. In Principatu *Fürstenberg* similis in montosis. In Alsatie montosis prope *Hochandlau* post *Mappu*

Flor. Augusto, Septembri. *Semina* Augusti, Octobri matura. z .

Radix horizontalis, ramosa, fibrigrica. *Caules* ex una radice plures, pedales et altiores, erecti, sulcato-angustifolii, saepe glabriusculi, simplicifolii superne subramosi. *Folia* alterna, imbricatoloba, obovato-lanceolata, obtusa; sessilia, lanceolata, rariter denticulata integra, acuta, glabra, aut villosiuscula terminalis, majusculus, luteus. *Calyx* minusculus, patens. *Squamae* lanceolatae, natæ. *Corollulae* radii ligulatae, patentes quinquedenticulatae; *Disci* infundibuliformes. *Semina* angulata. margine

SYNGENESIA. POLYG. FRUSTR. 493

ota. Ad rariores Floræ cives pertinet, planta
alias magis subalpina.

GEN. POLYG. FRUSTRANEA.

CCCC. HELIANTHUS.

Linn. Gen. pl. ed Schreb. n. 1322.

Calyx communis imbricatus, subquar-
to expansus: *squamis* oblongis, basi latiuscu-
lis hiscentibus undique apicibus. *Corolla*
posita radiata maxima: *Corollulae* herma-
ditae numerosissimae in disco maximo. *Fe-*
ae pauciores, multo longiores in radio.
ria hermaphroditis cylindracea, calyce
nuni brevior, basi ventricosa, orbiculata,
essa: *limbo* quinquedentato, acuto, pa-
e. *Femineis* ligulata, lanceolata, integerri-
longissima. *Calyx* fructiferus subimmutatus.
na oblonga, obtusa, tetragona, angulis
stitis compressa, coronata *paleis* duabus
colatis, acutis, deciduis. *Receptaculum*
iceum magnum, planum: *paleis* lanceola-
cutis, duabus singulum semen distinguen-
e, deciduis.

5. HELIANTHUS *Annuus* foliis om-
nibus cordatis trinervatis, (pedunculis
incrassatis), floribus cernuis. *Linn. Sp.*
pl. 2. p. 1276.

elenium indicum maximum. *Bauh. pin.*
276.

326. HELIANTHUS SOLYCE. FRUIT.

Herb. maxima, *Dodon. pempt.* 264.

Chrysis. *Renealm. spec.* 84. t. 83.

Corona Solis I. *Tabernam. p.* 1146.

Icon. *Regnault. bot.* 371. 101. 1146.

Germ. Grasse Sonnenblume. H. 3000.

Gall. *Helianthe annuelle, Fleur de Soleil*
Tournesol.

Hab. In Peru et Mexico. Colitur passim in her-
tis, agris et vineis.

Flor. Julio, Augusto. *Semina* Septembri, Oc-
tobri matura. ☉

Planta maximis floribus natantibus aureis
biens.

Semina oleum unguentum largiantur. *Re-*
cepta.

Flores apibus melliferi.

326. HELIANTHUS tuberosus

ovato-cordatis triplinerviis. Linn.
pl. 2. p. 1277.

Helenium indicum tuberosum. *Bauh.*
277.

Flos Solis farnesianus. *Col. ecphraf.* 1. p.
t. 13.

Icon. *Jacqu. Hort. Vind. t.* 161.

Germ. Knollige Sonnenblume. *Erdäpfelre-*
gende Sonnenblume.

Gall. *Hélianthe tubéreuse. Topinambour.*
tichaut de Canada.

Hab. In Brasilia. Colitur passim in oleraceis
agris.

Flor. Julio, Augusto. *Semina* Septembri ma-
tura. ☿

SYNGENESIA. POLYG. FRUST. 426

Radix tuberosa esculenta. Flores multo minores quam in praecedente specie, lutei.

Flores apibus melliferi.

Nota. Helianthus multiflorus Linn. praesertim in hortis anglicis Carlsruhanis cum aliis speciebus virginicis, colitur.

CCCCI. CENTAUREA.

Linn. Gen. pl. ed. Schreb. n. 1331.

Calyx communis imbricatus, subrotundus: *sepalis* saepius varie terminatis. *Corolla* composita, flosculosa, difformis: *Corollulae* hermaphroditae plurimae in disco. *Femininae* pauciores, majores, laxae, in radio. *Propriae* hermaphroditi monopetala: tubo filiformi; *Limbo* intricato, oblongo, erecto terminato. *Laciniae* quinque linearibus, erectis. *Femineis* monopetala, infundibuliformis: *tubo* tenui sensim ampliato, recurvo; *limbo* oblongo, obliquo, aequaliter diviso. *Calyx* fructiferus, immutatus, connivens. *Semina* solitaria. *Pappus* plerumque plumosus seu capillaris. *Receptaculum* osum.

CYANI calycinis squamis ferrato-ciliatis.

327 *CENTAUREA phrygia* calycibus recurvato-plumosis, foliis indivisis. Linn. Sp. pl. 2. p. 1287. — Lamarck et Decand. fl. franc. 4. p. 92. — Roth. fl. germ. 3. p. 359.

390 SINGENTISIA. POLYQ. RUBR.

Cyathis foliis subhirsutis; ovato-lanceolatis, amplexicaulibus; dentatis, plumbeis reflexis. *Hall. stirp. helv. n.*

Succa latifolia et angustifolia; capite
Bauh. pin. 271. — Mapp. fl. Alf.

Icon. Oeder. fl. Dan. t. 520.

Germ. Phrygische Flockenblume.

Gall. Centaurée plumbeuse.

Hab. In pratis montosis sylvaticis retro
auf dem Kaltenbrunn, auf der Herr
den Blättle; in convallibus Böhler
fibus, Oppenauensibus, Griesbacensibus
et passim frequens. In Principa
tenbergensi retro Pilsenberg et
pratis montosis. In Alsatia Vogesen
prope Barr et in dem Urbiethal.

Flor. Julio, Augusto. *Semina* Septembri
tura.

Radix ramosa, fibrosa. *Caulis* pedalis, cul-
erectus; angulatus, hirsutus, scaber, ram-
Rami erecti, angulati, foliosi. *Folia*
petiolata; reliqua alterna, sessilia, am-
plexicaulia, ovali oblonga, integra, aut rarite-
tata, firma, hirsutis scabra. *Flores* cauli-
mosque terminantes, solitarii, majusculi
purei, rarius albidi. *Calycinae* squamae
to-ciliatae, plumosae, cinereae, ciliae
vae, longae, elegantissimae.

Nota. Variatio Caule simplici unifloro no-
fundenda cum *C. uniflora* quam in regi-
montosis calidioribus Cataloniae et Galliae
et quae sine dubio nullibi in Germania cre-

Obs. *Centaurea uniflora* Linn. in horto Bo-
Carlsruhano culta, constantissima permansit.
non progenies *C. Phrygiae* alienata.

8. *CENTAUREA nigra* calycibus ciliatis: squamula ovata: ciliis capillaribus erectis, foliis lyrato-angulatis, floribus flosculosis. *Linn. Sp. pl.* 2. p. 1288. — *Poll. fl. pal.* n. 822.

Centaurea foliis radicalibus semipinnatis, caulinis ovato-lanceolatis, caulibus angulatis. *Hall. stirp. helv.* n. 184. an?

Jacea nigra laciniata. *Bauh. pin.* 271. — *Mapp. fl. Als.* p. 150.

Jacea austriaca VI. *Clus. hist.* 2. p. 7. — *n. Oeder. fl. Dan.* t. 996.

m. Schwarz-Kelchige Flockenblume.

l. Jacée noire. Centaurée noire.

l. Utrinque in collibus dumetosis apricis, nemorosis graminosis, ad viarum marginalia passim frequens. Circa *Carlsruhe*, am *Beurtheimer Wald*, im *Hauacker* et alibi passim frequens.

r. Julio, Augusto. Semina Septembri maturatione. ♂

A *Centaurea Jacea* affini facile distinguitur, et firmiore, Foliis magis duris, Calycibus obovatis, flosculis globosis, Squamis arctis, incumbens, nigris, ciliatis: Ciliis erectis, nigris, Flosculis omnibus hermaphroditis, et purpureis.

at. Foliis integris et laciniatis. Capitulis majoribus et minoribus.

lores melliferi.

9. *CENTAUREA montana* calycibus ferratis, foliis lanceolatis decurrentibus, caule simplicissimo. *Linn. Sp. pl.* 2. p. 1289. — *Poll. fl. pal.* n. 823.

2. Bed. Tom. III.

li

Cyanus caule unifloro alato, foliis ellipticis lanceolatis. *Hall. stirp. helv. n. 190.*

Cyanus montanus latifolius, vel Verbascum cyanoides. *Bauh. pin. 273.* — *Mapp. fl. Als. p. 85.*

Cyanus montanus. *Tabernamont. p. 432.*

Icon. Jacqu. fl. Austr. t. 371. — *Blackw. t. 66.*

Germ. Blaue Bergflockenblume.

Gall. Jacée ailée. Centaurée de montagne.

Hab. In montosis altioribus. In Marggr. superiore in monte *Belchen* rarius, in adscensu montis *Blauen* nec non in monte *Feldberg* *Brigoviae*; in monte *Wasserfallen* infrequens. In *Alfatia* retro *Barr* et *Rappolsweyher*; in monte *Ballon*. In *Palatinatu* circa *Lautern* in sylvis umbrosis; prope *Frankenstein*; in sylvis inter *Kirnbeckerbach* et *Rathsweyer*, nec non in monte *Donnersberg* post *Pollichium* vidi.

Flor. Junio, Julio. *Semina* Julio, Augusto matura. ☞

Radix ramosa, fibrosa, fusca. *Caulis* pedalis et altior, erectus, angulatus, villosus, quandoque tomentosus, simplex uniflorus, rarius quandoque ramo uno alterove auctus. *Folia* alterna, per caulem decurrentia, ovali-lanceolata, utrinque villosa, scabra, integra, rarius denticulata, pallide viridia. *Flos* terminalis, erectus, speciosus, solitarius, paucissime ex violaceo caeruleus, raro albus. *Calycinae* squamae ovatae, glabriusculae, margine nigricantes, ciliatae. *Corollulae* radii femineae, praelongae, amoene caeruleae. *Dissectae* hermaphroditae, violaceo-caeruleae.

SYNGENESIA. POLYG. FRUSTR. 499

Nota. In subalpinis individua magis viridia, villosa; in demissioribus magis lanuginosa sunt.

In horto botanico culta, ramosa, multiflora est.

Egregium præsertim in hortis anglicis decus.

Flores apibus melliferi.

30. CENTAUREA *Cyanus* calycibus ferratis, foliis linearibus integerrimis; infimis dentatis. *Linn. Sp. pl. 2. p. 1289.* — *Poll. fl. pal. n. 824.*

Cyanus foliis infimis ellipticis dentatis, supermis linearibus; semisflosculis latissimis. *Hall. stirp. helv. n. 191.*

Cyanus segetum flore caeruleo albo disco coeruleo cum corona candida. *Bauh. pin. 273.* — *Mapp. fl. Als. p. 85.*

Cyanus caeruleus arvensis. *Tabernæmont. p. 428. Icon.*

non. Regnault. bot. — Bulliard. herb. t. 221.
— *Oeder. fl. Dan. t. 993. — Blackw. t. 270.*

erm. Kornblume. Blaue Kornblume.

all. Jacée des blés. Cassé lunette. Bleuret.
Bluet. Barbeau. Aubifoin.

ab. Utrunque inter segetes, in arvis agris et vineis passim frequens.

lor. Junio, Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Flores inodori vulgo pulcherrime coerulei, quædamque albi, rarius in disco caerulei et in radio albi, vice versa.

Multae variationes in hortis pulcherrimæ coluntur.
arm. CYANI Flores.

Pulcherrimum sæpe in agris neglectis cum *Papaver Rhoeas* decus.

Agricolis invisa, ante florescentiam secundam diminuenda.

Flores apibus melliferi.

In odoramentis ficcis Pot-pourri et nigrata.

Pigmentum cæruleum præbent. Gentl. m. a. 1748.

1331. *CENTAUREA paniculata*

bus ciliatis: squamis planis, foliis natifidis, rameis pinnatifidis linearibus caule paniculato. *Linn. Sp. pl.* 2. p. *Pollich. fl. pal. n.* 825.

Cyanus pinnulis foliorum pinnatis linearibus. *Hall. stirp. helv. n.* 187.

Jacea foliis candicantibus lacinjatis, caliculis non splendentibus. *Mapp. fl. p.* 151.

Stoebe major calyculis non splendentibus. *Bauh. pin.* 273.

Centaurii majoris species tenuifolia. *hist.* 3. p. 31.

Jacea II. *Tabernæm. p.* 435. *Icon.*

Icon. Jacqu. fl. Austr. t. 320.

Germ. Rispenförmige Flockenblume.

Gall. Jaccée paniculée.

Hab. In Rheni regionibus sabulosis apertis ad viarum marginalia. Prope *Germ.* et *Philippsburg* copiose. In *Marggra* superiore prope *Neuenburg*, *Zienkenbreybach*; ad montem *Kaiserstuhl* 1 *Limburg* et alibi ad Rhenum frequenter. *Alsatia* in arenosis prope *Marckels*, *Othmarsheim* et alibi versus Rhenum.

Palatinatu ad vias circa *Worms*, *Franckenthal*, *Klein-Niedesheim*, *Oppenheim*, et alibi copiose.

o r. Junio, Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ♂.

Radix ramosa, dura. *Caulis* sesquipedalis, bitalis, tripedalis et altior, erectus, quaque decumbens, angulatus, striatus, firmus, canus, scabriusculus, a basi sæpe ad apicem noso-paniculatus. *Folia* sparsa, alterna, sessa, ad caulem sæpe reflexa, sicca, incanoidia, scabriuscula, pinnata: pinnis profunde pinnatifidis, angustis, linearibus; *summa* ramea pinnatifida, integerrimaque, linearia, vel oblonga. *Flores* caulem ramosque terminantes ciliati, numerosi, mediocres, pallide purpurescentes, rarius albid. *Calycinae* squamato-lanceolatae, adpressae, erectae, incanæ, virgine fusco-ciliatæ. *Semina* pappo brevi inæquali, capillari coronata. *Receptaculum* villosum.

In præcipitiis ad Rhenum sabulosis arenam figit.

Flores melliferi.

32. *CENTAUREA Scabiosa* calycibus ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Linn. Sp. pl.* 2. p. 1291. — *Pollich. fl. pal.* n. 826.

Cyanus foliis pinnatis, pinnis dentatis obtusis, squamis calycinis triangulis ciliatis. *Hall. stirp. helv.* n. 186.

Jacea vulgaris laciniata, flore purpureo et albo. *Mapp. fl. Alf. p.* 150.

Scabiosa major squamatis capitulis. *Bauh. pin.* 269.

Jacea III. *Tabernaemont. p.* 435. *Icon.*

talis, tripedalis et altior, erectus, quam
 subdeclinatus, angulatus, striatus, hirsu-
 tus, simpliciusculus, aut superne ramosus
 radicalia et caulina inferiora petiolata,
 pinnatifida, laciniata, dentata; *superiora*
 fida: *segmentis* integriusculis, omnia
 pilosaeve hirsuta, laete viridia. *Flores* ter-
 recti, longe pedunculati, magni, dilu-
 puri, rarius albi. *Calycinae* squamæ
 arcte incumbentes, villosiusculae, mar-
 gatae, fusco-nigricantes, rarius fusco-
Corollulae radii, disci multo longiores,
 Pappo parvo inæquali coronata.

Flores melliferi.

† Rha pontica *calycinis squamis men-*
ceis, aridis, integris, muticis.

1333. CENTAUREA *Jacea* calyci
 riosis laceris, foliis lanceolatis:
 libus sinuato-dentatis, ramis ar-
 ...

SYNGENESIA. POLYG. FRUSTR. 503

Jacea I. *Tabernaemont. p. 435. Icon.*

3. Jacea nigra angustifolia, lithospermi arvensis foliis, caule aspero. *Bauh. pin. 271.*
— *Prodr. p. 127.*

con. Bulliard. herb. t. 227. — Oeder. fl. Dan. t. 519.

herm. Gemeine Flockenblume.

all. Jacée des prés.

ab. Utrisque in pratis, aggeribus, ad viarum marginalia frequens.

lor. Junio, Julio, Augusto, Septembri. Semina Augusto, Septembri, Octobri matura. 7

Radix ramosa, fibrosa. *Caulis* pedalis, sessilis, et altior, erectus, sulcatus, superangulatus, villosusculus, ramosus. *Folia* ternata, semiamplexicaulia, sessilia, lanceolata, inferiora saepe inciso dentata, *summa* integra, omnia laete viridia, quandoque ex viridi-incana, aspera. *Flores* caulem ramosque terminantes, erecti, solitarii, purpurei, pro albidi, majusculi. *Calycinae* squamae ovalimucronatae, obtusae, aridae, margine fuscae, aut nigricantes, lacerae, scariosae.

variat Foliis latioribus et angustioribus fere linearibus, Calycibus fuscis et nigricantibus, minoribus et majoribus,

Nota. Variatio calycibus majoribus nigricantibus non confundenda cum *Centaurea nigra* affini, a qua differt calycibus non subglobosis majoribus, sed ovalibus minoribus. Hanc variationem sine dubio Hallerus sub n. 184. descripsit. Est *Centaurea dubia* Sutter. fl. helv. 2. p. 202.

334. CENTAUREA *amara* calycibus scariosis, caulibus decumbentibus, foliis

... Cyanus repens latifolius. *Bauh. pin*
Cyanus repens. *Lobel. Icon. p. 548.*

Germ. Bittere Flockenblume.

Gall. Centaurée amère. Jacée amère.

Hab. In Brisgoviae collibus macilent
am *Kaiserstuhl* ad arcem *Limbur*
rarius. In Alsatia in pratis siccis ma
circa Argentinam versus *Offwald*
... *Eckolsheimer Wald t. Mappo.*

Flor. Junio, Julio, Augusto, Septemb
na Septembri, Octobri matura. 7

Nimis omnino affinis præcedenti
a qua mere differre videtur, *Caulibus* d
ribus decumbentibus, simpliciusculis, u

Vix pro distincta specie declaranda.

††† *Calcitrapae calycinis spinis comp*

1335. *CENTAUREA Benedicta*

SYNGENESIA. POLYG. FRUSTR. \ 505

Carduus sylvestris hirsutus f. **Carduus benedictus.** *Bauh. pin.* 378.

Carduus benedictus. *Tabernæm. p.* 1063. *Icon.*

Icon. Blackw. t. 476.

Ver m. Benedikten - Flockenblume.

Wall. Centaurée chardon-béni.

Tab. In Oriente et Hispania ad verfuras. Colitur passim in rusticorum hortis.

Flor. Junio, Julio. *Semina* Augusto matura. ☉

Calyx foliaceo-involucratus, lanatus. *Flos* levis.

Ver m. CARDUI BENEDICTI Herba.

Herba amara, stomachica, roborans.

336. CENTAUREA Calcitrapa calycibus subduplicato-spinosis sessilibus, foliis linearibus pinnatifidis tateribus dentatis caule piloso. *Linn. Sp. pl.* 2. p. 1297. — *Poll fl. pal.* n. 828.

Calcitrapa foliis hirsutis pinnatis, pinnis dentatis, floribus in alis sessilibus. *Hall. stirp. helv.* n. 194.

Carduus stellatus f. **Calcitrapa.** *Bauh. hist.* 3. p. 89.

Spina stella. *Tabern. p.* 1080. *Icon*

Icon. Regnault. bot.

Ver m. Stern - Flockenblume. Calcitrape.

Wall. Calcitrape. Chauffe - trape étoillée.

Tab. In Rheni campestribus sabulosis macilentis hinc inde abunde. Prope *Dachsland, Au, Kehl, Altbreisach* et alibi. Circa Argen-

306 SYNGENESIA POLYG. FEUSTE

tinam am Schief-Rein et alibi abunde. |
Palatinatu circa Heidelberg, Worms, O
penheim, inter Dürkheim et Lamshei
in pascuis ficcis, et alibi passim abunde.

Flor. Julio, Augusto. *Semina* Septembri m
tura. ☉

Caulis sesquipedalis, bipedalis, erectu
erassiusculus, teres, pilosus, ex viridi albidu
ramosus. *Rami* numerosi, alterni, divarica
divisi. *Folia* radicalia plurima, pinnatifida, ma
na; *caulina* alterna, sessilia, semiamplexico
lia, profunde pinnatifida: *pinnis* lineari lance
latis longis, integris dentatisve, inermibus, p
briusculis. *Costa* folii dorsalis alba. *Flores*
lares, terminalesque, sessiles, solitarii, pale
carnei, seu albi. *Calyx* ovatus, rigidus. *Squa*
calycinae apice longe spinosæ; *Spinæ* div
gentes validæ, pungentes, stellam imitant
albidæ, basi spinulis rigidiusculis armatæ. *S*
mina pappo destituta. *Receptaculum* villosu

Pharm. CALCITRAPÆ Radix; Herba; Semina

Planta amara, saponacea, resolvens, tonica, di
retica.

Plantæ efficaciam in febribus intermittentibus
ab antiquis temporibus ex Baptista Sardo mon
rat J. Bauhinus.

Succus herbæ expressus, pulvis vel extractu
præcipue in Gallia contra febres intermittentes exhibe
tur; quantum in hisce sanandis valeant in nosocomio
Veredanenſi in frequenti ægrotorum numero An. 178
exploratum a Cl. Plouet, vid. Buchoz. Diff. sur
la Chauffe-trappe, reconnue comme spécifique
dans les fievres intermittentes 1787. c. fig.

Apud nos nimis a medicis neglecta, vix cognita.

Condimentum Agni paschalis apud Judæos fuit.

Caules juniores edules in Aegypto. Forskhl
in Niebuhrs Beschreib. von Arabien, præf.
p. 44.

537. CENTAUREA solstitialis calycibus duplicato-spinosis solitariis, foliis rameis decurrentibus inermibus lanceolatis, radicalibus lyrato-pinnatifidis, *Linn. Sp. pl. 2. p. 1297. — Lamarck et Decand. fl. franc. 4. p. 103.*

Calcitrapa caule alato foliis tomentosis, radicalibus femipinnatis, caulinis lanceolatis dentatis. *Hall. stirp. helv. n. 193.*

Carduus stellatus luteus foliis cyani. *Bauh. pin. 387.*

Spina solstitialis. *Bauh. hist. 3. p. 91. Icon. Germ. Gelbe Sommer-Flockenblume.*

11. Centaurée du solstice.

12. In ditionis Basileensis agris ante portam Eschianam et extra portam St. Johannis et circa Hüningen t. Lachenalio.

13. Junio, Julio. Semina Augusto matura. ☉

Caulis pedalis, sesquipedalis, erectus, angulosus, foliis alatus, ramosus, hirsutus. *Folia* infima lyrato-pinnatifida, lobis lanceolatis, extremo maximo; *caulina* alterna, decurrentia, elliptico-lanceolata, dentata, summa linearia, integra, omnia hirsuta. *Flores* caulem ramosque terminantes, solitarii, flavi, erecti. *Calycinae* quatuor exteriores aculeatae, sequentes aculeo minore, basi aculeolato praeditae, intimae mites.

Nota. Planta in regionibus calidioribus Galliae, Italiae et Hispaniae in agris abunde crescens.

**SYNGEN. POLYGAM. NECESSARIA,
CCCCII. CALENDULA.**

Linn. Gen. pl. ed. Schreb. n. 1339.

Calyx communis simplex, polyphyllus, erectus: segmentis lineari-lanceolatis, sub-

lata, tridentata, basi villosa. *Calyx* fructu connivens, subrotundus, depressus. *Corollae* disci nulla: *peripheriae* propria, medio obcordata, compressa. *Femineis* majoribus longioribus, incurva, triangularia, angulis mucronatis. *Pappus* nullus. *Receptaculum* nunc planum.

1338. *CALENDULA arvensis* sen
cymbiformibus muricatis incurva
timis erectis protensis. *Linn. Sp. p.*
— *Poll. fl. pal. n.* 829.

Calendula foliis lanceolatis amplexica
dentatis, floribus concoloribus.
stirp. helv. n. 94.

Caltha arvensis. *Bauh. pin.* 275.

Icon. Bulliard. herb. t. 239.

Germ. Kleine Ringelblume.

Gall. Souci des champs.

Hab. In arvis, agris et vineis passim. In
prope *Durlach* et *Grözingen*. In Pal:

SYNGENESIA. POLYG. NECESS. 509

Caulis digitalis, palmaris, et altior, ad-
 endens, angulosus, ramosus. *Folia* alterna,
 simplicicaulia, cordato-ovata, acuminata,
 argine subsinuato-dentata, scabriuscula, pu-
 bescentia. *Flores* terminales, solitarii, aurantii,
 ediocres. *Corollulae* radii, ligulatae, patentes.
Albicinae squamæ parallelæ, lineares, villosi-
 sculae. *Semina* aculeata; exteriora longiora,
 parva, rectiora; interiora breviora, incurva,
 affiora, pappo destituta.

Nota. Calendula officinalis feminibus omnibus
 incurvatis, floribus multo specioribus aurantiis,
 in hortis passim colitur.

YNGEN. POLYGAM. SEGREGATA.

CCCCIII. ECHINOPS.

Linn. Gen. pl. ed. Schreb. n. 1353.

Calyx communis polyphyllus: *squamis* su-
 peris reflexis, multos flores continens. *Peri-
 thium* partiale uniflorum, oblongum, imbric-
 tum, angulatum; foliolis subulatis, superne
 erectis, persistentibus. *Corolla* monope-
 ala, longitudine calycis, tubulosa: *limbo* quin-
 cefido, reflexo-patente. *Calyx* fructiferus im-
 mitatus, major. *Semen* unicum, ovato-oblon-
 gum, basi angustius: apice obtuso. *Pappus* ob-
 tetus. *Receptaculum* commune globosum, se-
 dum.

539. *ECHINOPS sphaerocephalus*
 capitulis globosis, foliis pubescentibus.
Linn. Sp. pl. 2. p. 1314. — Lamarck et
Desand. fl. franc. 4. p. 71.

Bauh. pin. 381.

Ritro seu Rutro Theophrasti. *Lob.*
2. p. 8.

Ger m. Kugeldistel.

Gall. Echinope a tête ronde.

Hab. In Brisgovia am Kaiserstuhl pro-
gen t. generosissimo Jitnero.

Flor. Julio, Augusto. Semina Septem-
tura. ♀

Caulis 2-3 pedalis, erectus, sulcatus,
erectus, ramosus. Folia alterna, pinnatifid-
gine spinulosa, superne viridia, subtus
tomentosa. Capitula caulem ramosque
nantia, erecta, solitaria, capitata, r-
Flosculi albi.

Flores melliferi.

Ad rariores omnino Florae cives pertine-
alias in calidioribus Galliae et Hispaniae regionibus

SYNGENESIA MONOGAMI

CCCCIV I A S I O N E

SYNGENESIA. MONOGAMIA. 311

Genus. *Perianthium* proprium quinque-dentatum, superum, persistens. *Corolla* propria monopetala, regularis, profunde quinquesida: *laminae* lanceolatis erectis. *Capsula* subrotunda, quinqueangularis, coronata calyce proprio, bibilocularis, apice foramine rotundo dehiscens. *Dissepimentum* ad axem divisum. *Semina* parva subovata. *Receptaculum* subglobosum, pedicellatum, liberum, in basi capsulae.

340. *JASIONE montana* foliis linearilanceolatis obsolete serratis, floribus terminalibus in capitulum congestis.

Jasione montana. Linn. *Sp. pl.* 2. p. 1317. — Pollich. *fl. pal.* n. 834.

Rapunculus foliis linearibus subsperis, spica planiuscula, petalis liberis. Hall. *fl. stirp. helv.* n. 678.

Rapunculus scabiosae capitulo coeruleo. Bauh. *pin.* 92. — Mapp. *fl. Als.* p. 267.

Scabiosa minor. Tabern. *p.* 444 *Icon.*

Mon. Curtis. *fl. Lond.* t. 243. — Oeder. *fl. Dan.* t. 319. — Schkuhr. *bot. hand.* t. 266.

erm. *Jasione.* Acker-*Jasione.* Berg-*Jasione.* all. *Jasione ondulée.* *Jasione des montaignes.*

a b. Utrunque in pascuis macilentis graminibus apricis, collibus aridis, arvis et aggeribus neglectis, sylvis, ericetis, et juniperitis sabulosis, passim frequens.

lor. Julio, Augusto, Septembri. *Semina* Septembri, Octobri matura. ☉

Radix fibrosa, albida. *Caules* ex una radice saepe plures, palmares, spithamei, pedales

311 SYNGNESIA MONOGAMIA

et altiores, erectiusculi, ramosi, hirsuti. Folia sparsa, alterna, sessilia, linearilanceolata, obtusiuscula, integriuscula, undulata, hirsuta. Flores caulem ramosque terminantes, capitati, corulei aut albi.

Bonum et gratum ovibus pabulum.

Flores melliferi.

In sabulosis arenam figit.

CCCCV. V I O L A.

Linn. Gen. pl. ed. Schreb. n. 1364.

Calyx pentaphyllus, brevis persistens. *Corolla* pentapetala; irregularis: *petalis* interioribus. *Petalum* supremum rectum, deorsum spectans, latius, obtusius emarginatum, densum basi in *Nectarium* corniculatum, obtusum, inter calycis foliola prominens. *Filamenta* quinque minima. *Antherae* subconnexae, membranaceae ad apicem auctae. *Capsula* ovata, trigona, obtusa, unilocularis, trivalvis. *Semina* plerumque, ovata, appendiculata.

† *Acaules.*

1341. *VIOLA hirta* acaulis, foliis cordatis, piloso-hispidis. *Linn. Sp. pl.* 2. p. 1324. — *Poll. fl. pal.* n. 831. — *Habstirp. helv.* n. 559.

Viola hirta, acaulis, foliis petiolisque hirsutis, bracteis infra medium pedunculi. *Curtis. fl. Lond.*

Viola

SYNGENESIA. MONOGAMIA. 313

Viola martia hirsuta inodora. *Morif. hist.* 2. p. 475. — *Sect. 5. t. 35. f. 4.*

Viola Trachelii folio vulgo. *Raj. hist.* 1051. *n. Curtis. fl. Lond.* — *Oeder. fl. Dan. t. 618. r m. Rauhharriges Veilchen.*

11. *Violette velue.*

b. In nemoribus et campestribus sabulosis passim non infrequens. Circa *Carlsruhe* in nemoribus. ¶

Radix articulata, fibrosa. *Stolones* nulli. *Radicalia* plurima, longe petiolata, cor-
a, oblonga, obtuse crenulata, utrinque hir-
juniora pilosa. *Petioli* hirti. *Pedunculi* ra-
ales, erecti, digitales, et altiores, glabrius-
i, medio *Bracteis* binis muniti, uniflori,
erne tetragoni, nutantes. *Corolla* majuscula,
urate vel dilute violacea, inodora.

42. *VIOLA palustris* acaulis, foliis re-
niformibus. *Linn. Sp. pl.* 2. p. 1324. —
Poll. fl. pal. n. 836. — *Hall. stirp. helv.*
n. 560.

Viola palustris rotundifolia, glabra. *Morif.*
hist. 2. p. 475. — *Sect. 5. t. 35. f. 5.*

n. Curtis. fl. Lond. t. 217. — *Oeder. fl.*
Dan. t. 83.

r m. Sumpf-Veilchen.

11. *Violette des marais.*

b. In montosis udis *Sphagno impletis*, et
demissioribus locis passim turfáceis, auf dem
Kaltenbrunn, der *Herrnwiese*, auf dem
Plättie, in pratis udis in convallibus *Büh-*
lerthalensibus, *Griesbacensibus*; in monte
Kniebis an der Schwedenschanze et alibi.

prope *Lantern* in pratis paludosis
vae *Reiswald* vocatæ paludibus, *Reiswald*
versus situatis. t. *Pollichio*.

Flor. Aprili, Majo, Junio. *Semina* Ju
lio matura. 7

Radix repens, dentata, fibrosa.
radicales plurimæ, ovato-acutæ, ferru
lia radicalia, petiolata, subreniform
nata, venosa, tenera, nitida, subtus sæp
purpurascentia. *Pedunculi* radicales, p
petiolis multo longiores, erecti, subte
teneri, *Bracteis* duabus suboppositis
latis, basi ferrulatis infra medium instrui
flori. *Flos* medipetris, terminalis, nuta
dus, lineis purpureis ramosis, pulch
pictus. *Calcar* breve, obtusum.

Obs. Planta in horto botanico culta, muta
succubuit.

1343. *VIOLA odorata* acaulis, fo
datis, stolonibus reptantibus. *L*
pl. 2. p. 1324. — *Pollich. fl. pal*
— *Hall. stirp. helv.* n. 558.

SYNGENESIA. MONOGAMIA. 515

n. *Curtis. fl. Lond.* — *Bulliard. herb. t.* 169. — *Regnault. bot.* — *Blackw. t.* 55.

r m. *Merzen-Veilchen. Wohlriechende-Veilchen.*

l l. *Violette de Mars. Violette odorante.*

s b. Utrunque in sepibus et ad viarum marginalia aprica collicosa et alia passim frequens.

o r. Febuario, Martio, Aprili. *Semina* Aprili, *Majo* matura. ¶

Stolonibus reptantibus, *Floribus* suaviter oratis, facile a *V. hirta* distinguitur.

riat. Corollis violaceis, purpureis, carneis, rubellis, et albis.

In hortis obvia floribus impletis pulcherrimis.

arm. VIOLAE ODORATAE s. MARTIAE Flores; Herba; Semina; Radix.

Radix emetica vires habet.

Flores melliferi, apibus accepti.

†† *Caulescentes.*

544. *VIOLA canina* caule adultiore ascendente, foliis oblongo-cordatis. *Linn. Sp. pl. 2. p.* 1324. — *Poll. fl. pal. n.* 838.

Viola caule procumbente ramoso, foliis petiolatis, cordatis. *Hall. stirp. helv. n.* 563.

Viola martia inodora sylvestris, foliis majoribus et rotundioribus. *Bauh. pin.* 199. — *Mapp. fl. Alf. p.* 326.

Viola martia inodora sylvestris foliis mucronatis oblongis et strictioribus. *Bauh. pin.* 199. — *Mapp. l. c.*

Viola canina. *Tabern. p.* 681. *Icon.*

Viola martia alba inodora. *Bauh. pin.* 199. *Mapp. fl. Alf. p.* 326.

Icon. Curtis. fl. Lond. t. 108.

Germ. Hunds-Veilchen.

Gall. Violette sauvage.

Hab. Utrunque in sylvis, nemoribus, sepil
passim frequens.

Flor. Martio, Aprili, Majo. *Semina* Ma
Junio matura. 7

Radix dentata, obliqua, fibrosa. *Ca*
digitalis et altior, adscendens, laevis, sub
gulosus, folia floresque ferens. *Folia* petiola
cordata, acutiufcula, crenata, lævia. *Stipul*
caulinae lanceolatae, acutae, ereetae, fer
latae. *Flores* in pedunculis tetragonis, erect
superne nutantibus, solitarii, majusculi, viol
ceo-purpurei, aut albidii, inodori.

Pharm. VIOLÆ CANINÆ Radix.

Radix subemetica vix usitata. conf. Niemey
Diff. de Violæ caninae in medicina
Götting. 1785.

In officinis pharmaceuticis sæpe hisce violis Sy
pum Violarum odoratarum corrumpunt.

1345. *VIOLA montana* caule erecto, fo
liis cordatis oblongis. * *Linn. Sp. pl.*
p. 1325.

Viola caule lato erecto, foliis ellipticis cre
natis, stipulis semipinnatis. *Hall. stir*
helv. n. 567.

Viola flore cæruleo longifolia. *Riv. t. 10*

Viola erecta flore cæruleo et albo. *Mon*
hist. 2. p. 475. — Sect. 5. t. 7. f. 7.

Viola arborescens. *Camerar. Epit. 911.*

Viola martia arborefcens purpurea. *Bau*
pin. 199.

Viola martia arborescens. *Tabernaemont.*
p. 681. *Icon.*

erm. Blaue Bergveilchen.

a ll. Violette de montagne.

a b. Prope *Dachsland* in dumetis rarius. Circa Argentinam passim in sepibus *im Ostwinkel* et alibi An. 1779. et sequentibus annis vidi; nec non intra *Kehl* et *Goldschier*, rarius. In *Vogesi* montosis sylvaticis prope *Hochandlau*. In Palatinatu prope *Heidelberg* in dumetis rarius.

Caules ex una radice plures, herbacei pedales, sesquipedales, erecti, simplices, angulati, glabri, tenues. *Folia* alterna, petiolata, inferiora cordato-ovata; superiora cordato, vel ovato-lanceolata, serrata, late viridia, glabra. *Stipulae* magnae, dentato-serratae, magis minusve incisae. *Flores* axillares, longe pedunculati. *Pedunculi* solitarii angulati, nudi, univiri. *Corolla* pallide caerulea, venis saturationis violaceo-caeruleis picta, majuscula, odora.

b f. Planta in Hispania a me lecta, differt a nostrata *Caulibus* ex radice multo copiosioribus longioribus, 3-4 pedalibus, firmioribus fere lignosis; *Foliis* majoribus. *Corollis* speciosioribus pallide caeruleis. In Horto Botanico Carlsruhano ex feminibus culta per quatuor annos vix mutata; post hac sensim humilior, in omnibus partibus nostrati simillima evasit.

346. *VIOLA persicifolia* caule erecto, foliis ovato-lanceolatis, serratis. *Böhm. fl. Lips. n. 456.* — *Hoffm. fl. germ. p. 311.* — *Roth. fl. germ. 2. p. 271.*

Viola lancifolia. Lamarek et Decand. fl. franc. 4. p. 807.

**Viola palustris angustis periclymenae
cristatis et serratis. Rupp. fl. Jen.**

**Viola sylvestris longifolia. Taberna-
p. 682. Icon bona.**

Icon. Schkuhr. bot. handb. t. 269.

Germ. Pferfingblatt-Veilchen.

Gall. Violette fer de lance.

**Hab. Circa Linkenheim, Hochstädt e
heim in arvis humidis sabulosis, fa-
undatis rarius, ubi vidi cum b. c.
ckerto.**

Flor. Majo, Junio. Semina Julio ma

Affinis præcedenti, a qua differt,
bus demissioribus, digitalibus, semipe-
Folius multo angustioribus, omnibus ov-
ceolatis, Stipulis minoribus, Corollis mi-
pallidioribus, ex cæruleo-albidis. Por-
natali arvenſi humido, inundato; no-
dumetoso.

Obs. Planta a b. Schwevckerto in hori

SYNGENESIA. MONOGAMIA. 519

Viola floribus radicalibus corollatis aborti-
entibus, caulinis apetalis seminiferis.
Hort. Cliff. p. 427.

Viola montana latifolia, flores ex radice,
semina in cacumine ferens. *Dill. hort.*
Elth. p. 408. t. 303. f. 390. — Cat. pl.
Giff. app. p. 36.

non. Jacqu. fl. Austr. t. 19. — Oeder. fl. Dan.
t. 1045.

erm. Wunderbares Veilchen.

all. Violette admirable.

a b. In Marggraviatu superiore in monte pro-
pe *Crenzach*. In ditione Basileensi prope
Chrschonen, supra *Arisdorff*. In Princi-
patu superiore in montosis sylvaticis apri-
cis retro *Oeningen*. In Principatu *Fürsten-*
*bergen*si prope *Engen* in montosis ad *Krü-*
genthal et alibi nascentem abunde nuper
vidi.

lor. Aprili, Majo. *Semina* Junio matura. 7

Radix dentata, fibrosa multiceps. *Caulis*
erectus, palmaris, erectus, acute triangulus,
obtusiusculus. *Folia* radicalia numerosa, cæspi-
osa, petiolata, reniformi-cordata, subcucul-
ata, acuta, crenulata, utrinque villosiuscula,
petiolina 1-2 alterna, minora. *Flores* caulini,
simplices, apetalæ, seriores, fertiles; *radicati*
steriles, numerosi, petalis instructi, albidi, aut
purpureo-violacei: *Petalum* infimum purpu-
reum striis pictum, pedunculati. *Pedunculi* his-
pidi, trianguli, villosi, uniflori. *Stipulae* ad
basin pedunculorum caulinarum, lanceolatae,
integrae. *Semina* ovata, nitida, fus-
cescentia.

os. Vidi in Principatu *Fürstenbergen*si prope
Krügenthal specimina nostratibus villosiora,

4. p. 808.

Viola caule debili, paucifloro, foliis
latis, reniformibus obtuse dentatis
stirp. helv. n. 564.

Viola alpina rotundifolia minor lutea
cken. alm. 388. t. 233. f. 1.

Viola alpina rotundifolia, lutea. *Ba*
199. — *Mapp. fl. Alf. p. 327.*

Viola montana L. *Clus. hist. 1. p. 309*
Icon. Oeder. fl. Dan. t. 46.

Germ. Gelbes zweyblüthiges Veilchen.

Gall. Violette à deux fleurs.

Hab. In Alsatiaë Vogelis summis montium
retro *Dietselhausen* versus *dem weiff*
t. Mappo.

Flor. Julio, Augusto. *Semina* Septem-
tura. ¶

Caulis digitalis, palmaris, decumbens
doque erectiusculus, tener glabriusculus
folius, subunislorus. *Folia* caulina altera

SYNGENESIA. MONOGAMIA. 521

Planta alias alpina, in monte Jura abundans, nec non in pyrenaeis lecta, semper bifolia, saepissime uniflora, rarius biflora. Idem vidi in plantulis in horto botanico cultis.

Ad rariores omnino Florae cives pertinet.

Stipulis pinnatifidis; Stigmate urceolato.

9. *VIOLA arvensis* caule inferne diffuso, angulato, foliis oblongis retusis, stipulis pinnatifidis, corolla calycis fere longitudine. *Willd. prodr. fl. Berol. p. 282.* — *Lamarck et Decand. fl. franc. 4. p. 808.* — *Roth. fl. germ. 2. p. 273.*

Viola tricolor. a. *Linn. Sp. pl. 2. p. 1326.*
Pollich. fl. pal. n. 839.

Viola caule diffuso, ramoso, foliis ovatis, dentatis, flore calyce paulo majori. *Hall. stirp. helv. n. 569.*

Viola bicolor arvensis, flore candido et luteo - flore coeruleo et luteo - flore coeruleo et candido. *Bauh. pin. 200.* — *Mapp. fl. Als. p. 326.*

Viola trinitatis minor. III. *Tabernaemont. p. 690. Icon.*

1. *Curtis. fl. Lond. (sub V. tricolore.)*

m. *Acherviole. Ackerfreyssamkraut.*

l. *Violette des champs.*

. Utrunque in agris, cultis, arvis, vineis passim frequens.

. Majo, Junio, Julio, Augusto. *Semina* Julio, Augusto, Septembri matura. ☉

A sequente specie facile dignoscitur, *Caulerectoribus, tenuioribus, Corollis multo* ribus, *vix calycem superantibus,*

1350. *VIOLA tricolor* caule
diffuso, foliis oblongis, retusi-
lis pinnatifidis, corolla calyce ma-
jori. *Willd. prodr. fl. Berol.* 1
Lamarck et Decand. fl. franc.
— *Roth. fl. germ.* 2. p. 272.

Viola tricolor var. β . *Linn. Sp. pl.* 1
— *Poll. fl. pal.* n. 839. β .

Viola caule angulato diffuso, foli-
dentatis, flore calyce duplo
Hall. stirp. helv. n. 568.

Viola tricolor hortensis repens. *B*
199.

Viola trinitatis. I. II. *Tabernæmor.*
Icon.

Icon. *Oeder. fl. Dan.* t. 623. — *Blau*
Germ. Dreyfarbiges Veilchen. Gart-
samkraut. Dreyfaltigkeits-Veilchen-
mütterchen.

CLASSIS XX.
YNANDRIA.

DIANDRIA.

CVII. ORCHIS. *Nectarium* corniculatum.

CVIII. SATYRIUM. *Nectar.* scrotiforme.

CIX. OPHRYS. *Nectar.* subcarinatum.

CX. SERAPIAS. *Nectar.* ovatum subtus
gibbum.

CXI. CYPRIPIEDUM. *Nectar.* inflato-ven-
tricosum.

HEXANDRIA.

CXII. ARISTOLOCHIA. 6-gyna. *Cal* nul-
lus. *Cor.* 1-petala. *Capsf.* sex-
locularis.

giore, foliis rubovatis, impunis
Linn. Sp. pl. 2. p. 1325. — Lan
Decand. fl. franc. 4. p. 810.

Viola alpina purpurea, exiguis foli
stirp. helv. n. 566: a. tab. 17. p.
Bauh. pin. 199.

Viola montana caerulea, grandiflora
nesf. inst. p. 420.

β. *Viola montana tricolor odoratissima*
pin. 199. — Mapp. fl. Alf. p.
Clus. hist. 1. p. 310.

γ. *Viola montana grandiflora, foliis c*
flore variegato. Mapp. fl. Alf. p.

δ. *Viola montana alba grandiflora. H*
var. III.

Germ. Langgesporntes Veilchen.

Gall. Violette à long éperon,

Hab, In montibus Feldberg et Kande
viae. β. In excelsi montis Vogesi
des Vogelsbergs Belchen; in c

SYNGENESIA. MONOGAMIA. 525

tert, *Stipulis* erectis, non pinnatifidis, sed articulatissimis integrifusculisve, *Calcare* longiore, *lycinis* foliolis inferioribus postice ferratis. *Corolla* etiam speciosa purpureo-violacea, trilobor, variegata, et albida, odorata.

Nota. Alpina alias planta, ad rariores omnino florae cives pertinet.

Non confundenda cum *V. cornuta*. Linn. quae ille distinguitur Caulibus longioribus, ramosis, foliis radicalibus caulinisque cordato-ovatis obtusis, puberulis, villosis, Stipulis subquinquefidis, aut indentatis, Pedunculis longioribus, Nectario longe sublongiore.

»f. *Viola cornuta*, in pyrenaeis abs me lecta, in hortum botanicum Carlsruhanum transplantata, mutationi non succubuit.

CCCCVI. IMPATIENS.

Linn. Gen. pl. ed. Schreb. n. 1365.

Calyx diphyllus, minimus coloratus, deducens. *Corolla* 4-5 pedala, ringens. *Petalis* aequalibus. *Nectarium* monopetalum, cuculli fiat fundum floris recipiens, calcaratum. *Filamenta* quinque brevissima, incurva. *Antherae* si divisae, superne connatae. *Capsula* monocularis, quinquevalvis, elastice diffiliens, valvis in spiram convolutis. *Semina* plura, subrotunda, receptaculo columnari affixa.

353. IMPATIENS *Noli tangere* pedunculis multifloris solitariis, foliis ovatis, geniculis caulinis tumentibus. *Linn. Sp. pl. 2. p. 1329. — Poll. fl. pal. n. 840.*

Orchis radicibus subrotundis, spica d
calcare longissimo, labello tricornia
liter tripartito. *Hall. stirp. helv. n.*

Cynosorchis militaris montana, spica rul
conglomerata. *Bauh. pin. 81. prodr.*

Orchis militaris montana, spica rubente
glomerata. *Mapp. fl. Alf. p. 215.*

β. Orchis militaris montana spica congl
rata alba. *Mapp. l. c.*

γ. Orchis militaris montana, spica congl
rata ex albo et purpureo - varie
Mapp. l. c.

*Icon. Jacqu. fl. Austr. t. 266. — Hall.
t. 35. f. sinistra.*

*Germ. Pyramidenförmiges Knabenkraut.
Gall. Orchis pyramidal.*

Hab. In collibus graminosis retro Grötzin
prope Weingarten et Bruchsal cum
β. et γ. In Marggraviatu superiore retro
purg in prato Hexmatte et prope Gen
bach t. Vulpio. In montosis herbidis n
tis Muteli. t. Bauhino. In Alsatia prope L
in collibus montosis gramineis; in pi
prope Haslach versus Niedeck.

Flor. Majo, Junio. *Semina* Julio matura.

Caulis sesquipedalis et altior, erectus. S
in planta juniore conica, conferta, in ad
ovato-pyramidalis. *Corollae* amoene purpur
rarius albae, albido-purpureo-variegatae.
Lobum inferius trifidum: segmentis subrotun
intermedio subangustiore. *Calcar* cylindric
longum.

Nota. Planta omnino elegans, amans colles apr
graminosas, quae apud nos hodie maxime
parte Viticulturae destinantur, hinc prod





Orchides, Satyria, aliaeque plantae pulcherrimae, hisce in locis tam copiose crescentes, intra viginti annos facile rarissimae erunt.

f. Plantam nostram a Mappo ad Argentinam in dem Ruprechtsauer Wald citatam, frustra ibi quaesivi.

56. *ORCHIS globosa* bulbis indivisis, nectarii labio trifido, medio trilobo, cornu brevi, petalis apicē subulatis. *Linn. Sp. pl. 2. p. 1332. — Lamarck et Decand. fl. franc. 3. p. 245.*

Orchis radicibus subrotundis, spica densissima, petalis exterioribus aristatis. *Hall. stirp. helv. n. 1272.*

Orchis flore globoso. *Bauh. pin. 81.*

n. *Jacqu. fl. Austr. t. 265. — Hall. l. c. t. 27. f. sinistra.*

m. *Kugelartiges Stendelkraut.*

l. *Orchis globuleux.*

b. In ditione Basileensi in monte *Vogelberg* versus *Baswang*, nec non in summis palcuis montis *Ballon* Alsatiae t. *Lachenalio.*

or. Junio, Julio. *Semina* Augusto matura. ¶

Caulis pedalis, cubitalis, erectus, foliosus. *lia* lanceolata. *Spica* pyramidali - globosa, fertissima, brevis. *Corollae* laete purpureae, spinatae. *Petala* apicē aristata. *Calcar* ger-
ne brevius.

Ad rariores Florae cives pertinet, copiose in Hel-
ae alpinis obveniens.

57. *ORCHIS coriophora* bulbis indivi-
visis, nectarii labio trifido reflexo, cre-

nato : cornu brevi, petalis conniventibus
Linn. Sp. pl. 2. p. 1332. — Pollich. fl.
pal. n. 842.

Orchis radicibus subrotundis, galea conni-
 vente, labello trifido reflexo. *Hall. stirp.*
helv. n. 1284.

Orchis odore Hirci minor. *Bauh. pin. 82. —*
Mapp. fl. Alf. p. 217.

Icon. Jacqu. fl. Austr. t. 122. — Oeder. fl.
Dan. t. 224. — Hall. l. c. t. 34. fig. media.

Germ. Wantzen - Knabenkraut.

Gall. Orchis punais.

Hab. In pratis apricis praesertim montosis. Re-
 tro *Ettlingen auf dem Dobel*. In Brisgovia
am Kaiserstuhl in pratis montosis prope
Schellingen. In Marggr. superioris pratis
 montanis passim non infrequens. In Alsatia
 prope *Lingelsheim* in pratis, nec non in pra-
 tis montosis retro *Barr* et alibi passim. In
 Palatinatu in pratis siccis inter *Dürkheim*,
Lambsheim, *Frankenthal* et *dem Holzhof*
 post *Pollichium* vidi.

Orchis radicibus subrotundis, galeae petalis lineatis, labello trifido crenato, medio segmento emarginato. *Hall. stirp. helv.* n. 1283.

Orchis Morio femina. *Bauh. pin.* 82. — *Mapp. fl. Alf.* p. 216.

6. Orchis Morio foemina, flore roseo, *Mapp. l. c.*

7. Orchis Morio foemina flore niveo et versicolore. *Mapp. l. c.*

8. Orchis Morio foemina parva. *Mapp. l. c.*

9. Orchis Morio, foemina testiculis ternis, *Mapp. l. c.*

Icon. *Curtis. fl. Lond.* t. 174. — *Oeder. fl. Dan.* t. 253. — *Hall. l. c. t.* 33. — *Vail. parif.* t. 31. f. 13. 14.

Germ. *Weibliches Knabenkraut.*

Call. *Orchis bouffon.*

Hab. Utrinque in pratis ficcioribus apricis passim abunde cum multis variationibus.

Flor. Aprili, Majo. *Semina* Junio, Julio matura. 7

Bulbi subrotundi. *Caulis* digitalis, palmaris altior, erectus. *Folia* alterna, amplexicaulia, superiora erecta, infima reflexa, lanceolata, lineata. *Spica* cylindrica, laxa, pauciflora. *Florae* 6-8-14 vulgo purpurei, interdum rosei, incarnati, albidi, candidi, et versicolores. *Labium* inferius trifidum, crenulatum, maculatum. *Laciniae* laterales retrorsum plicatae, intermedia angustiore, emarginata, sursum plicata, crenulata. *Calcar* adscendens, obtusum, longitudine fere germinis.

Variat. Bulbis rarius tribus, Caule altiore, Floribus in spica longiore majoribus, paucioribus saepe violaceis.

534 GYNANDRIA: DIANDRIA.

Pharm. SALEP Radix.

Radices maxime nutrientes, mucilaginosi, involucentes.

Planta florens mire hinc inde prata ornat.

Flores melliferi.

1359. *ORCHIS mascula* bulbis indivisi, nectarii labio quadrilobo crenulato: cornu obtuso, petalis dorsalibus reflexis. *Linn. Sp. pl. 2. p. 1333. — Poll. fl. pal. n. 844.*

Orchis radicibus subrotundis, petalis lateralibus reflexis, labello trifido, segmento medio longiori bifido. *Hall. stirp. helv. n. 1283.*

Orchis Morio mas foliis sessilibus non maculatis. *Bauh. pin. 215. — Mapp. fl. Alf. p. 215.*

Cynosorchis Morio II. *Tabernæm. p. 1048. Icon.*

β. *Orchis Morio* mas foliis maculatis. *Bauh.*



Affinis praecedenti speciei, a qua distingui-
ar, *Bulbis* majoribus. *Caule* altiore, *Foliis*
stioribus, *Spica* longiore, laxiore, *Floribus*
majoribus, copiosioribus, *Labio* inferiore lon-
giore, *lacinia* media longius ultra lateralia
producta.

ariat. *Foliis* maculatis. *Corollae* purpureo-vio-
laceae, rubrae, roseae, incarnatae, albae, ni-
veae et varicolores.

harm. SALEP. Radix f. Bulbus.

Ex hac specie praesertim Salep venit, quamvis praet-
er hanc, omnes species bulbis majoribus instructae
idem subministrant.

Flores apibus melliferis.

360. ORCHIS *Tabernaemontani*

bulbis indivisis, nectarii labio trilobo
crenulato, lobis lateralibus productioni-
bus, petalis reflexis, foliis lineari en-
siformibus.

Orchis laxiflora, Lamarck et Decand. fl.
franc. 3. p. 247. — Lamarck Encyclop.
4. p. 553. — Flor. franc. 3. p. 504.

Orchis ensifolia radicibus subrotundis, pe-
talis reflexis, nectarii labio bilobo cre-
nato, foliis ensiformibus. Villars. fl. Delph.
2. p. 29. Willd. Linn. Sp. pl. 2. p. 25.

Orchis Morio foemina calcare extuberante,
bifariam diviso. Sequ. pl. Veron. 2. p.
125. t. 15.

Orchis Morio foemina procerior, majori
flore. Vaill. parif. p. 150. t. 31. f. 33. 34.

Orchis angustifolia Bauh. pin. 83.

Testiculus XVI. Tabernaemont. p. 1052.
Icon.

Calcare adscendente, germine brevioris
fo; apice saepe bifido, *Habitu* Scirp-
i foliae, aut Gladioli.

Nota. Nomen dedi a Tabernaemont
plantae, sine dubio in regionibus Rhen-
neationem primus a se lectae, exhibuit

1361. *ORCHIS ustulata* bulbis
nectarii labio quadrifido, punctis
cornu obtuso, petalis distinctis
Sp. pl. 2. p. 1333. — *Pollich.*
n. 845.

Orchis radicibus subrotundis, labe-
drifido; *calcare* brevissimo. *Ha-*
helv. n. 1273.

Orchis militaris pratensis humilior, 1
Alf. p. 215.

Orchis pannonica IV. *Clus. hist.* 2.
Icon. Oeder. fl. Dan. t. 103. — *Hall.*
Germ. Kleines Knabenkraut.

Caulis palmaris et altior, erectus *Spica* à, parva, superne quasi adusta, fusco-nigra, adulta longior, laxior. *Flores* numerosi, i, nitidi. *Labium* inferius quadrifidum, nigra, margine purpureum, punctis purpurascibus, asperis, elevatis, exaratum. *Calcar* e, rubellum.

12. *ORCHIS tephrosanthos* bulbis subrotundis, nectarii labio subquinquefido, angustissimo, petalis confluentibus. *Vill. fl. Delph.* 2. p. 32. — *Desfont. fl. Atl.* 2. p. 319. — *Willd. Linn. Sp. pl.* 4. p. 81.

Orchis simia. *Lamarck et Decand. fl. franc.* 3. p. 249.

Orchis cinerea. *Schrank. Bajerisch. flor.*

Orchis militaris var. 1. *Linn. Sp. pl.* 2. p. 1334.

Orchis flore simiam referens. *Bauh. pin.* 82. — *Mapp. fl. Als.* p. 216.

Orchis zoophora cercopithecum exprimens oreades. *Col. Ecphr.* 1. 320. *Icon.*

2. *Vaill. bot. parif. t.* 31. f. 25. 26.

m. *Affenartiges Knabenkraut. Graues-Knabenkraut.*

l. *Orchis finge. Orchis cendrée.*

. In collibus apricis asperis graminosis passim non infrequens.

r. Majo, Junio. *Semina* Julio matura. 7

Caulis palmaris, pedalis, erectus. *Folia* 2a, parva, ex viridi-cinerea. *Spica* terilis, elliptica, cinerea. *Bractea* breves, 3tae, albidæ. *Labium* inferius quinquefidum, laciniis linearibus, longis; lacinia media

1503. *ORCHIS variegata* Dun
sis, spica brevi, nectarii labio q
lobis apice ferratis, disco glab
lis confluentibus. *Jacqu. Collect*
Hoff. Synops. 485. — *Willd.*
pl. 4. *p.* 21.

Orchis simia. *Villars. fl. Delph.* 2.

Orchis militaris. γ . *Linn.*

Orchis radicalibus subrotundis, spi
sima, labello breviter quadrifid
ferrato, punctato. *Hall. stirp. hel*

Orchis militaris minor. *Mapp. fl.* 2
— *Rupp. fl. Jen. ed. Hall.* 295.

Orchis militaris pratensis elatior,
variegatis. *Mapp. fl. Als. p.* 21.

Cynosorchis militaris pratensis, ela
ribus variegatis. *Bauh. pin.* 81. *f*

Icon. Jacqu. Icon. rar. 3. *t.* 599. —
t. 30.

Germ. Bntes Knabenkraut.

Gall. Orchis panaché.

Hab. Utrinqve passim in collibus app

dia. Spica subrotunda, densa; deflorata elongata. *Petala* quinque erecta, cucullata, conniventia, dilute purpureo-carnea, tria exteriora paulo longiora. *Labium* inferius quadrifidum, petalis concolor, segmentis ad apicem ferrulatis, lateralibus oblongis; intermediis rotundatis, omnibus punctis purpureis vix elevatis adspersis. *Stylar* rectum, subulatum, germine brevius. *Saccae* dilute carnae, acutæ, germine breviores.

Inf. In Horto culta, mutationi non succubuit.

564. *ORCHIS militaris* bulbis indivisis, spica oblonga, nectarii labello quadrifido, lobis integerrimis, disco muricato, petalis confluentibus. *Jacqu. Coll.* 2. p. 268.

Orchis militaris bulbis indivisis, nectarii labio quinquefido, punctis scabro, cornu obtuso, petalis confluentibus. *Linn. Sp. pl.* 2. p. 1333. — *Poll. fl. pal.* n. 446.

Orchis radicibus subrotundis, spica conica, labello quadrifido, brachiolis et pectore perangustis. *Hall. stirp. helv.* n. 1277.

Orchis Rivini bulbis subrotundis, spica conica, petalis conniventibus, nectarii labio trifido, lacinia media productiore bifida. *Gouan. Ill.* p. 74.

Orchis latifolia hiante cucullo, major. *Vaill. bot. parif.* p. 148. t. 31. f. 24. — *Mapp. fl. Als.* p. 214.

Orchis mas latifolia. *Fuchs. hist.* p. 554. *Icon.*

Orchis galea et alis cinereis. *Bauh. hist.* 2. p. 755.

Satyrium trifolium. *Tabernæmont.* p. 1056. *Icon.*

Flor. Majo, Junio, Julio. *Semina* Junio
gusto matura. 7

Differt a præcedenti specie, *Caule* sesqui-
pedali, *Foliis* majoribus, magis viridibus, *Spica* conica,
globose, *Labii* inferioris laciniis omnibus
obtusis, saturatius purpureis, in discis
punctis elevatis, muricatis, purpureis.

Variat. Corollis dilute carneis, et albidis.

Obs. Planta in horto culta, mutationi non

Nota. Ex hac specie, sæpe etiam Radix
officinis pharmaceuticis.

1365. *ORCHIS fusca* bulbis indi-
carii labio quadrifido, laciniis
cornu brevi, petalis confluentibus
Linn. Syst. veg. ed. 14. p. 809.

Orchis militaris. var. B. Linn. Syst.
1334. — *Pollich. fl. pal. n. 84*

Orchis radicibus subrotundis, spicis
labello quadrifido, brachiolis

n. Jacqu. *fl. Austr.* t. 307. — *Hall.* l. c. t. 31.

r m. *Braunes großes Knabenkraut.*

ll. *Orchis brun.*

b. In collibus graminosis dumetosis præsertim calcareo-argillaceis. Prope Grözingen, Berghausen, Weingarten, Bruchsal et alibi passim non infrequens. In Brisgovia am Kaiserstuhl, ad sylvarum marginalia graminea, dumetosa.

r. Majo, Junio. *Semina* Julio matura. *?*

Bulbi gemini, oblongi, majusculi. *Caulis* vitalis, tripedalis, erectus, superne aphyllus, se fusco purpurascens. *Folia* inferiora, longa, lata, late viridia, obtusiuscula, semipedalis-4 uncias lata. *Spica* speciosa, magna, lata vel cylindrica. *Bracteae* pusillae. *Petala* inque conniventia, fusco-viridia, sæpe exaricante fusca. *Labium* inferius quadrifidum inius lateralibus angustioribus, interioribus, obtusis, crenulatis, cum denticulo medio acuminato, albidum, niveum, margine fusco, punctis villosis-penicilliformibus, asperis, elevatis, purpurascentibus adpersum. *Cal.* breve, fuscum, obtusum.

c. In horto botanico culta, mutationi non succumbit. Verum omnino hortorum decus.

66. *ORCHIS moravica* bulbis indivisis, nectarii labio trifido, lacinia media subrotunda emarginata, amplissima, petalis conniventibus. *Jacqu. Collect.* 1. p. 61. — *Murr. Linn. Syst. veg.* p. 809.

n. Jacqu. *Icon. rar.* 1. t. 182.

r m. *Großes Mährisches Knabenkraut.*

Gall. Orchis de Moravie.

Hab. Cum præcedente specie passim, præ-
tim retro *Grötzingen* et *Weingarten*.

Flor. Majo, Junio. *Semina* Julio matura. 7

Affinis præcedenti, a qua differt, *Fol*
paucioribus, *Petalis* pallide virentibus, *pr*
pureo striatis, *Labio* inferiore trifido, pallid
rubello seu carneo, *laciniis* lateralibus lineis
bus obtusis; *intermedia* maxima, ampla, sul
rotunda, emarginata, crenulata.

Obs. Planta in horto botanico culta, speciosa, mot
tioni non succubuit.

Nota. Haec species sine dubio passim pro *O. pa-*
pilionacea Linn. habitæ est, quam nescio
an vix Germania ferat spontaneam.

1367. *ORCHIS palustris* bulbis indivisi-
nectarii labio inverse triangulari, apice
obiter trilobo, lobis rotundatis, medio
emarginato, cornu obtuso, petalis obtrun-
catis exterioribus patulis. *Jacqu. Collect.*
1. p. 75.

Orchis mascula *Granta* 9. *Andria* 5.

inantibus, lineari-lanceolatis, canaliculatis, rectis, *Spica* brevior 3-7 flora, *Labii* amplius laciniis lateralibus rotundatis, obtusis, *media* multo minore, bifida, *Calcare* gerere subbreviore, quandoque sublongiore, obtusum. *Flores* inodori, saturate purpurei, aut albi.

Nota. Planta omnino ab *O. mascula* et *O. Morione* specie diversa.

¶ In horto culta, mutationi non succubuit.

Nota. Orchidem pallentem bulbis indivisis, nectarium labio trifido integerrimo, cornu obtuso mediocri, petalis patentibus. *Linn. Mant.* 292. *Jacqu. fl. Austr.* 1. t. 45. — *Hall. stirp. helv.* n. 1281. t. 30. (sed *Orchis foetida* sylvatica proecox flore albo barba luteola *Rupp. fl. Jen.* p. 297) frustra hucusque in terris Badensibus et confinibus quaesivi.

†† *Bulbis palmatis.*

68. *ORCHIS latifolia* bulbis subpalmatis rectis, nectarium cornu conico: labio trilobo lateribus reflexo, bracteis flore longioribus. *Linn. Sp. pl.* 2. p. 1334. — *Poll. fl. pal.* n. 847.

Orchis radicibus palmatis, caule fistuloso, bracteis maximis, labello trifido, serrato, medio segmento obtuso. *Hall. stirp. helv.* n. 1279.

Orchis palmata pratensis latifolia, longis calcaribus. *Bauh. pin.* 85. — *Mapp. fl. Als.* p. 220.

Orchis palmata palustris latifolia. *Bauh. pin.* 86. *Mapp. l. c.*

Orchis palmata palustris altera. *Bauh. pin.* 86. — *Mapp. l. c.* 223.

GYNANDRIA. DIANDRIA.

4. *Orchis palmata palustris maculata*. Bauh. pin. 86. — Mapp. fl. Alf. p. 223.

5. *Orchis palmata palustris; tota rubra*. Bauh. pin. 86. — Mapp. p. 223.

Icon. Curtis. fl. Lond. t. 250 — Hall. l. t. 32. — Oeder. fl. Dan. t. 266. — Black. t. 405.

Germ. Breitblättrichtes Händleinkraut.

Gall. *Orchis à feuilles larges*.

Hab. Utrunque in pratis riguis, humidis, virginosis, locisque graminosis sylvaticis passim abunde cum multis variationibus.

Flor. Majo, Junio. Semine Julio matura.

Planta pro diverso loco natali magnitudine et colore ludit, facile distinguenda. Spica longe bracteata. Bracteis flore semper longioribus.

Variat. Foliis latioribus et angustioribus, maculatis et immaculatis. Floribus carnis, rubellis, albis, purpureis, violaceis, pallide roseis, albis variegatis.

1369. *ORCHIS maculata* bulbis palmatis, patentibus, nectarii cornu germinebus brevioribus: labio plano, petalis dorsilibus patulis. Linn. Sp. pl. 2. p. 1335. Poll. fl. pal. n. 849.

Orchis radicibus palmatis, caule solido, labello trifido ferrato, medio segmento acuminato. Hall. stirp. helv. n. 1278.

Orchis palmata pratensis maculata. Bauh. pin. 85. — Mapp. fl. Alf. p. 221.

β. *Orchis palmata pratensis maculata flore alba*. Bauh. pin. 85. — Mapp. fl. Alf. p. 221. γ. Flores

7. Flore variegato. *Bauh. pin.* 85. — *Mapp. l. c.*

4. *Orchis palmata maculata*, flore albido, rubellis punctis et striis leviter notato. *Mapp. fl. Alf. p.* 221. (Caulis et spica insignis magnitudinis.)

Icon. Oeder. fl. Dan. t. 933. — *Hall. l. c. t.* 32

erm. Geflecktes Händleinkraut.

all. Orchis tachetée.

Tab. Utrunque in pratis sylvaticis, humidis et siccioribus, locisque graminosis ad sylvarum marginalia passim frequens.

Flor. Majo, Junio. *Semina* Julio mature. 7

Distinguitur a praecedente, *Caule* solidiori, subangulato, saepe altiori, *Foliis* subangustioribus, *Bracteis* minoribus, *Floribus* pallidioribus, *Labio* inferiore plano, profundius trifido, *Laciniis* lateralibus non reflexis, *Calcare* magiore.

Folia saepissime pulcherrime maculata.

370. *ORCHIS odoratissima* bulbis palmatis nectarium cornu recurvo brevior, labio trilobo, foliis linearibus. *Linn. Sp. pl. 2. p.* 1336.

Orchis radicibus palmatis, flore concolore, labello obtuse trifido, calcare germine brevior. *Hall. stirp. helv. n.* 1274.

Orchis palmata, angustifolia minor odoratissima. *Bauh. pin.* 86. — *Prodr. p.* 30. *Icon.* — *Mapp. fl. Alf. p.* 221.

Icon. Jacqu. fl. Austr. t. 264. — *Hall. l. c. t.* 29

erm. Geruchvolles Händleinkraut.

all. Orchis odorant.

Hab. In Margraviatus superioris collibus prope Kembs rarius t. *Vulpio*, nec non in monte prope Crenzach. In ditione Basileensi in pascuis montis Dietisberg, Michelsfeldae. t. C. Bauhino; in monte Wasserfallen t. Strehelino. In Alsatia prope Eschau in pratis et Hangenbietenheim et auf dem Pafsberg prope Buchsweiler t. Mappo.

Flor. Majo, Junio. *Semina* Julio matura. 7

Affinis sequenti, a qua differt, *Caule* foliisque durioribus, *Labio* inferiore longiore, angustiore; *lacinia* media magis producta; *Calcare* duplo brevior, *Odore* forti, singulari, grato.

Nota. Citat Mappus plantam circa Argentoratum in der Ruprechtsau et in dem Illkircher Wört, ubi hodie vix crescit.

Ad rariores omnino Florae cives pertinet.

1371. *ORCHIS Conopsea* bulbis palmatis, nectarii cornu setaceo germinibus longiore: labio trifido; petalis duobus

GYNANDRIA. DIANDRIA. 247

m. Stechfliegenartiges Händleinkraut.

l. Orchis à long éperon.

. Utrunque in pratis colliculosis, montosis
siccioribus apricis, sylvis ericetosis grami-
neis passim frequens.

*r. Majo, Junio, Julio. Semina Julio, Au-
gusto matura. 7*

*Radix palmata. Caulis pedalis, sesquipeda-
litalis et altior, erectus. Folia lanceo-
Spica 2-3-4 uncialis et longior, multiflo-
rubragracilis. Bracteae germine sublongiores,
ratae. Corollae dilute purpureae, carnea-
um inferius trifidum: laciniis lateralibus
ulatis, intermedia ovata, integra. Calcar-
en longe superans, praelongum, tenue,
osum.*

ia t. Corollis albis.

*et a. Omnes orchidum bulbi indivisi seu testiculati,
et palmati, vires habent nutrientes, inspissantes,
hinc promiscue saepe Radicem Salep officinis
pharmaceuticis subministrant.*

2. ORCHIS sambucina bulbis sub-
palmatis rectis, nectarij cornu conico,
labio ovato subtrilobo, bracteis longitu-
dine florum. *Linn. Sp. pl. 2. p. 1334. —
Pollich. fl. pal. n. 848.*

*Orchis radicibus palmatis, bracteis colo-
ratis, labello circumferrato trilobato, lo-
bo medio emarginato. Hall. stirp. helv.
n. 1280.*

*Orchis palmata lutea; labio floris maculato
Sequ. pl. Veron. 349. t. 8. f. 5.*

*Orchis palmata sambuci odore. Bauh. pin. 86.
n. Jacqu. fl. Austr. t. 108.*

M m 2

Germ. Hollunderduftendes Händleinkraut.
bes Händleinkraut.

Gall. *Orchis fureau.*

Hab. In palatinatu transrheno in montosis
sylvaticis circa *Steinbach* versus dem
nersberg t. *Pollichio*. Ni fallor vidi in
tos in Principatu *Fürstenbergenfi*.

Flor. Majo, Junio. *Semina* Julio matur

Radicis unus bulbus palmatus, alter
tus. *Caulis* digitalis, palmaris, erectus.
lia ovali - lanceolata, laete viridia. *Sp*
ovata, pauciflora. *Stipulae* floribus longior
lanceolatae, erectae. *Corollae* pallide flav
Labium inferius trifidum, *lacinia* media en
ginata, crenulata. *Calcar* germine longius,
res, obtusum.

Odor florum Sambuci.

Ad rariores florum cives pertinet.

CCCCVIII. SATYRIUM.

Linn. Gen. pl. ed. Schreb. n. 1368.

Spathae vagae. *Spadix* simplex. *Periant*
um nullum. *Corolla* petala quinque, ovato-
longa: tria exteriora, duo interiora sursum co
niventia in galeam. *Nectarium* monophyll
a latere inferiore inter divisuram petalorum
septaculo adnexum. *Labio superiore* erec
brevissimo; *inferiore* plano, pendulo, basi bu
scrotiformi postice prominente. *Capsula* obli
ga, unilocularis, tricarinata, trivalvis, sub
rinis trifariam dehiscens, apice et basi coh
rens. *Semina* numerosa, minima, scobiform

Nota. *Caulis* f. *Spapus* simplex. *Fol*
integra.

373. SATYRIUM *hircinum* bulbis indivisis, foliis lanceolatis, nectarii labio trifido: intermedia lineari elongata, obliqua praemorsa. *Linn. Sp. pl. 2. p. 1337. Pollich. fl. pal. n. 851.*

Orchis hircina Swartz. *Act. Holm. 1800. p. 207. — Willd. Linn. Sp. pl. 4. p. 28. — Scopol. fl. Carn. 2. n. 1113.*

Orchis radicibus subrotundis, labello longissimo plicato tripartito. Hall. stirp. helv. n. 1268.

Orchis barbata odore Hirci brevior latioreque folio. *Bauh. pin. 82. — Mapp. fl. Alf. p. 217.*

Testiculus hircinus I. Tabernaemont. p. 1055. Icon.

B. *Orchis* odore Hirci, longior et angustior folio. *Bauh. pin. 82. — Mapp. fl. Alf. p. 217.*

Testiculus hircinus II. Tabernaemont. p. 1055. Icon.

non. Jacqu. fl. Austr. t. 367. — Hall. l. c. t. 25. — Vaill. bot. par. t. 30. f. 6.

erm. Bocksgeilen. Grofse Bocksgeilen.

all. Satyrion bouquin.

a b. In collibus argillaceo calcareis dumetosis gramineis aridis retro Grözingen, prope Weingarten et Bruchsal infrequens. In Marggraviatu superiore prope Sulzburg et Lauffen; in montolis inter Kembs et Itstein; am Rötteler Schloß frequens, et infra Esringen. In Alsatia prope Bischofsheim; inter Buchweiler et Zabern, in monte Passberg copiose, in montolis prope Rappolsweiler t. Mappo, In Palatinatu in collibus

juniperinis siccis dumetosis circa *Dürkheim*, *Umstein*, *Calstatt* et *Herzheim* post *Pölichium* vidi. In *Sponheimia* prope *Weyler* et *Martinstein*.

Flor. Majo, Junio. *Semina* Julio matura. 7

Bulbi gemelli ex rotundo-ovati, magni. *Caulis* sesquipedalis et longior, erectus, crassifuscus. *Folia* radicalia, caulinaque inferiora ampla, late lanceolata. *Spica* speciosa, multiflora, adulta saepe curvata. *Galea* petalis conniventibus, quinque ovatis obtusis, externe pallide albidis, interne striis purpureis viridibusque pictis, constructa. *Labium* inferius trifidum: *lacinia* intermedia longissima, biunciali, triunciali et longiori, lineari, angusta, obtusa, ex albido-viridi, anterieus maculis rubiginosis aut purpureis, picta; lateralibus brevioribus, undulato-crispatis. *Germen* virens, viscidum,

Flores fortem hircosum quidem odorem spirant.

1374. *SATYRIUM viride* bulbis palmatis, foliis oblongis, obtusis: nectarii labio linearitrisido: intermedia obsoleta. *Linn. Sp. pl.* 2. p. 1337. — *Poll. fl. pal.* n. 832.

Orchis viridis. *Swartz. Act. Holm.* 1800. p. 207. — *Willd. Linn. Sp. pl.* 4. p. 33. — *Scopol. fl. Carn.* n. 1122.

Orchis radicibus palmatis, galea connivente, labello trifulco, calcare brevissimo. *Hall. stirp. helv.* n. 1269.

Orchis palmata flore viridi. *Bauh. pin.* 86. *Prodr.* 30. *Mapp. fl. Als.* p. 222.

Icon. *Oeder. fl. Dan.* t. 77.

Germ. Grüne Bocksgeilen.

Gall. *Satyrium verdatre.*

ab. In ditione Basileensi in pascuis montis *Vogelberg* et *Ramstein*. In Principatu *Fürstenbergensi* in collibus juniperinis prope *Donaueschingen* cum *Carlina acauli* et *Osmunda Lunaria* nupervidi. In Alsatia circa *Argentinam* prope *Niederhausbergen* t. *Mappo*. In Palatinatu in pratis siccioribus inter *Dürkheim*, *Lambsheim*, *Frankenthal* et *dem Holzhof*; circa *Mannheim*; in pratis circa *Cusel*. Circa *Alzey*, *Germersheim*, *Neustadt etc.* t. *Pollichio*. In Sponhemiae pratis passim, frequens prope *Kirchberg* cum *Osmunda Lunaria*.

lor. Majo, Junio. *Semina* Julio matura. 7.

Bulbi palmati. *Caulis* palmaris, pedalis, erectus. *Folia* ovali-lanceolata, obtusa, infimae viora. *Spica* uncialis, biuncialis, laxa. *Bractae* lanceolatae, germine longiores. *Corollae* pallido luteae virescentes. *Labium* inferius fidum: *Laciniis* lateralibus acutis, intermedia evissima.

variat. Floribus submajoribus, Bracteis longioribus.

575. *SATYRIUM nigrum* bulbis palmatis, foliis linearibus, floribus resupinatis, nectarium labio indiviso ovato acuminato. *Linn. Sp. pl.* 2. p. 1338.

Orchis nigra Swartz. Act. holm. 1800. p. 207. *Willd. Linn. Sp. pl.* 4. p. 35. — *Scopol. fl. Carn. n.* 1123.

Orchis radicibus palmatis, spica densissima, flore resupinato, calcare brevissimo. *Hall. stirp. helv. n.* 1271.

Orchis palmata angustifolia alpina, nigro flore. *Bauh. pin.* 86. — *Mapp. fl. Alf. p.* 222.

Icon. Jacqu. fl. Austr. t. 363. — Hall. l. 6 t. 27.

Germ. Schwarze Bocksgeilen.

Gall. Satyrion noir.

Hab. In Alfatiae jugo Vogeli auf dem Hochgebürg retro dem schwarzen See in summo cacumine t. Mappo.

Flor. Junio, Julio. *Semina* Augusto matura. 7

Bulbi palmati. *Caulis* digitalis, palmaris, vix altior, erectus. *Folia* linearia, canaliculata, superiora acuta. *Spica* ovata vel subrotunda, parva, densissima. *Bracteae* longitudine florum, fusco-purpureae. *Flores* resupinati, fusco-purpurei, læpissime expurpureo-nigricantes, odorati.

Nota. Planta alias alpina ad rariores florae cives pertinet.

1376. SATYRIUM *albidum* bulbis fasciculatis, foliis lanceolatis, nectarii labio trifido acuto: lacinia intermedia obtusa. *Linn. Sp. pl. 2. p. 1338.*

Orchis albida. *Swartz. Act. Holm. 1800. p. 207. Willd. Linn. Sp. pl. 4. p. 38.*

Orchis radicibus confertis teretibus, calcare brevissimo, labello trifido. *Hall. stirp. helv. n. 1270.*

Pseudo Orchis alpina, flore herbaceo. *Michel. Gen. 30. t. 26.*

Limodorum montanum, flore albo dilute virescente. *Chom. Act. paris. 1705. p. 517.*

Icon. *Oeder. fl. Dan. t. 115.*

Germ. Weisse Bocksgeilen.

Gall. Satyrion à fleurs blanches.

6. Prope *Frauenalb*, *Herrnalb*; auf dem *Tobel* in pratis montosis ad sylvarum marginalia, et in locis gramineis ericetosis cum *Lycopodio complanato*, non infrequens. In *Marggraviatu* superiore in monte *Feldberg* et *Belchen* non infrequens. In ditione *Basileensi* in monte *Gelderkinden* et *Farnsburg*.

r. Majo, Junio, Julio. *Semina* Julio, Augusto, matura. 7

Bulbi fasciculati, teretes. *Caulis* palmaris, lis, erectus. *Folia* lanceolata. *Spica* juniora, demum laxa, pollicaris, tripollicaris. es ex viridescente albidi, deflexi. *Labium* trifidum: *laciniis* brevibus, acuminatis: a majore, obtusiore.

7. SATYRIUM *Epipogium* bulbis compressis dentatis, caule vaginato, corollis resupinatis, nectarii labio indiviso. *Linn. Sp. pl.* 2. p. 1338.

imodorum *Epipogium* aphyllum, scapo vaginato paucifloro, floribus pendulis resupinatis, labello trilobo concavo, cornu adscendente ovato. *Swartz Nov. Act. Upsl.* 6. p. 80. — *Willd. Linn. Sp. pl.* 4. p. 129.

pipactis caule aphylo, flore supinato, labello ovato-lanceolato, calcare ovato turgido. *Hall. stirp. helv. n.* 1289.

Epipogium. *Gmel. fl. Sibir.* 1. p. 12. t. 2. f. 2.

1. *Jacqu. fl. Austr.* t. 84.

n. *Saftige Bocksgeilen.*

1. *Satyrium epipoga.*

Hab. In sylvaticis fagineis umbrosis, in te
ex putrefactis foliis orta prope *dem Stein*
cher Baad versus templum *Sanctae B*
barae dicatum rarius vidi 1798. et insequ
tibus annis. In ditione Basileensi supra *Re*
stein in sylvis umbrosis. *t. Lachenalio.*

Flor. Julio, Augusto. *Semina* Augusto
tura. ¶

Radix coralloidea, compressa, carnosa, l
co-albida. *Scapus* spithamaeus, erectus, infer
subbulbosus, teres, simplex, aphyllus sordi
ex flavescente albidus diaphanus, squamis me
branaceis, vaginantibus, ovatis truncatis retic
latis luteo-fuscis per intervalla praeditus. *Spit*
fubracemosa, simplex, laxa, pauciflora, saep
2-3-4 flora. *Flores* mediocres, penduli, su
secundi, inversi, luride ochroleuci, labio
facco scrotiformi crasso obtuso, sursum directi
Petala quinque, patentia, sordide flavescents
punctis longitudinalibus purpureis margine
asperata, duo exteriora lineari lanceolata; du
interiora latiora et sublongiora, quintum inte
medium sublineare brevius. *Labium* inferi
ovatum acutum, integrum, appendiculatum
lineis purpureis duabus penicilliformibus, pilis
ad latera pictum.

Habitus totius plantae singularis. In Hortum bot
nicum omni studio repetitis vicibus transplantata, pen
sine dubio parasitica.

Nota. Ad rariores florae cives pertinet.

1378. *SATYRIUM repens* bulbis fibrosis
foliis ovatis radicalibus, floribus secat
dis. *Linn. Sp. pl. 2. p. 1339.*

Neottia repens foliis radicalibus ovatis pen
latis, reticulatis, scapo vaginato, floribus

GYNANDRIA. DIANDRIA. 555

bus pubescentibus, secundis, labello petalisque lanceolatis. *Willd. Linn. Sp. pl.* 2. p. 75. — *Swartz. Act. Holm.* 1800. p. 226.

Epipactis foliis petiolatis ovato-lanceolatis, floribus tetrapetalis hirsutis. *Hall. stirp. helv. n.* 1295.

Orchis minor flosculis albis f. radice repente. *Camerar. Epit. III, t.* 35.

n. Jacqu. fl. Austr. t. 369. — *Lightf. fl. Scot. 2. t.* 22. — *Oeder. fl. Dan. t.* 812. — *Schkuhr. bot. handb. t.* 272.

m. Kriechende Bockageilen.

l. Satyrion rampante.

b. In Margraviatu superiore in pineto supra *Kembs* in solo mulcoso copiose *t. Vulpio.* In ditione Basileensi in monte *Vogelberg, Wasserfallen*, inter *Alschwyler* et *Neubad*, prope *dem Wendenhof*, circa *Arisdorf* in pinetis. In Principatu superiore in pinetis passim frequens, inter *Mörsburg* et *Ueberlingen* nuper copiose vidi.

r. Majo, Junio. *Semina* Julio matura. 7

Radix repens articulata, bulbosa, bulbi teres, carnosus, villosus. *Caulis* palmaris, spithameus, erectus, teres, inferne foliosus, glaber, tunc villosus, aphyllus, squamis foliaceis antheris, erectis, praeditus. *Folia* radicalia caulique inferiora petiolata, ovata, subtus quinque-*ervia*. *Spica* secunda, biuncialis et longior, glaberrima. *Bractee* lanceolatae, germine brevioribus. *Flores* parvi, albidi. *Petala* quinque connatis, villosa. *Villi* apice glanduliferi.

t. a. Partes plantae fructificationis egregie exhibit acutissimus *Schkuhr. l. c.* et 3. p. 201,

386 GYNANDRIA: SCANDIA.

CCCCIX. OPHRYS.

Linn. Gen. plant. ed. Schreb. n. 1369.

Spathae vagae. *Spadix* simplex. *Perianthium* nullum. *Corolla* petala quinque oblonga, sursum conniventia, aequalia: duo horum exteriora. *Nectarium* petalis longius, dependens, postice tantum subcarinatum. *Capsula* subovata, trigona, obtusa, firiata, trivalvis, unilocularis, angulis carinatis debiscens. *Semina* numerosa, scobiformia.

† *Bulbis ramosis.*

1379. OPHRYS *Nidus avis* bulbis fi-

broso-fasciculatis, caule vaginato, nectarii labio bifido. Linn. Sp. pl. 2. p. 1339.
— Poll. fl. pal. n. 853.

Epipactis Nidus avis. Swartz. Act. Holm. 1800. p. 232. Willd. Linn. Sp. pl. 4. p. 87.

Epipactis aphylla, flore inermi, labello bifido. Hall. stirp. helv. n. 1290.

Nidus avis. Mapp. fl. Alf. p. 210.

Orobanche affinis Nidus avis. Bauh. hist. p. 782.

Icon. Oeder. fl. Dan. t. t. 181.

Germ. Vogelneft.

Gall. Ophris Nid-d'oiseaux.

Hab. Utrunque passim in nemoribus umbrosis ad arborum radices, parasitica.

Flor. Majo, Junio. Semina Julio matura.

Radix lateralis, dense fibroso-fasciculata. Nidum avis imitans. Caulis palmaris, pedalis.

altior, erectus, crassiusculus, squamis vagitus luridus, ex luteo-fuscus, demum fusco-panicatus. *Folia* nulla. *Spica* longa, inferne raris, superne densa, multiflora, cylindrica. *Stamina* fere omnia, sordide ex luteo-fusci. *Labium* bifidum.

380. *OPHRYS Corallorhiza* bulbis

ramosis flexuosis, caule vaginato, nectario trifido. *Linn. Sp. pl.* 2. p. 1339.

Ophrys corallorhiza bulbis ramosis flexuoso-divaricatis, caule vaginato aphyllis, nectarii labio indiviso. *Smith. fl. brit.* 3. p. 932.

Cymbidium Corallorhizon. *Swartz. Act. Holm.* 1800. p. 238. — *Willd. Linn. Sp. pl.* 4. p. 109.

Corallorhiza. *Hall. stirp. helv. n.* 1301. — *Gmel. Sibir.* 1. p. 26.

Neottia bulbis reticulatis, nectario labio trifido. *Act. Upsal.* 1740. p. 34. *Linn. fl. Succ.* 743.

Orobanche radice coralloide. *Bauh. pin.* 88. *non. Oeder. fl. Dan. t.* 451. — *Hall. l. c. t.* 44. *Lightf. fl. Scot.* 1. t. 23.

erm. Corallenwurz.

211. *Ophris coralloide.*

212. In ditione Basileensi ad radices abietum in locis umbrosis supra *Ramstein. t. Lachenalia*, perraro.

213. Majo, Junio. *Semina* Julio matura. 7

Radix lateralis, ramosa, corallina, appendiculata, succiplena, alba. *Caulis* spithameus, erectus, squamis vaginatus, aphyllus. *Spica* densa, laxa, 9-13 flori. *Bracteae* ovato-aeolatae, minutae. *Flores* breviter pedun-

culati, albid. *Labium* oblongum integrifolium. *Fructus* penduli.

Nota. Ad rarissimos florae cives pertinet.

1381. OPHRYS *spiralis* bulbis aggregatis oblongis, caule subsolioso, floribus secundis, nectarii labio indiviso crenato. *Linn. Sp. pl.* 2. p. 1340. — *Poll. fl. pal.* n. 855

Neottia spiralis foliis radicalibus oblongis subpetiolatis, spica tortiti, floribus secundis, labio ovato. *Swartz. fl. Ind. occid.* 3. p. 1408. — *Willd. Linn. Sp. pl.* 4. p. 73.

Neottia bulbis fasciculatis, nectarii labio indiviso. *Act. Ups.* 1740. p. 32.

Epipactis bulbis cylindricis, spica spirali, labello crenulato. *Hall. stirp. helv.* n. 1294.

Orchis spiralis alba odorata. *Bauh. hist.* 2. p. 769. — *Mapp. fl. Als.* p. 217.

Icon. Curtis. fl. Lond. t. 270. — *Oeder. fl. Dan. t.* 387. — *Hall. l. c. t.* 38.

Germ. Spiralförmige Ophrys.

Gall. Ophris en spirale.

Hab. Utrunque passim in pascuis gramineis. Circa *Beuertheim*, *Rintheim* et *Grötzingen*. In *Brisgovia am Kaiserstuhl* passim. In *Margraviatu superiore* prope *Krentzsch* et alibi. In *Alsatia* circa *Argentinam* in der *Gansau* et alibi. In *Palatinatu*, in montibus *Donnersberg* pascuis graminosis t. *Pollich.*

Flor. Julio, Augusto. **Semina** Septembri maturantur. ♀

Radix bulbosa. *Bulbi* 1-2-3-4 oblongi, minati, inaequales. *Caulis* palmaris, spithaeus, erectus, squamis foliaceis, pubescenti, vaginatus. *Folia* radicalia ad terram exarsa, ovata, acuta. *Spica* gracilis, spiralis. *Fructus* secundi, parvi, albidi, odorati. *Petala* quinque, subaequalia, villosiuscula. *Labium* oblongum, crenulatum, concavum.

82. *OPHRYS ovata* bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido. *Linn. Sp. pl.* 2. p. 1340. — *Poll. fl. pal.* n. 855.

Epipactis ovata. *Swartz. Act. Holm.* 1800. p. 232. — *Willd. Linn. Sp. pl.* 4. p. 87.

Epipactis foliis binis ovatis, labello bifido. *Hall. stirp. helv.* n. 1291.

Ophris bifolia. *Bauh. pin.* 87. — *Mapp. fl. Als.* p. 213: — *Tabernaemont.* p. 1101. *Icon.*

Ophris trifolia. *Bauh. pin.* 87. — *Mapp. l. c.* — *Tabernaemont.* p. 1101. *Icon.*

o n. *Curtis fl. Lond.* t. 177. — *Oeder. fl. Dan.* t. 137. — *Hall. l. c. t.* 37. — *Schkuhr. bot. Hantb.* t. 273.

r m. *Zweyblättrige Ophris.*

l l. *Ophris double feville.*

r b. Utrinque in pratis praesertim montosis sylvaticis et sylvis graminosis passim frequens.

r r. Majo, Junio. *Semina* Julio matura. 3

Radix fibrosa, fibris copiosis cylindricis, tortis. *Caulis* pedalis, sesquipedalis, bipedalis, et altior, erectus, sub medio diphyllus.

GYNANDRIA. DIANDRIA.

Folia duo opposita, sessilia, ovalia, quinque-
nervia, lata. *Spica* longa, laxa, angusta. *Flo-*
res ex luteo-viriduli. *Labium* inferius profunde
bifidum, descendens.

Variet foliis ternis.

1383. OPHRYS cordata bulbo fibroso,
caule bifolio, foliis cordatis. *Linn. Sp.*
pl. 2. p. 1340.

Epipactis cordata. Swartz. Act. Holm. 1800
p. 232. — *Willd. Linn. Sp. pl.* 4. p. 18.

Epipactis foliis binis cordatis, labello bifido
postice bidentato. *Hall. Stirp. helv.* 2.
1392.

Ophris minima. Bauh. pin. 87. *Prodr.* 2.
Icon. Hall. l. c. t. 22. f. 4.

Germ. Kleine zweyblättrige Ophris.

Gall. Ophris petite à double feville.

Hab. In montosis umbrosis auf dem Kall-
brunn. In Marggraviatu superiore in mon-
tis Belchen umbrosis rarius t. *Vulpio.*

Flor. Junio, Julio: *Semina* Augusto matura.

Radix fibrosa, fibris cylindricis, carnosae.
Caulis biuncialis, digitalis semipedalis et altior,
erectus, delicatulus, bifolius, caeterum nudus.
Folia opposita, cordata, laete viridia, respectu
plantulae magna. *Spica* parva, laxa, 5-7 florum.
Flores parvi, patuli, subvirescentes, demum
fusco rubelli. *Labium* inferius profunde bifidum,
postice utrinque dentatum: lacinulis acuminatis.

Nota. Planta pulchella ad rariores omnino flores
cives pertinet.

†† *Bulb.*

GYNANDRIA. DIANDRIA. 561

†† *Bulbis rotundis.*

14. *OPHRIS Loeseli* bulbo subrotundo: scapo nudo trigono, nectarii labello ovato. *Linn. Sp. pl.* 2. p. 1341. — *Roth. fl. germ.* 2. p. 402.

Malaxis Loeseli foliis binis ovato-lanceolatis, scapo trigono, labello apice ovato recurvato. *Swartz. Act. Holm.* 1806. p. 235. — *Willd. Linn. Sp. pl.* 4. p. 92.

Ophris liliifolia. β. *Loeseli.* *Hudf. fl. Angl.* 2. p. 390.

Ophris diphyll bulbosa. *Loesel. fl. pruss.* 180. t. 58.

Orchis liliifolius minor fabuletorum Zelandiae et Bataviae. *Bauh. hist.* 2. p. 770.

seudo-*Orchis bifolia* palustris. *Raj. Synops.* 382.

15. *Oeder. fl. Dan.* t. 877. sub *Ophride paludosa.*

m. *Loesels Ophris.*

l. *Ophris de Mr. Loesel.*

. In Alsatia prope *Hagenau* ad stagnum in fabulosis udis, ubi legi An. 1779.

r. Junio, Julio. *Semina* Augusto matura. 7

Differt a sequenti specie affini, *Scapo* non quegono sed exacte trigono, *Foliis* radicalibus, lanceolatis, vaginatis, longioribus, que glabris, longitudine fere scapi, *Spica* racemosa laxa, 4-6 flora. *Floribus* majoribus luteo-albidis. *Labium* inferius ovatum, recurvatum.

16. a. Ad rariores Florae cives pertinet. Foran et in fabulosis udis ad Lacum Bodamicum crescit.

17. a. Bod. Tom. III,

N 2

1385. *OPHRIS paludosa* bulbo tundo, scapo subnudo pentagoniorum apicibus scabris, nectarii integro. *Linn. Sp. pl. 2. p. 1341.* — *fl. pal. n. 855.* — *Act. Palat. II.*

Malaxis paludosa foliis subquaternis scabris, scapo pentagono; labello cavo acuto. *Swartz. Act. Holm. 1235.* — *Willd. Linn. Sp. pl. 4. p.*

Icon. Oeder. fl. Dan. t. 1234.

Germ. Sumpf-Ophris.

Gall. Ophris des marais.

Hab. In Palatinatu in praefectura *Lau-*
gr. in pratis paludosis inter der *Effel-*
Gersweiler, Ottersberg et der Be-
mühl. t. Pollichio.

Flor. Julio, Augusto. *Semina* Augusto
tembri matura. ¶

Radix bulbosa. *Bulbus* ovatus unicus.
lis digitalis, palmaris, erectus, pentag-
nusus, versus radicem foliosus. *Folia* r-
lia, caulinaque infima plura, ovali-lance-
obtusula, apice tuberculis scabra, cauli
breviora. *Spica* subracemosa, 9-11 flora
cillis. *Flores* ex luteo-viridescentes. *Petala*
que, tria exteriora linearia, reflexa, du-
riora angustiora. *Labium* inferius laceolati-
tum, integrum, reflexum.

Nota. Ad rarissimos Florae cives pertinet,
adhuc in udis ad Lacum Bodamicum cresc

1386. *OPHRIS Monorchis* bulbo
boso, scapo nudo, nectarii labio t
cruciato. *Linn. Sp. pl. 2. p. 1342.* —

fl. pal. n. 857. — Swartz. Act. Holm. 1800. p. 222. — Willd. Linn. Sp. pl. 4. p. 61.

Orchis bulbo unico subrotundo, labello cruciformi. Hall. stirp. helv. n. 1262.

Monorchis flore calcari destituto ac radice unico tantum praedita bulbo. Mapp. fl. Alf. p. 220.

Orchis odorata moschata f. Monorchis. Bduh. pin. 84.

Orchis lutea hirsuto folio. Bauh. pin. 84. Testiculus luteus; Orchis lutea l. Tabernaemont. p. 1054. Icon.

Triorchis lutea folio glabro. Bauh. pin. 84.

Triorchis lutea l. II. III. Tabernaemont. p. 1058. Icon.

n. Oeder. fl. Dan. t. 102. — Hall. l. c. t. 22.

r. m. Einhodige Ophris. Wohlriechende Ophris.

l. Ophris unibulbe.

Utrinque in collibus calcareo-argillaceis graminosis humidis passim frequens. In monte Thurnberg prope Durlach frequens, nec non prope Grötzingen et Berghausen. In Brisgovia am Kaiserstuhl prope Rothweil ubi cum Hebelia collina crescentem abunde nuper vidi.

r. Junio, Julio. Semina Augusto matura.

Bulbus unicus subrotundus. Caulis palmapedalis, erectus, teres, subnudus, gracilis. radicalia 2-3 ovato lanceolata. Spica longigracilis, multiflora. Flores subobliqui, evirentes, odoratissimi, mellei, suaveolens. Labium inferius trifidum, cruciforme.

564 GYNANDRIA. DIANDRIA.

Variat. Foliis glabris et villosis. Bulbis tribus.
Flores gratissimum odorem melleum spirant.

1387. *OPHRIS anthropophora* bulbis subrotundis, scapo folioso, nectario labio lineari tripartito: medio elongato bifido. *Linn. Sp. pl. 2. p. 1343.* — *Willd. Linn. Sp. pl. 4. p. 63.*

Orchis radicibus subrotundis, spica longiore flore inermi, labello perangustio quadrifido. *Hall. stirp. helv. n. 1264.*

Orchis flore nudi hominis effigiem representans mas. *Bauh. pin. 82.* — *Mapp. Alf. p. 216.*

Orchis anthropophora oreades. Col. Ecphr. 1. p. 320.

Icon. Hall. l. c. t. 23. — *Mapp. l. c. p. 216.*
Icon bona.

Germ. Menschenähnliche Ophris.

Gall. Ophris homme.

Hab. In Margraviatu superiore in monte dicto Schwärtze retro Oberweiler raro *t. Vulpio* in monte prope Grentzach *t. Stæhelino.* In Alsatia in prato montoso prope Urbisthal *t. Mappo.*

Flor. Majo, Junio. *Semina* Julio matura. 7

Radix testiculata. *Caulis* pedalis et altior, erectus, teres, firmus, supra medium a basi foliosus. *Folia* lanceolata. *Spica* laxa, 3-4 uncias longa. *Flores* numerosi, perpendiculares, odorati. *Petala* quinque, tria exteriora conniventia, ovato-lanceolata, ore purpurea; duo interiora linearia, erecta. *Labium* inferius longum, sublineare, ex luteo-virescens, splendidum, versus basin subvillosum, profunde tripartitum.

ia media elongata, profunde bifida, coo-
l. Sic Galea caput; Lacinae labii latera-
brachia; media: crura cum trunco homi-
aliquatenus repraesentant.

ta. Ad rariores florae oives pertinet, In horto
cultura major evadit.

Mappus plantam circa Argentinam in der
uprechtsau in prato et loco herbido gegen
em Auenheimer-Wörlt citat; ubi eam fac-
ius quidem, sed semper frustra quaesivi.

8. *OPHRIS myodes* bulbis subrotun-
dis, scapo folioso, nectarii labio oblon-
go quadrifido, laciniis duabus mediis
lanceolatis elongatis. *Murr. Linn. Syst.*
veg. p. 813.

Orchis musciflora bulbis subrotundis, caule
folioso, nectarii labio quadrilobo elon-
gato convexiusculo subpubescente, co-
lumna obtusa. *Smith. fl. brit. 3. p. 937.* —
Huds. fl. angl. 2. p. 391.

Ophris insectifera. var. *a. Myodes. Linn.*
Sp. pl. 2. p. 1343

Orchis radicibus subrotundis, labello feri-
ceo bisulco, brachiolis brevibus subula-
tis. *Hall. stirp. helv. n. 1265.*

Orchis muscae corpus referens, minor, ga-
lea et alis herbidis. *Bauh. pin. 83.* — *Mapp.*
fl. Alf. p. 219.

Vesiculus muscarius II. *Tabernaemont. p.*
1048. Icon.

n. Jacqu. Icon. rar. 1, t. 184. — *Hall. l.*
c. t. 24.

m. Fliegenartige Ophris,

II. Ophris mouche.

Hab. Utrunque praesertim in collibus graminosis dumetosis apricis, ad sylvarum marginalia graminea aprica passim. Retro de *Thurnberg*, prope *Grötzingen*, *Weingarten*, *Berghausen* et alibi.

Flor. Majo, Junio. *Semina* Julio. 7

Radix bulbosa, testiculata. *Caulis* subpedalis, erectus, inferne foliosus. *Folia* lanceolata. *Spica* rara 3-4-6 flora, gracilis. *Flora* distantes, perpendiculares, ex violaceo-atropurpurei, muscam referentes. *Galea* aperta. *Petalis* tribus exterioribus viridibus, interioribus duobus, minoribus, angustis, atropurpureis. *Labium* inferius atropurpureum serici toni modo villosum, nitens, caerulea macula angulata, demum albicante notatum, oblongum, quatripartitum, ad basin in foveolam depressum. *laciniis* duabus mediis perpendicularibus, elongatis, lanceolatis, obtusiusculis.

O b. f. Planta cum sequentibus in horto botanico Carlsruhano culta, mutationi non succubuit.

1389. *OPHRIS apifera* bulbis subrotundis, scapo folioso, nectarii labio quinquelobo, lobis subtus inflexis, *Hudsf. fl. Angl.* 2. p. 391. — *Smith. fl. brit.* 3. p. 938. — *Willd. Linn. Sp. pl.* 4. p. 66.

Ophrys insectifera var. i. *Linn. Sp. pl.* 2. p. 1343.

Orchis fucum referens major, foliolis superioribus candidis et purpurascensibus. *Bauh. pin.* 83. — *Mapp. fl. Als.* p. 218. — *Vaill. bot. parif.* p. 146. t. 30. f. 9.

Testiculus VII. sphegodes. *Tabern. p.* 1051. *Icon.*

Orchis fucum referens, major, foliolis superioribus candidis, linea in medio viridi notatis. *Mapp.* 219.

erm. Bienenartige Ophris.

all. Ophris abeille.

a b. In graminosis praesertim calcareo-argillaceis apricis collicosis. Prope *Durlach am Eisenhafen*, retro *Grötzingen*, prope *Berghausen* et alibi passim.

lor. Majo, Junio. Semina Julio matura. ¶

Affinis sequenti speciei a qua differre videtur, *Petalis* tribus exterioribus albidis, *Labio* inferiori convexo, sericeo-villoso, quinque-
do, *lacinia* media subulata recurvata.

390. OPHRIS *aranifera* bulbis subrotundis, caule folioso, nectarii labio convexo villosa, quadrifido, mutico, emarginato. *Smith. fl. brit.* 3. p. 939. — *Linn. Sp. pl.* 4. p. 66.

Ophrys insectifera var. *a.* *Linn. Sp. pl.* 2. p. 219.

Orchis fucum referens, colore rubiginoso. *Bauh. pin.* 83. — *Mapp. fl. Als.* p. 219. *Vaill. bot. parif.* 146. t. 31. f. 15. 16.

Orchis fucum referens, floribus variegatis, *Bauh. pin.* 83. — *Mapp. fl. Als.* p. 219.

erm. Spinnen-Opris.

all. Ophris aranifera.

a b. et *Flor.* Passim cum praecedente. ¶

Nimis affinis praecedenti et sequenti speciei a quibus differre videtur, *Labio* inferiori subrotundo integro, emarginato, convexiori.

Nota. Inter hanc, praecedentem et sequentem, omnino variationes adhuc intermediae sunt, hinc vix specie diversae, quanquam in horto cultae, sat constantes. Hinc Haller et aliique forsitan non immerito sub meris variis sequentis speciei citant. Conf. Hall. *helv.* n. 1266. *B.* p. 135. Idem fere sententiam Ophride crucigera, tenthredinifer, banifera, bombylifera, picta, f. Scolopace et lutea, Willd. *Linn.* pl. 4. p. 64-70.

Obs. Qui amant speciem Ophridis Arachnitis multiplicari, adeant Colles argillaceo gypseas, limari impregnatas apricas gramineas regni Iunii et Arragoniae, ubi certe magnam messurationum pulcherrimarum invenient quod ipsi. Ex nimia multiplicatione specierum sine necessitate inextricabilis confusio oritur, et damnum irremediabile!

1391. OPHRIS *arachnites* bulbis rotundis, scapo folioso, nectarii lato emarginato appendiculato. *Linn. Syst. veg.* p. 813. — *Willd. Sp. pl.* 2. p. 67.

Ophris infectifera var. *γ. Arachnites* *Linn. Sp. pl.* 2. p. 1343. — *Pollic. pal.* n. 858. (*sub Ophride infectifera*)

Orchis fuciflora radicibus subrotundis bello holofericeo emarginato appendiculato. *Hall. stirp. helv.* n. 1266.

Orchis araneam referens petalis superius magnis intense purpureis, rotundis longiori aviculam mentiente, apice in extremitate petali inferioris, *M. fl. Als.* p. 219.

Orchis araneam referens, *Bauh. pin.* 8

on. Curtis. fl. Lond. (sub Ophride apifera).

Hall. l. c. t. 24.

erm. Spinnentragende Ophris.

all. Ophris araignée.

ab. et Flor. Passim cum praecedentibus. ¶

Differt ab *Ophride aranifera*, *Caulc* mai-
folioso, crassiori saepe humiliori, *Spica* bre-
re 2-3-4 flora, *Floribus* majoribus, ara-
m seu Fucum referentibus, *Galea* majori:
alis duobus interioribus latioribus, *Labio*
teriori lato, obtuso subrotundo, emarginato-
pendiculato, sericeo-pubescente, fusco-spa-
co nitente, versus basin tumoribus duobus
is, fuscis notato.

CCCCX. S E R A P I A S.

Linn. Gen. pl. ed. Schreb. n. 1371.

Spathae vagae. *Spadix* simplex. *Perianthi-*
nullum. *Corolla* petala quinque, ovato ob-
ga, erecto-patula, sursum conniventia. *Ne-*
trium longitudine petalorum, basi excavatum,
liferum, ovatum, inferne gibbum, trifi-
a, acutum, *Lacinia* intermedia cordata, ba-
icatrice bifida, tridentata. *Capsula* obovata,
use trigona, tribus carinis adnatis, trivalvis,
carinis dehiscens, unilocularis, *Semina* nu-
rosa, scobiformia.

92. *SERAPIAS latifolia* bulbis fibrosis
foliis ovatis amplexicaulibus, floribus
pendulis. *Murr. Linn. Syst. veg. p. 814.* —
Poll. fl. pal. n. 859.

Epipactis latifolia. Swartz. Act. Holm. 1800,
p. 232. — Willd. Linn. Sp. pl. 4, p. 83.

Epipactis foliis amplexicaulibus ovato lanceolatis, labello lanceolato. *Hall. fl. helv. n.* 1297.

Serapias *Helleborine* α . *latifolia*. *Linn. ph. 2. p.* 1344.

α . Planta junior flore viridi et albescente.

Helleborine latifolia montana. *Bauh. pin.* 186. — *Mapp. fl. Als. p.* 140.

β . Planta paulo vetustior, flore purpureo carneo dilutiori.

γ . Flore et imprimis unguine atrorubente, rotunda, praelonga spica.

Helleborine altera flore atrorubente. *Bauh. pin.* 186. — *Mapp. fl. Als. p.* 141.

Icon. Oeder. fl. Dan. t. 811. — *Hall. l. c. t.*

Germ. Breitblättrige Serapias.

Gall. Helleborine à feuilles larges.

Hab. In sylvis umbrosis passim.

Flor. Julio, Augusto. *Semina* Septembri tura. γ

Radix fibrosa, fibris cylindricis, simplicibus. *Caulis* pedalis, sesquipedalis et altior, erectus, firmus, subvillosus, foliosus. *Folia* alterna, amplexicaulia, lata, breviter, inferiora obtusa, superiora acuta, angustiora, firma, nervosa. *Spica* subracemosa, rotunda, longa, rara, multiflora. *Bracteae* lanceolatas inferiores floribus longiores. *Flores* subnates, subpedicellati, minores quam sequentes specierum. *Galea* petalis ovato-lanceolata, virescenti-albidis, extus villosiusculis. *Lobus* inferius galea brevius, cordiforme, albidius carneum, per senectutem ex fusco purpureo. *Germina* brevia, 2-3 lineas longa, subnata, villosiuscula.

3. *SERAPIAS longifolia* bulbis fibrosis, foliis ensiformibus sessilibus, floribus pendulis. *Murr. Linn. Syst. veg. p. 815.*
— *Poll. fl. pal. n. 860.*

Scirp. palustris foliis lanceolatis amplexicaulibus, bracteis flore brevioribus, floribus pendulis, labello crenato obtuso petalis aequali, germinibus pubescentibus. *Willd. Linn. Sp. pl. 4. p. 84.* — *Swartz. Act. Holm. 1800. p. 232.*

Scirp. foliis ensiformibus caulinis, floribus pendulis, labello obtuso per oras plicato. Hall. stirp. helv. n. 1296.

Scirp. palustris. Scopol. fl. Carn. ed. 2 n. 1129.

Helleborine angustifolia palustris f. pratensis. Bauh. pin. 87. — *Mapp. fl. Alf. p. 141.*

n. Oeder. fl. Dan. t. 267. Hall. l. c. t. 39.

m. Langblättrige Scirpias.

l. Helleborine des marais.

Utrique praesertim in Rheni vicinia ad prata dumetosa uliginosa, nec non in sylvis adjacentibus aquosis, humidis passim frequens. Circa Dachsland, Knielingen, Linkenheim, Kehl et alibi.

Junio, Julio. Semina Augusto matura.

Distinguitur a praecedente specie cum qua us confunditur, *Radice* sub terra longe late reptante, *Caule* crassiore, subflexioso, longioribus, ovali late ensiformibus, ampliori, laxiori, rariori, *Bracteis* latioribus, subbrevioribus, magis reflexis, *Floribus* majoribus, magis pendulis, *Labio* inferogalea longiori, albido, ovato, obtuso,

marginē plicato, *Germinibus* longioribus, uncialibus, *Loco* demum natali demiffiori, mīdo, paluftri, aquofo.

1394. *SERAPIAS lancifolia* bulbis fiffis, foliis ovato-lanceolatis, bracteis longitudine capsulae, floribus erectis, nectarii labio obtuso petalis brevibus. *Murr. Linn. Syst. veg. p. 815. — L. fl. pal. n. 861. var. β.*

Epipactis pallens. Willd. Linn. Sp. pl. 85. — Swartz. Act. Holm. 1800. p. 21.

Epipactis caule paucifloro, lineis in obtusis, labello laevibus. *var. α. latifolia, foliis ovato-lanceolatis. Hall. stirp. helv. 1798.*

Helleborine alba, barba luteola. Rivin. xap. t. 4.

Helleborine flore albo, vel *Damason montanum latifolium. Bauh. pin. 187. Mapp. fl. Alf. p. 140.*

Helleborine flore albo. *Tabernæm. p. 1. Icon bona.*

Icon. Hall. l. c. t. 41. individuum magnus

Germ. Weisgelbe Serapias.

Gall. Helleborine lanciforme.

Hab. In fylvis montosis gramineis et collumetosis gramineis passim non infrequenter. *Retro Grözingen, Berghausen, Weinthal* et alibi cum *Serapiade rubra.*

Flor. Majo, Junio. *Semina* Julio matura

Diffinguitur a sequente specie, *Foliis* ovato-lanceolatis, latioribus, non distichis, & brevioribus, laxioribus, pauciflora, *Bracteis* laevibus.

oribus, longitudine fere floris, inferioribus
gioribus, *Floribus* grandioribus, pauciori-
magis erectis, non totum niveis, sed ex
luteo albidis: *Labio* inferiori apice luteo
fulvo.

95. *SERAPIAS ensifolia* bulbis fibro-
fis, foliis ensiformibus distichis, bracteis
minutissimis, floribus erectis, nectarii
labio obtuso, petalis brevioribus. *Murr.*
Linn. Syst. veg. p. 815. — *Pollich. fl. pal.*
n. 861. var. α . (sub *f. grandiflora*.)

Epipactis ensifolia. *Swartz. Act. Holm.* 1800.
p. 232. — *Willd. Linn. Sp. pl.* 4. p. 85.

Epipactis var. β . foliis ensiformibus. *Hall.*
stirp. helv. n. 1298.

Helleborine montana angustifolia, spicata.
Bauh. pin. 187.

Helleborine angustifolia. *Tabernaemont. p.*
1100. *Icon.*

Oeder. fl. Dan. t. 506. — *Schkuhr. bot.*
handb. t. 274.

erm. Weisse Degenblättrige Serapias.

ll. Helleborine ensiforme.

α *b.* In nemoribus gramineis praesim frequens.
Retro *Wohlfartsweyer, Ettlingen*. Prope
Bauschlott in sylvulis cum *Pyrolae* diversis
speciebus abunde.

lor. Majo, Junio. Semina Julio matura. γ

Distinguitur a praecedente specie, *Foliis*
distichis, lineari-ensiformibus, angustioribus,
magioribus. *Bracteis* multo minoribus, *Flori-*
bus in *Spica* copiosioribus, erectiusculis, ex
toto niveis, minoribus.

b. f. Planta in horto culta, constans.

1396. *SERAPIAS rubra* bulbis fibrosis,
liis ensiformibus, floribus erectis,
ctarii labio acuto. *Linn. Syst. veg. p. 8*
— *Pollich. fl. pal. n. 862.*

Epipactis rubra. Swartz. *Act. Holm. 18*
p. 232. — *Willd. Linn. Sp. pl. 4. p. 8*

Epipactis caule paucifloro, labello lance-
lato, lineis undulatis. *Hall. stirp. helv.*
1299.

Helleborine montana angustifolia purpu-
rens. *Bauh. pin. 187.* — *Mapp. fl. A*
p. 140.

Damafonium flore roseo. *Rivin. hexap. t.*
Icon. Oeder. fl. Dan. t. 345. — *Hall. l.c. t. 4*
Germ. Rothe Serapias.

Gall. Helleborine rouge.

Hab. In sylvaticis, gramineisque collibus da-
metolis passim. Prope *Grötzingen*, *Wein-*
garten, *Berghausen* et alibi.

Radix fibrosa, fibris filiformibus carnosis.
Caulis pedalis et altior, erectus, soliosus. *Fo-*
lia sessilia, alterna, amplexicaulia, lanceolata
acuta. *Spica* rara 3-4-9 flora. *Bracteae* gra-
minibus longiores. *Flores* erecti, magni, amo-
purpurei vel rosei.

Obs. Planta in horto culta mutationi non succubuit.

CCCCXI. CYPRIPEDIUM.

Linn Gen. plant. ed. Schreb. n. 1376.

Spathae vagae. *Spadix* simplex. *Perian-*
thium nullum. *Corolla* petala 4-5 lanceolata
longissima, patentia, erecta. *Nectarium* intra

GYNANDRIA. DIANDRIA. 579

in infimum calceiforme, inflatum, ob-
 , cavum, petalis brevius, latius: *labio*
ore ovato, plano, inflexo, parvo. *Cap-*
 obovata, obtuse trigona, tribus futuris,
 ribus dehiscens angulis, trivalvis, unilo-
 s. *Semina* numerosa, minima. *Recepta-*
 lineare, singulae valvulae pericarpium lon-
 galiter adnatum.

1. **CYPRIPEDIUM** *Calceolus* radi-
 cibus fibrosis, foliis ovato-lanceolatis
 caulinis. *Linn. Sp. pl.* 2. p. 1346.

Cypripedium lobo styli ovali, concavo, sub-
 tus canaliculato carinato, labello petal-
 is breviori compresso. *Swartz. Act.*
Solm. 1800. p. 251.

Calceolus radicibus fibrosis, foliis ovato-
 lanceolatis. *Hall. stirp. helv. n.* 1300.

Calceolus marianus. *Mapp. fl. Als.* p. 49.

Calceolus flore majore. *Mapp. l. c.*

Calceolus marianus flore exalbido. *Mapp. l. c.*

Redouté plant. liliac. — *Hall. l. c. t.* 43.

— *Schkuhr. bot. handb. t.* 275.

n. Frauenschuh. Marienschuh.

Sabot.

In ditone Basileensi prope *Wallenburg*
Zwingero. In Principatu superiore prope
alem et alibi in sylvis passim non infre-
 uens. In Principatu *Fürstenbergensi* pas-
 m. In Alsatiae monticulis dumetolis gra-
 minosis inter *Dorelsheim* et *Mutzig* fre-
 uens. *β.* retro *Barr* in valle in vineis et

prope *Heiligenstein*. γ. in vineis prope
Heiligenstein t. Mappo.

Flor. Majo, Junio. *Semina* Julio matur

Radix fibrosa, repens. *Caulis* pedali
quipedalis, erectus, foliosus. *Folia* sessili
plexicaulia ovato lanceolata, nervosa.
1-2 speciosi, pulcherrimi, pallide luteo
purpureo tincti, raro exalbidi. *Petala* qu
lanceolata, patentia. *Labium* praeamplum
ceum ligneum repraesentans, ovatum, c
flavum, intus maculis lineisque pilosis
ditum.

Planta omnino elegans, in hortis facile cie

HEXANDRIA.

CCCCXII. ARISTOLOCHIA.

Linn. Gen. pl. ed. Schreb. n. 138.

Calyx nullus. *Corolla* monopetala,
loba, irregularis: basis ventricosa, subgl
torulosa: *Tubus* oblongus, hexagono-cyl
ceus. *Limbus* dilatatus, inferne extensus
quam longam. *Stamina Filamenta* nulla
therae sex, subtus stigmatibus adnatae, q
loculares. *Capsula* sexangularis, sexloc
Semina plura, depressa, incumbentia.

1398. ARISTOLOCHIA *Clematit*

liis cordatis, caule erecto, floribu
laribus confertis. *Linn. Sp. pl. 2. p.*
Poll fl. pal. n. 863. — Hall. stirp
n. 1029.

Aristolochia Clematidis recta. *Bauh. pi*
— Mapp. fl. Alf. p. 27.

Aristolochia Clematidis. *Tabern. p. 1141. Ic.*
o n. Bulliard. herb. t. 39. — Blackw. t. 255.
Schkuhr. bot. handb. t. 276.

r m. Osterlúcey.

ll. Aristolochie Clematite.

b. Utrunque passim in vineis frequens. Colitur passim in hortis.

or. Julio, Augusto. Semina Septembri matuta. 7

Radix simplex, longa, cylindrica, tetra-
da, descendens, extus ex viridi-lutea, intus
ida. Caules ex una radice plures, cubitales,
pedales et altiores, erecti, teretiusculi, striati,
pliciusculi, glabri. Folia alterna, petiolata,
obcordata, obtusa, integra, venosa, gla-
bra, margine aspera. Flores axillares, conferti
et pedunculati, nudi, lutei.

arm. ARISTOLOCHIAE VULGARIS Ra-
dix; Herba.

Planta trita, aromatica, graveolens, amara, in-

tes, resolventes, emmenagogas, et vulnerarias

habet. Domestica nostratibus medicina, hinc saepe

ficorum hortis colitur.

Equorum ulceribus, foliorum pulvis et succus imit-

r. Praestans saepe in lue bovilla medicina, nimis in

his contagiosis neglecta, magis adhuc propaganda,

Flores apibus mel praebent.

ota. Aristolochia Siphon L'Heritier stirp.
13. t. 7. seu A. macrophylla foliis cordato-
subrotundis amplissimis, caule frutescente volu-
bili, floribus binis incurvatis, limbo triangulari,
plano. Lamarck. Encycl. 1. p. 252. in Pen-
sylvania obveniens, hospes in hortis Carlsruha-
nis. Individuum pulcherrimum praealtum in horto
Generosissimi Lttneri in Heitersheim Bris-
goviae nunquam satis admirandum, Populum
italicam excelsam obvolvens, fere suffocans.

CLASSIS XXI.
MONOECIA.

MONANDRIA.

CCCCXIII. ARUM. *Spatha monophylla, cucullata. Spadix supra nudus, inferne foemineus, medio masculus.*

CCCCXIV. CALLA. *Spatha plana. Spadix nudus flosculis. Cal. o. Cor. 6. Baccae polyspermae.*

CCCCXV. ZANNICHELLIA. ♂ *Cal. nullus. Corolla nulla.*

♀ *Cal. 1-phyllus. Corolla nulla. Pist. 4. Sem. 4.*

CCCCXVI. ITTNERA. ♂ *Cal. nullus. Corolla nulla. Stam. Filamentum multum. Anthera denticulata.*

♀ *Juxta masc. florem. Cal. nullus. Corolla nulla. Stylus filiformis. Stigma 2-3-fidum acutum. Caps. ovali-cylindrica, 1-locularis, crustacea, monosperma.*

MONOECIA.

579

CXVII. CHARA. ♂ *Cal.* nullus. *Corolla*
nulla.

♀ *Cal.* 4-phyllus. *Cor.* nulla.
Stigm. 3-fidum. *Sem.* unicum.

• *Callitriche verna.*

DIANDRIA.

CXVIII. LEMNA. ♂ *Cal.* 1-phyllus. *Cor.*
nulla.

♀ *Cal.* 1-phyllus. *Cor.* nulla.
Styl. 1. *Capf.* 1-locularis.

TRIANDRIA.

CXXI. ZEA. ♂ *Gluma.* 2-flora. *Glum* 2-
valvis.

♀ *Glum.* 1-flora. *Glum.* 2-
valvis. *Styl.* 1. *Sem.* 1. nu-
dum, subrotundum.

CXXII. CAREX. ♂ *Ament.* 1 florum. *Cor.*
nulla.

♀ *Ament.* 1-florum. *Cor.* o.
Stylus 1. *Sem.* 1-tunicatum.

CXX. SPARGANIUM. ♂ *Cal.* 3-phyllus.
Cor. nulla.

♀ *Cal.* 3-phyllus. *Cor.* nulla.
Stigm. 2. *Sem.* 1-sperma.

CXIX. TYPHA. ♂ *Cal.* 3-phyllus *Cor.* nulla.

♀ *Cal.* capillaris. *Cor.* nulla.
Styl. 1. *Sem.* 1. pappigerum.

TETRANDRIA.

CCCCXXV. *URTICA*. ♂ *Cal.* 4-phyllus. *Cor.* nulla. *Nectar.* cyathiforme.

♀ *Cal.* 2 valvis. *Cor.* o. *Stigma* villosum. *Sem.* 1. ovatum.

CCCCXXVI. *MORUS*. ♂ *Cal.* 4-partitus. *Cor.* nulla.

♀ *Cal.* 4-phyllus. *Cor.* o. *Styl.* 2. *Sem.* 1. baccatum.

CCCCXXIV. *BUXUS*. ♂ *Cal.* 3-phyllus. *Cor.* 2-petala.

♀ *Cal.* 4-phyllus. *Cor.* 3-petala. *Stigma* 3. *Caps.* 3-locularis.

CCCCXXIII. *BETULA*. ♂ *Ament.* 5-florum. *Cor.* 4-partita.

♀ *Ament.* 2-florum. *Cor.* o. *Styl.* 2. *Sem.* 1. ovatum.

PENTANDRIA.

CCCCXXVII. *XANTHIUM*. ♂ *Cal.* commut. polyphyll. *Cor.* 5-fida. *Filam.* connexa.

♀ *Cal.* o. *Cor.* o. *Styli.* 2. *Drupa* 2-locularis.

CCCCXXVIII. *AMARANTHUS*. ♂ *Cal.* propr. 5-phyllus. *Cor.* o. *Stam.* 3 5.

♀ *Cal.* propr. 5-phyllus. *Cor.* o. *Styl.* 3. *Caps.* circumscissa.

MONOECIA.

581

POLYANDRIA,

XXXI. *SAGITTARIA*. ♂ *Cal.* 3-phyll.
Cor. 3-petala. *Stam.* 24 cir-
citer.

♀ *Cal.* 3-phyll. *Cor.* 3-peta-
la. *Pistilla* numerosa. *Semi-*
na numerosa.

XXX. *MYRIOPHYLLUM*. ♂ *Cal.* 4-
phyllus. *Cor.* o. *Stam.* 8,
♀ *Cal.* 4-phyllus, *Cor.* o. *Pist.*
4. *Sem.* 4.

XXIX. *CERATOPHYLLUM*. ♂ *Cal.* sub-
7-partit. *Cor.* o. *Stam.* 18.
circiter,
♀ *Cal.* sub-7-partit. *Cor.* o.
Pistil. 1. *Sem.* 1.

XXXII. *POTERIUM*. ♂ *Cal.* 4-phyll.
Cor. 4-partitu. *Stam.* 32. cir-
citer.

XXXV. *FAGUS*. ♂ *Cal.* 5-fidus. *Cor.*
nulla. *Stam.* 12. circiter.
♀ *Cal.* 4-fidus. *Cor.* o. *Styl.* 3.
Caps. 2-sperma.

XXXIII. *QUERCUS*. ♂ *Cal.* 5-fidus. *Cor.*
o. *Stam.* 10. circiter.
♀ *Cal.* integer. *Cor.* o. *Styl.*
5. *Nux* coriacea.

Arum italicum acaule, foliis hastatis, auriculis divaricatis, spadice drico luteo *Lamarck. Encycl.* 3. p.

Arum folio lato atro-viridi margine cincto, sparsis maculis albis gatis. *Sabbat hort. rom.* 2.

Arum venis albis italicum maximum. *nes. inst.* 158.

Icon. Sabbat. l. c. t. 75.

Germ. Ueberwinternder Aron. Großer Italiänischer Aron.

Gall. Pied-de veau hyemale ou d'Italie

Hab. In dumetis apricis prope *Beurtheim* *Carlsruhe* et alibi passim frequens. In *govia* ad montem *Kaiserstuhl* in du prope *Burgheim* et *Sponeck*.

Flor. Majo, Junio. *Baccas* Julio, Augusto matura. 7

Radix tuberosa, fibrillosa, alba, ca *Folia* radicalia longe petiolata, maxima lato-sagittata, auriculis divaricatis magn perne saturate viridia, eleganter ex viri bide venosa, subtus pallidiora, utrinque splendidia, vernice quasi obducta, nov cias ad pedem longa, septem ad octo lata, hyeme perdurantia, laetissima. S cylindrico-clavatus, spatha brevior, spe vitellinus, seu luteolus. *Spatha* erecta, ciosa, pedalis, intus albida, extus pallide *Baccas* magnas, coccineas. *Semina* rot reticulata.

Speciosa omnino planta ob folia pulcherrima venosa, per hyemem perdurantia, auriculis prædita.

Specie omnino a sequente diversa. Folia in hyemantia, semper multo majora, in altera,

lla, Spadix major semper luteus, non violaceo-
pureus.

2 f. Planta jam ultra viginti annos in horto Botanico
cultā, speciosissima, mutationi non succubuit.

Nota. Incalescentiam Spadicis, durante fructifica-
tione, de qua primus Ill. Lamarck mentionem
fecit, in nostra specie italica ultra octode-
cim annos in horto botanico Carlsruhano cum
Schweyckerto hortorum Inspectorē obser-
vavi.

Planta degustata acerrima, mordens. Sequentis
es habet.

400. *ARUM maculatum* acaule, foliis
hastatis integerrimis: spadice clavato.
Linn. Sp. pl. 2. p. 1370. — Poll. fl. pal.
n. 864.

Arum foliis sagittatis, spatha recta, clava
cylindrica. *Hall. flirp. helv. n. 1302.*

Arum vulgare maculatum et non maculatum.
Bauh. pin. 195. — Mapp. fl. Alf. p. 28.
29. 30.

Arum I. II. III. *Tabernaemont. p. 1123. 1124.*
Icones.

• *Arum* folio subrotundo maculatum et non
maculatum, *Mapp. fl. Alf. p. 29.*

• *o n. Curtis. fl. Lond. 114. — Bulliard, herb.*
t. 25. — Blackw. t. 228.

erm. Arum. Aronskraut,

gll. Pied-de-veau.

• *ab. Utrique passim in sylvis umbrosis, ad*
sylvarum marginalia, in saepibus humidius-
culis frequens.

• *or. Junio, Julio. Baccae Augusto Septembri*
maturæ. 7

Radix tuberosa, subrotunda, fibrillosa,
tus nigra, intus alba, carnosa: fiocata farina.

586. MONOECIA. MONANDRIA.

cea. *Folia* radicalia petiolata, hastata, inter viridia, maculis nigris, cinereis aut nullis tuta, juniora minora, subflagittata, saepe ovata. *Spatha* erecta, intus albida. *Spadix* clavatus, superne nudus, ex violaceo-amore fime purpureus. *Baccae* coccineae.

Pharm. ARI Radix.

Tota planta degustata, initio mitis, post acerrima, pungens.

Radix siccatione mitiganda, saponacea, incipit resolvens, stomachica. Siccata farinacea, amilacea triens, mitis.

CCCCXIV. CALLA.

Linn. Gen. pl. ed. Schreb. n. 1388.

Masculi flores foemineis intermixti.

tha monophylla, ovato-cordata, acuminata superne colorata, maxima, patens, persistens. *Spadix* digitiformis, simplex, erectus, fructificationibus tectus. *Perianthium proprium* nullum. *Corolla* nulla. *Stamina Filamentum* singulum compressum, truncatum. *Anthera* depressa. *Feminei Spatha* et *Spadix* communis cum culis. *Perianthium proprium* nullum. *Corolla* nulla. *Germen* singulum subrotundum, obtusum. *Stigma* acutum. *Bacca* tetragono-globosa, posita, unilocularis. *Semina* plura, oblonga, lindracea, utrinque obtusa.

1401. *CALLA palustris* foliis cordatis, spatha plana, spadice undique herboso. *Linn. Sp. pl. 2. p. 1373*
Poll. fl. pal. n. 865:

MONOECIA. MONANDRIA 587

Dracunculus palustris f. radice arundinacea.
Bauh. pin. 195.

Arum palustre radice arundinacea. *Mapp.*
fl. Alf. p. 30.

Dracontium palustre. *Tabernæmont. p.* 1122,
Icon.

On. *Oeder. fl. Dan. t.* 422. — *Schkuhr. bot.*
handb. t. 278.

rm. *Froschlöffel. Wasser-Aron. Wasser-*
Schlangenwurz.

ll. *Serpentaire aquatique.*

b. In Margraviatu superiore in paludosis ad
rivulos in alnetis prope *Haussen t. Hebelio.*
In Alfatia juxta stagnum prope *Still. t. Map-*
po. In Platinatu circa *Lautern* in palu-
dibus turfosis copiose cum *Scheuchzeria*
palustri. t. Pollichio.

or. Junio, Julio. *Semina* Augusto, Septem-
bri matura. z

Radix cylindrica, elongata, longe lateque
tens, obliqua, crassitie digiti, laevis, nitida,
dicea, geniculata, fibrosa: *fibris* filiformi-
bus approximatis, numerosis, elongatis, albis,
geniculis subtus exeuntibus, descendentibus.
Radicalia radicalia longe petiolata, cordata, acuti-
cula, integerrima, firma, venosa, laete viri-
da, glabra. *Scapus* palmaris et altior, erectus,
terres, teres, crassiusculus, glaber. *Spadix* ova-
tus, obtusus, terminalis. *Spatha* ovato-subro-
bata, integra, plana, superne alba, glabra.
Pericarpia 6-9 spermae, amoene rubro-purpureae,
viscidulae, mucilaginosae.

Ad rariores omnino Florae cives pertinet.

Radix acris, masticata primum insipida, dein pun-
gentius, perdurans, fervida, molestissima, manifestans

rectum. *Anthera* sagittata, erecta.
juxta masculum florem. *Calyx* mono-
ventricosus, tridentatus. *Corolla* nu-
mina 4: 8 corniculata, conniventia. S-
dem, patentiusculi. *Stigmata* ovata, p-
tentia extrorsum. *Capsulae* totidem sul-
erecto-patentes, stylo reclinato rostrati-
tuberculatae, crustaceo-coriaceae, co-
uniloculares, e valves. *Semina* solitaria
ga, hinc gibba,

1402. *ZANNICHELLIA palustris*
ra quadriloculari, stigmatibus
mis, feminibus dorso denticula-
libus filiformibus articulatis, fol-
fitis angustis linearibus, apice em-

Zannichellia palustris. *Linn. Sp. pl.*
— *Poll. fl. pal. n.* 866. — *H*
helv. n. 1604.

Zannichellia palustris anthera qu

.MONOECIA. MONANDRIA. 589

Jgoides vulgaris. *Vaill. act.* 1719. p. 15.
t. 1. f. 1.

otamogeton capillaceum capitulis ad alas
trifidis. *Bauh. pin.* 193. *Prodr.* 101.

n. *Oeder. fl. Dan.* t. 67. — *Schkuhr. bot.*
handb. t. 280.

m. *Sumpf-Zannichellia.*

l. *Zannichelle aquatique.*

i. In paludibus, stagnis, fossis aquaticis passim frequens. Inter *Carlsruhe* et *Rintheim* in foveis aquosis cum *Lindernia pyxidaria*. In Margraviatu superiore in rivulo inter *Buggingen* et *Dattingen*. In ditione Basileensi prope *Riechen* in rivulo, prope *der Holzmatt.* t. *Stachelino*. In Allatia circa Argentinam in loco ubi Brusca Ellam intrat frequens, nec non in *dem Rheingiesen*. In Palatinatu in paludibus stagnantibus passim inter *Oggersheim* et *Frankenthal* abunde post *Pollichium* vidi. In Sponhemia in fossis aquaticis prope *Kirchberg* maxima in copia.

r. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caules filiformes, tenues, ramosi, glabri, virides, saepe albidi. *Folia* opposita, liliifolia, apice emarginata, glabra, sessile ad binas uncias longa. *Flores* axillares, subsessiles, minuti. *Germina* in floribus inclusis saepissime quatuor, glabra, stylo perite incurvo coronata.

Juxta antheram nudam albidam quadrilocularem non raro flos foemineus ad marginem conspicius.

t a. Non confundenda cum *Potamogetone*.

lanta avibus insectisque aquaticis accepta.

nullus. *Corolla* nulla. *Stylus* filiformis
2-3 fidum acutum. *Capsula* ovali-cylindrica
locularis, evalvis, glabra, crustacea
sperma.

1403. *ITTNERA Najas* caule
foliis oppositis, fasciculatisque
bus, sinuato denticulatis
floribus sessilibus axillaribus
busque.

Najas marina. Linn. *Sp. pl.* 2. p.

Najas monosperma caule muricato
monospermo. Willd. Linn. 8
331.

Najas fructu monospermo. Hall.
n. 556.

Fluvialis vulgaris latifolia. Vaill.
p. 17. t. 1. f. 2.

Fluvialis latifolia fructu minus olivaceo
nospermo. Michel. Nov. Gen.
t. 8. f. 2.

MONOECIA. MONANDRIA. 891

ab. In regionibus rhenanis in rivis segnioribus et locis aquosis inundatis prope *Dachland in der Federbach* Rhenum versus, et in confiniis rivulis passim cum sequente frequens. Circa Argentinam cum sequente.

o r. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Caulis pedalis, cubitalis, et altior cre-
s, dichotomus, articulatus, glaber aut den-
sus foliaceo - carilagineis raris praeditus. *Fo-*
opposita, terna, suprema fasciculata, om-
i sessilia, linearia, sinuato - dentata, subcar-
iginea, laete viridia, sesquilineam ad binas
eas lata, ultra unciam ad sesquiunciam lon-
, fragilia. *Flores* axillares feminei, subsolita-
Flores masculi femineique Calyce et Corolla
rituti. *Filamentum* nullum. *Anthera* superne
trifida, denticulata, initio subrubella termi-
les feminei, masculique tecti. *Capsula* ovata,
ea, stigmate recto, bifido saepe trifido ter-
nata.

sf. Planta saepe in regionibus rhenanis lecta, omni-
no monoica nec dioica est. *Antherae* non qua-
tuor sed unica, subtrifida.

Nota. Hinc nostram plantam in Monecia non tetra-
tia, sed monandria posui.

Linneus in Horto Cliffortiano p. 437.
recte ad monoeciam monandriam citat; in *Spe-*
ciebus plantarum autem Dioeciae monandriae
amandavit.

sf. Nudis oculis flores masculi non conspicui sunt,
sed Microscopio in ramulis extremis plantae omni-
no inveniuntur.

sf. Planta, excepta fructificatione omni ex parte egre-
gie quadrat cum figuris *Michelianis* t. 8. f. 1. 2.
Corollam autem tubulosam limbo quatripartito re-
flexo praeditam, l. c. exhibitam, nunquam vidi

qui Horticulturam, Pomologiam et Viticulturam
vise, adauxit. Artium et scientiarum prome-
nitorum Maecenas!

1404. *ITTNERA minor* caule di-
foliis caulinis oppositis, rame-
summis congestis, lineari-subu-
curvatis, sinuato denticulatis rig-
bus sessilibus axillaribus termina-

Najas marina. var. B. Linn. Sp. pl.


Najas minor. Roth. fl. germ. 2. p. 1.
Lamarck et Decand. fl. franc.

Najas subulata. Thuil. fl. paris. 2. p. 1.

Caulinia fragilis foliis ternis oppo-
neari-subulatis recurvatis, acul-
tatis rigidis. *Willd. Linn. Sp. pl.*

Fluvialis minor foliis angustissimis
latis deorsum reflexis, fructu a-
jori monospermo. *Michel. gen.*
t. 8. f. 3.

Fluvialis angusto longoque folio. V.
paris. 1719. p. 17.



1

2

3

4

5

6

7

all. Ittnera mineur.

ab. Cis et trans Rhenum in rivulis segnioribus passim frequens. Prope *Dachsland in der Federbach* et in aquis adjacentibus cum *Marsilea quadrifolia*, *Riccia fluitante* et *Conservea reticulata* non infrequens; Prope *Knielingen*, *Eckenstein*, *Schröck* et alibi, prope *Hochstädt* cum *Salvinia natante*. In Alsatia circa Argentoratum versus *Oßwald* in Ellae fluvio et alibi.

or. Julio, Augusto. *Semina* Augusto, Septembri matura. ☉

Planta elegantissima, cartilaginosa, rigida, de viridis. *Caulis* digitalis, palmaris et longior in fundo aquarum, erectus, maxime ramosis, dichotomus, rigidus, articulatus. *Folia* alina opposita, lineari-fetacea, sinuato-artedentato-aculeolata, recurva, fragilia; *radix* subterna, suprema in rosulam congesta, niunciam adunciam longa, lineam dimidiam a. *Flores* 1-2-3 axillares, sessiles, terminaque, minuti, foliis occulti. *Anthera* antehiscentiam amoene rubella, apice albida. *Stigma* acutum simplex, bifidum, rarius tritum.

Nota. Caulinae nomen Zofterae oceanicae Linn. Mant. 123. a Lamarckio et De-candollio in fl. franc. 3. p. 156. datum est. quia Caulini hanc plantam in peculiari Dissertatione 1792. descripsit.

CCCCXVII. C H A R A.

Linn. Gen. plant. ed. Schreb. n. 1397.

Masculus flos ad basin feminei; extra caly-
m. *Calyx* nullus. *Corolla* nulla. *Stamen* Fi-
Flora. Bed. Tom. III. P p

MONOECIA. MONANDRIA.

lamentum nullum. *Anthera* globosa, germini extra calycem subjecta, unilo non dehiscens. *Femineus* flos. *Calyx* tet-
lus: *foliolis* subulatis, inaequalibus, pa-
tibus; vel nullus. *Corolla* nulla. *Germe*
natum, striis quinque spiralibus inse-
Stylus nullus. *Stigmata* quinque, fir-
Bacca ovato oblonga, spiraliter striata,
cularis, intra crustam tenuissimam semina-
tinens. *Semina* plurima, sphaerica, minu-

1405. *CHARA vulgaris* caulibus
bus (frondibus) foliis interne d
Linn. Sp. pl. 2. p. 1624. — *Poll.*
n. 1135.

Chara caule laevi fragili. *Hall. Stirp.*
n. 1681.

Chara caule herbaceo rigido, ramuli-
ceis patulis. *Weiss. Crypt.* p. 32.

Chara vulgaris foetida. *Vaill. Act.* 1719
t. 3. f. 1.

Hippuris foetida. *Dillen. Giesf. p.* 105
gen. p. 88. t. 2. — *Mapp. fl. Als.* 1

Equisetum foetidum sub aqua repens.
pin. 16. *Prodr.* 25.

Icon. Oeder. fl. Dan. t. 150.

Ger m. Gemeiner Armleuchter.

Gall. Charagne vulgaire.

Hab. Utrunque in aquis stagnantibus et
aquaticis Rheno vicinis passim abund

Flor. Junio, Julio. *Semina* Augusto matu-

• **MONOECIA. MONANDRIA.** 595

Caules filiformes, articulati, glabriusculi, el scabro-crustaceo-tuberculati, ramosi, fra-
iles, in aqua natantes, extra aquam cinereo-
lbidi, siccitate fragilissimi, albidii. *Folia* ver-
cillata, tereti-subulata, rigida, obscure viri-
ia. *Semina* in axillis foliorum orbicularia, phoe-
icea, pulchella.

Maxima saepe in quantitate fossas aquaticas implet.

Odor plantae peculiaris, foetidus.

406. *CHARA tomentosa* caulibus sub-
tomentosis fragilibus.

Chara tomentosa aculeis caulinis ovatis.
Linn. Sp. pl. 2. p. 1624.

Chara caule subtomentoso fragili. *Hall. stirp.
helv. n. 1683.*

Equisetum fragile majus subcinereum aquis
immersum. *Morif. hist. 3. p. 621. Sect.
15. t. 4. f. 9.*

Equisetum seu *Hippuris lacustris*, foliis manfu
arenosis. *Plucken. Alm. 135. t. 29. f. 4.*

erm. Filziger Armleuchter.

all. Charagne cotonneuse.

ab. In Stagnis et paludibus passim.

lor. Junio, Julio. Semina Augusto matura. ☉

Caules filiformes, striati, articulati, ramosi,
auci, farinoso-subtomentosi, fragiles. *Rami*
articillati 6-7. *Folia* tereti-subulata. *Semina*
cillaria, phoenicea, orbiculata.

407. *CHARA hispida* caulibus hispidis,
ramulis scabris basi subfoliosis, articulis
foliosis teretibus, subulatis, bracteis (ca-

lyce) bacca longioribus. *Willd. Sp. pl.* 4. p. 185.

Chara hispida aculeis caulinis capilla confertis. *Linn. Sp. pl.* 2. p. 1624.

Chara caule fragili spinoso. *Hall. stirp.* n. 1682.

Chara major, caulibus spinosis. *Vaill.* 1719. p. 18. t. 3. f. 3.

Hippuris muscosa setis per extremum fistis. *Dill. Eph. N. C. Cent.* 6. app. p. t. 13. f. 1. — *Mapp. fl. Alf.* p. 146.

Icon. Oeder. fl. Dan. t. 154.

Germ. Stachlichter Armleuchter.

Gall. Charagne heriffée.

Hab. et Flor. cum praecedente passim.

Affinis praecedenti a qua differt, *Cau-* spinulosis, *Spinulis* copiosis tereti - subul patentibus, demum reflexis. *Calycinis* foli seu *Bractea* *Bacca* longiore.

Obs. Amat stagna praesertim salsa; maxima abundantia in Cataloniae salis, maria



Leon. Schkuhr. bot. Handb. t. 280.

Germ. Biegsamer Armleuchter.

Gall. Charagne flexible.

Hab. In rivulis limpidis passim. Prope *Ettlingen* in aquis limpidissimis, in aquis horti aulici *Ettlingensis* frequens.

Flor. Junio. *Semina* Julio matura. ☉

Caules tenues, filiformes, articulati; glari, laete virides, diaphani, aphylli. *Rami* verticillati 5-8-9, compressi, unciam ad binas uncias longi, inarticulati 2-3 furci, diaphani, glari. *Granula* feminifera in bifurcatione ramorum 3-4-5 aurantia, exigua, pulchella, *Bractea* destituta.

MONOECIA DIANDRIA.

CCCCXVIII. L E M N A.

Linn. Gen. pl. ed. Schreb. n. 1400.

Masculus flos. *Calyx* monophyllus, subrotundus, latere dehiscens, oblique extrorsum dilatatus, obtusus, patens, depressus, magnus, integer. *Corolla* nulla. *Stamina Filamenta* duo, tubulata, incurva, longitudine calycis. *Antherae* didymae, globosae. *Femineus* flos. *Calyx* 5 in masculo. *Corolla* nulla. *Germen* subovatum. *Stylus* brevis persistens. *Stigma* simplex. *Capsula* globosa, cum acumine, unilocularis. *Semina* nonnulla, oblonga, utrinque acuta, hinc triata.

b.f. Flores hermaphroditos viderunt Cl. EHRHART et HEDWIG.

MONOECIA. DIANDRIA.

1409. *LEMNA trifulca* foliis peti-
lanceolatis. *Linn. Sp. pl. 2. p. 137*
Poll. fl. pal. n. 867.

rophace caule filiformi, foliis elli-
anceolatis viviparis. *Hall. stirp. he*
1901.

Lenticularia ramosa, monorhiza, folii
longis, pediculis longioribus do
Michel. Nov. gen. 16. — Mapp. fl.
p. 167.

Lenticula equatica trifulca. *Bauh. pin.*
Hedera palustris. Tabernæmont. p.
Icon mala.

Icon. Wolf. Diff. de Lemna. f. 1. 2. 3. M
l. c. t. 11. f. 5.

Germ. Groses Entengrün.

Gall. Lentille d'eau trifulque.

Hab. Utrique in stagnis praesertim Rhei-
cinis passim abunde. Prope *Dachsla*
alibi cum *Marsilea quadrifolia* et *Sc*
ckerta nymphoide in superficie aquae
copiose.

Flor. Junio, Julio. *Semina* Augusto mat-
Radiculae solitariae. *Cauliculi* compl-
tenerrimi, decussati. *Folia* ovata, acuta
fulca, simpliciave, integra, in superficie
rum natantia, laete viridia, glabra.

Obs. Flores hujus speciei nuper primus detexit
et descripsit l. c.

1410. *LEMNA minor* foliis sessilibus
que planiusculis, radicibus foli-
Linn. Sp. pl. 2. p. 1376. — Polli
pal. n. 868.

Hydrophace monorhiza foliis planis ovatis.
Hall. stirp. helv. n. 1900.

Lenticularia minor monorhiza foliis utrinque
viridibus, subrotundis. *Michel, nov. gen.*
16. — *Mapp. fl. Alf. p. 167.*

Lenticularia palustris vulgaris. *Bauh, pin. 262.*
Tabernæmont. p. 890. Icon.

non. Wolf. l. c. f. 4-10. — Schkuhr. bot.
handb. t. 281. — Michel. l. c. t. 11. f. 3.

erm. Kleinè Meerlinse,

all. Petite Lentille d'eau,

ab. Utrinque in aquis stagnantibus passim
abunde.

lor. Junio, Julio, Semina Augusto matura.

Radicula solitaria, simplex. Folia elliptica,
inuta, utrinque planiuscula, laete viridia, na-
tia, aquas stagnantes saepe obducentia.

Planta avibus aquaticis, praesertim anatis ac-
ta.

LEMNA gibba foliis sessilibus sub-
tus hemisphaericis, radicibus solitariis.
Linn. Sp. pl. 2. p. 1377.

Hydrophace monorhiza, foliis pulvinatis in-
ferne convexis. *Hall. stirp. helv. n. 1898.*

Lenticularia palustris major, inferne magis
convexa, fructu polyspermo. *Michel.*
nov. pl. gen. 15. — Mapp. fl. Alf. p. 165.

non. Wolf. l. c. f. 16-21. — Schkuhr. bot
handb. t. 281. — Michel. l. c. t. 11. f. 2.

erm. Convexe Meerlinse.

all. Lentille d'eau hemisphaerique.

Hab. In aquis stagnantibus passim rarius. Pro
Dachsland, Knielingen et alibi in Rh
vicinia,

Flor. Junio, Julio. ☞

Affinis praecedenti, sed *Folia* conve
hemisphaerica, pulvinata.

1412. *LEMNA polyrhiza* foliis sessilib
radicibus confertis. *Linn. Sp. pl.* 2.
1376. — *Poll. fl. pal.* n. 869.

Hydrophace polyrhiza foliis ovatis pl
Hall. stirp. helv. n. 1899.

Lenticularia major polyrhiza, inferne at
purpurea. *Michel. nov. gen. pl.* 16.
Mapp. fl. Als. p. 167.

Icon. Schkuhr. bot. Handb. t. 281. — *Wo*
l. c. f. 16. 21. *Michel. l. c. t.* 11. f. 1.

Germ. Vielwurzliche Wasserlinsen.

Gall. Lentille d'eau a plusieurs racines.

Hab. In aquis segnioribus et stagnantibus c
L. minore passim abunde.

Flor. Junio, Julio. *Semina* Augusto matura

Radiculae fasciculatae. *Folia* oblongo-c
data, obtusa, crassiuscula, superne laete v
dia, inferne atropurpurascencia, supra aqu
natantia.

Obs. Omnes Lemnae sunt polygamae, ut Wol
Dissertatione egregia de Lemna, Altorfii i
edita, observationibus demonstravit.

MONOECIA TRIANDRIA.

CCCCXIX. TYPHA.

Linn. Gen. plant. ed. Schreb. n. 1401.

Flores masculi numerosi in amento
mum terminante. *Amentum* commune cy

um, densissimum, constans perianthiis pro-
s triphyllis, setaceis. *Corolla* nulla. *Stamina*
umenta tria longitudine calycis. *Antherae*
ongae, pendulae. *Feminei flores* numerosi
mento culmum eundem cingente, compac-
me digesti. *Calyx* nullus. *Corolla* nulla. *Ger-*
is insidens setae, ovatum. *Stylus* subulatus.
Stigma capillare, persistens. *Fructus* numerosi,
indrum constituunt. *Semen* unicum ovatum,
o instructum, setae insidens. *Pappus* capil-
latus, setae seminiferae a basi ad medium affixus,
gitudine pistilli.

13. *TYPHA latifolia* foliis subensiformibus spica mascula femineaeque approximatis, utraque cylindrica. *Willd., Linn., Sp. pl.* 4. p. 197. — *Linn. Sp. pl.* 2. p. 1377. — *Poll. fl. pal.* n. 870.

Typha clava unica. *Hall. stirp. helv.* n. 1305.

Typha palustris major. *Bauh. pin.* 20. *Mapp. fl. Als.* p. 315.

Don. Curtis. fl. Lond. t. 17 r, (*sub Typha majore*). — *Schkuhr. bot. handb.* 281.

Arm. Grofse breitblättrige Rohr - Kolben, Grofse Lieschkolben.

Il. Massettes à feuilles larges.

b. Utrinque passim ad Rheni aquas placidas et alibi.

r. Julio, Augusto. *Semina* Septembri, Octobri matura. ¶

Radix repens, articulata, fibrillosa. *Cul-*
mus 4-7 pedalis, erectus, simplex, foliosus,

teres, glaber. *Folia* alterna, erecta, prae vagina culmum involventia, ensiformia, sub plana, glauca, unciam fere lata, crassa *thae* duae deciduae, una ad spicae marginem, altera ad ejus medium. *Amentum* (spica terminalis) erectum, pedale et locum cylindricum, fuscum, ultra unciam latum.

Vietoribus nostratibus folia ad obturandas doliorum inserviunt.

Pappus copiosus cum lana mixtus pro tegumentorum aptus, quo etiam pulvinaria infarserunt rici. Kalm.

Lanugo Typhae, axungia permixta ab animalibus partibus tumorem et inflammationem arcet.

Nota. Planta in fossis ad ambulacra et vias publicas omni labore diminuenda, quia Pappus copiosus ejus, longe avolat, et cum spiritu combustus, suffocationis periculum minatur.

1414. *TYPHA media* foliis subsemicylindricis spica mascula femineaque rebus cylindricis.

Typha angustifolia foliis semicylindricis spica mascula femineaque remotis. *Sp. pl.* 2. p. 1377. *α.* — *Poll. fl.* p. 871. — *Lamarck et Decand. fl. fr.* p. 148.

Typha clava mascula a feminea remota. *fl. helv. n.* 1306. *var. α.*

Typha palustris clava gracili. *Bauh. pin.* *Mapp. fl. Alf.* p. 316.

Typha palustris media. *Bauh. hist.* 2. p. *Morif. hist.* 3. p. 246. *Sect.* 8. *t.* 13

Icon. Curtis. fl. Lond. t. 162. — *Oeder. Dan. t.* 815.

r m. Mittlere Rohrkolben.

ll. Massette mediaire.

b. Cum praecedente passim. Circa Carlsruhe in fossis aquaticis am Holzmagazin.

r. Julio, Augusto. Semina Septembri maturatione.

Differt a praecedente specie, *Foliis* subcylindricis canaliculatis angustioribus, *Spica* ciliore, mascula a feminea, septem lineas amentum remota. *Amentum* longe cylindricum.

f. Planta in horto botanico culta, constantissima.

15. TYPHA elliptica foliis linearibus canaliculatis, spica mascula femineaque remotis, feminea elliptica.

Typha minima. Lamarck et Decand. fl. franc. 3. p. 148.

Typha minor. Smith. fl. brit. 3. p. 960.

Typha angustifolia. var. β. Linn. Hall. stirp. helv. n. 1306. var. β.

Typha palustris minima duplici clava. Moris. hist. 3. Sect. 8. t. 13. f. 3. p. 246.

Typha palustris minor. Bauh. pin. 20. — Mapp. fl. Alf. p. 316.

Typha minor. Bauh. hist. 2. p. 540. Icon. Lobel. Ic. f. 81. t. 2.

Typha minor seu Typhula. Tabern. p. 559. (Icon optima).

r m. Elliptische kleine Rohrkolben.

ll. Massette elliptique.

b. In Insulis Rheni passim. In Principatu superiore ad Lacum Bodamicum, praeci-

pue inter *Roschach* et *Rheineck* abundanter vidi.

Flor. Junio, Julio. *Semina* Augusto maturantur.

Facile dignoscitur a praecedente, *Calyce* minoribus, gracilioribus, *Foliis* angustioribus brevioribus, glaucis, *Spica* feminea semine elliptica, multo brevior, unciam ad quinque lineas longa.

Obs. In horto botanico culta, mutationi non obnoxii cubuit.

CCCCXX. SPARGANIUM.

Linn Gen. plant. ed. Schreb. n. 1402.

Flores masculi numerosi, in capitulum collecti. *Amentum* commune subrotundum, demum imbricatum, constans perianthiis triphyllis, brevibus, deciduis. *Corolla* nulla. *Stamina* tria, capillaria calyce longiora. *Antherae* oblongae. *Feminei Calyx* ut in masculinis. *Receptaculum* commune subrotundum. *Corolla* nulla. *Germen* ovatum, desinens in stylum brevem, subulatum. *Stigmata* 1 2, acuminata, canaliculata, persistencia. *Drupa* exficcata, binata cum acumine, inferne angulata. *Nuxella* ossea, oblongo-ovata, angulata.

1416. SPARGANIUM *simplex* foliis triangularibus, lateribus planis, pediculis simplicibus. *Curtis. fl. Lond. N. 58.* — *Smith. fl. brit. 3. p. 961.* — *Willd. Linn. Sp. pl. 4. p. 199.* — *Lamarck Decand. fl. franc. 3. p. 149.*

MONOECIA. TRIANDRIA. 603

parganium simplex foliis ensiformibus, planis, caule simplici. *Huds. fl. Angl.* 2. p. 401.

parganium erectum β. non ramosum. *Linn. Sp. pl.* 2. p. 1378. — *Pollich. fl. pal.* n. 872. β.

parganium caule foliisque erectis. β. *Hall. stirp. helv.* n. 1303.

parganium non ramosum. *Bauh. pin.* 15. — *Mapp. fl. Als.* p. 294.

parganium alterum. *Tabernaemont.* p. 560. *Icon.*

n. *Curtis. fl. Lond.* t. 341. — *Oeder. fl. Dan.* t. 932. — *Schkuhr. bot. handb.* t. 282.

m. *Aufrechte einfache Igelkolbe.*

l. *Rubanier simple.*

b. Utrinque ad stagna, paludes, fossas aquaticas passim abunde, praesertim in Rhemi vicinia.

r. Junio, Julio. *Semina* Augusto, Septembri matura. ¶

Differt a sequente specie affini, *Caule* brevis, simplici, *Foliis* ensiformibus inferne tantriquetris, angustioribus, minus profunde liculatis. *Spicae* infimae femineae subpeculatae, superiores masculae, alternae, es.

In horto culta mutationi non succubuit.

7. *SPARGANIUM ramosum* foliis basi triangularibus, lateralibus concavis, pedunculis ramosis. *Curt. fl. Lond.* n. 58. — *Smith. fl. brit.* 3. p. 961. — *Lamarck et Decand. fl. franc.* 3. p. 149. — *Willd. Linn. Sp. pl.* 4. p. 199.

MONOECIA. TRIANDRIA.

Sparganium erectum foliis erectis triquetris.
Linn. Sp. pl. 4. p. 1378. — Poll. fl. pal. n. 872. — Hall. stirp. helv. n. 1303.

Sparganium ramosum. Bauh. pin. 15. — Mapp. fl. Alf. p. 294.

Sparganium l. Tabernæmont. p. 559. Icon. Curtis. fl. Lond. t. 342. — Schkuhr. handb. t. 282.

Germ. Aestige Igelskolbe.

Gall. Rubanier rameux.

Hab. Utrunque ad fossas aquaticas, stagna, paludes passim frequens.

Flor. Junio, Julio. Semina Augustio, Septembri matura. 7

Distinguitur a præcedente, *Caule* longioribus, *Foliis* longioribus, *latioribus*, *triquetris*, *Spicis* longe majoribus, *sessilibus*.

1418. *SPARGANIUM natans* foliis decumbentibus planis. *Linn. Sp. pl. 2. p. 1378. Poll. fl. pal. n. 873. — Lamarck et DeCand. fl. franc. 3. p. 150. — Smith. fl. brit. 3. p. 962.*

Sparganium foliis planis mollibus decumbentibus. Hall. stirp. helv. n. 1304.

Sparganium minimum. Mapp. fl. Alf. p. 294. Raj. hist. 1910. Angl. 3. p. 437.

Sparganium non ramosum minus. Dillenb. Giff. 130. Sp. 58.

Germ. Schwimmende Igelskolbe.

Gall. Rubanier flottant.

t. b. In fossis aquaticis argillaceis, et paludibus. Prope *Rintheim* rarius, ad paludes prope *Linkenheim*. Circa Argentinam in fodinis argillaceis in sylva prope *Ekbolsheim* post *Mappum* legi. In Principatu superiore ad Lacum Bodamicum. In Palatinatu in fossis, rivulis et piscinis circa *Lautern*, in aquis inter molam *imperialem* et dem *Hammer*. *t. Pollichio*.

or. Junio, Julio, Augusto. *Semina* Septembri, Octobri matura. **7**

Diffinguitur a praecedentibus, *Caule* tenebre, debili, *Folius* non erectis, nec triquetris, angustioribus, planiusculis, saepe in aquantibus, tunc ob aquae fluxum saepe praegis, *Spica* longe teneriore, *Spiculis* axillaris, sessilibus.

ota. Amat argillacea limosa.

Synonymon Caspari Bauhini ej. Prodr. 211. appo allegatum potius ad variationem pusillam simplicis pertinet.

CCCCXXI. Z E A.

Linn. Gen. pl. ed. Schreb. n. 1403.

Masculi flores in spicas distinctas laxas distincti. *Calyx* *Gluma* biflora, bivalvis. *Corolla* *ma* bivalvis: *valvulis* oblongis muticis longidine fere calycis: *exteriore* ventricosa ob; *interiore* apice bidentata. *Nectarium* dilium: *foliolis* carnosis, supra latioribus, truncis, vertice fulcatis, brevissimis. *Stamina* *menta* tria, capillaria. *Antherae* subprismatice, bifidae, apice dehiscentes. *Feminei* in

spica densissima infra masculos, foliis oblongis.
Calyx Gluma uniflora, bivalvis, persistens.
Corolla Gluma quadrivalvis: *valvulis* inaequalibus. *Stylus* filiformis omnium longissimus.
Stigma simplex, versus apicem sessile.
Receptaculum commune maximum, trigonum quinquangulare. *Semina* subrotunda, angulata, compressa, splendentia.

1419. *ZEAE Mays* culmo erecto, foliis linearibus, integris. *Linn. Sp. pl.* 2. p. 1378.

Frumentum indicum Mays dictum. *pin.* 25.

Frumentum indicum. *Tabernaem.* 1618. 619. *Icones.*

Icon. Regnault. bot. Blackw. t. 547. *Scot. bot. handb. t.* 283.

Germ. Mays. Welschkorn.

Gall. Froment de Turquie.

Hab. In America. Colitur passim praecipue in regionibus demissioribus sabulosis in abundantia.

Flor. Julio, Augusto. *Semina* Octobris. ☉

Planta speciosa nostratibus notissima pro frumento varietate, a longissimo retro tempore, in America culta, sub initium Seculi decimi sexti in Europam ducta, serius in Germania.

Multum Sacchari continet, et sine dubio pro nostris regionibus primaria; ex qua Saccharum possit.

Receptacula florum foemineorum tenella cum et aromate condita, suavissimum edulium cum comedendum, fiunt.

CCCC

MONOECIA. TRIANDRIA.

609

CCCCXXII. CAREX.

Linn. Gen. pl. ed. Schreb. n. 1407.

Masculi flores in spicam digesti. *Amentum* oblongum, imbricatum, constans *Squamis* unioris, lanceolatis, acutis, concavis, persistentibus. *Corolla* nulla. *Stamina* Filamenta tria caeca, erecta, calyce longiora. *Antherae* erectae, longae, lineares. *Feminei* flores in eadem vel in (quibusdam) in distincta planta. *Amentum* ut in masculis. *Corolla* nulla. *Nectarium* flatum, ovato oblongum, apice bidentatum, sperne contractum, ore dehiscens, persistens. *Semen* triquetrum intra nectarium. *Stylus* simplex. *Stigmata* tria vel duo, subulata, incurva, longa, acuminata, pubescentia. *Nectarium* majus factum semen fovet. *Semen* unicum, ovato-oblongum, triquetrum, altero angulo saepius minus.

Spica unica simplici, stigmatibus duobus.

20. *CAREX dioica* spica simplici dioica, capsulis striatis, marginibus hispidis reflexis. *Schkuhr. bot. handb. 3. p. 310.*

Carex dioica spica simplici dioica. *Linn. Sp. pl. 2. p. 1379.*

Carex spica simplici dioica, capsularum marginibus ferrulatis. Goodenough. Transact. Linn. Soc. 2. p. 139.

Carex spica unica terete, subrotunda, seminibus inflatis. Hall. stirp. helv. n. 1351.

Wiers Bad. Tom. III,

Q q

610 MONOECIA. TRIANDRIA.

Cyperoides parvum caulibus et foliis tenuis triangularibus, spica longiori, caulis oblongis, in angustum collum v. bifidum attenuatis. *Michx.* nov. gen. p. 56. t. 31. f. 1.

Icon. Schkuhr. bot. Handb. 3. t. 286. A. n. 1.

Ger. m. Getrennte Segge.

Gall. Carex dioica.

Hab. In pratis udis prope *Eckenstein*, *Knien*, *gen.*, *Schrök* et aliibi non infrequens. *Fr. Rastadt* ad *Favorite*, in pratis *auf dem Kienbrunnen*. In Principatu superiore cum *Bodanico* in pratis udis capiosa.

Flor. Majo, Junio. *Semine* Julio maturantur.

Radix repens. *Culmi* digitales, paleae pedales, erecti, triquetri, tenues, rigidiusculi, nudiusculi. *Folia* radicalia linearibulata, tenuia, culmi basin vestientia. *Sp.* dioica terminalis, simplicissima: *mascula*, ciliata longa, erecta, linearis, gracilis. *Glabra* pallide ex luteo-ferrugineae, margine membranaceae, ex albido-luteae. *Feminea* cylindrica, brevior, obtusa. *Capsulae* maturae patentes, demum reflexae, striatae, superne planiusculae, inferne convexae, rostratae, apice glandulosa, marginibus hispidulae, caeterum glabrae. *Stigmata* duo.

Obs. Rarius in basi *Spicae* masculae, flores 1-4 foeminei conspicui sunt.

1421. *CAREX pulicaris* spica simpliciter androgyna, superne mascula, capsula divaricatis, retroflexis. *Linn. Sp. pl. p. 1380.*

Carex spica unica terete, seminibus robustis. *Hall. stirp. helv. n. 1350.*

MONOECIA. TRIANDRIA. 611

Carex minima caulibus et foliis capillaceis, capitulo singulari tenuiori, capsulis oblongis, utrinque acuminatis et deorsum reflexis. *Michel. l. c. 66. t. 33. f. 1.*

n. Schkuhr. bot. handb. t. 286. A. n. 3. Leers. fl. Herb. t. 14. f. 1.

m. Floh. Segge.

l. Carex pucier.

o. In Marggraviatu superiore in valle dñe Mulde retro dem Boelchen t. Vulpio. In Principatu superiori ad Lacum Bodamicum in pascuis udis et pratis. In Sponhemia prope Kirchberg in pratis cum Satyrio viridi vidi.

r. Majo. Semina Junio, Julio matura. 7
Differt a praecedente, *Culmis* teretibus, semper androgyna, superne mascula, infeminea, *Capsulis* initio erectis, apprescenscentibus, demum patentibus, distichis, que retroflexis, utrinque longius acuminatis nigro rufescentibus, nitidis.

ta. Planta capsulis seminum maturis a quibusdam sub C. psyllophora salutatur. Sic Linneus in Suppl. 413.

Spica unica simplici, stigmatibus tribus.

2. CAREX pauciflora spica simplici androgyna, floribus foemineis subternis remotiusculis, patentibus: masculo subunico terminali. *Lightf. fl. Scot. 2. p. 543.*
— *Schkuhr. bot. handb. 3. p. 314. n. 4.*
— *Lamarck et Decand. fl. franc. 3. p. 102.*
Willd. Linn. Sp. pl. 4. p. 211.

Carex Leucoglochin spica simplici androgyna oblonga, superne mascula, capsulis reflexis, feminibus triquetris. *Linn.*

612 MONOECIA. TRIANDRIA

Suppl. 413. — *Ehrh. Beitr.* 1. p.
Roth. fl. germ. 2. p. 425.

Carex patula spica simplici andro
longa, superne mascula, capsul
gis acuminatis patulis. *Huds. fl.*
p. 402.

Icon. Schkuhr. l. c. t. 286. *A. n.* 4. —
l. c. t. 6. *f.* 2.


Germ. Wenigblumige Segge.

Gall. Carex pauciflore. Carex à quat.

Hab. In Brisgoviae montosis Sphag
tis prope *Triberg* cum *Eriophor*
nuper legi; In monte *Feldberg* ad
Feldsee dictum, cum *Scheuchzeria*
abunde vidi. In Principatu superio
Bohlingen in pascuis turfosis uligi
per vidi.

Flor. Majo. Semina Junio matura. ♀

Culmi digitates, palmares, trique
cti, tenues, nudiusculi. *Folia* radicalia
mos vestientia, eisque breviora, trique



Carex arenaria spica foliosa oblonga, acutiuscula, spiculis plurimis, terminalibus masculis, inferioribus foemineis, culmo incurvo. *Gooden. Act. Soc. Linn. Lond.* 2. p. 153.

Carex arenaria spica composita, spiculis androgynis; inferioribus remotioribus foliolo longiori instructis, culmo triquetro. *Linn. Sp. pl.* 2. p. 3381.

Carex spicis stipula brevioribus imis distinctis, superioribus congestis, *Hall. stirp. herb. n.* 1363.

Carex. Michel. nov. pl. gen. 67. n. 1, 2. t. 33. fig. 3. 4.

n. Schkuhr. l. c. t. 286, *B. n.* 6.

m. Sand-Segge. Deutsche Saffaparille.

l. Carex des sables.

h. Michelfeldae t. Casp. Bauhino.

Radix repens, cylindrica, articulata. Culpedalis, sesquipedalis et altior, erectus, modoque curvatus, triqueter versus radicem flexus. *Folia* linearia, canaliculata, acuta, rariuscula. *Squamae* foliaceae, versus radicem plurimae, fuscae. *Spica* terminalis, subcylindrica, densa, oblongo-ovata, composita. *Glumae* androgynae, sessiles, adpressae, summae saepe mere masculae, *Bracteae* foliaceae, ovatae, membranaceae, spiculis longiores. *Glumulae* marginatae, membranaceae, *Stigmata* duo.

Nota. Veram plantam hic descripsi. Plantam ab Hallero Michelfeldae citatam non vidi.

Alias copiose in sabulosis maritimis, nec non in arena mobili crescit Frustra eam hucusque in terris Badensibus quaesivi.

rm. CARICIS ARENARIAE Radix.

614 MONOECIA. TRIANDRIA.

Vires resolventes habet et radici Sarsap
quibusdam in locis substituitur.

1424. *CAREX intermedia* spica
ga obtusa, spiculis plurimis: infi
minalique femineis; intermediis
lis; culmo erecto. *Gooden. l. c. 8*
l. c. p. 320. — *Willd. Linn. Sp*
p. 224.

Carex spicata spica composita, spi
subrotundo - ovatis numerosis c
in summo scapo cohaerentibus.
bus androgynis nudis. *Poll. fl. pa*

Carex disticha spica composita sub
spiculis ovatis imbricatis androg
lio longiori instructis, culmo tr
Huds. fl. Angl. 2. p. 347. — *Schr*
fl. Lipsf. p. 63. — *Roth. fl. ger*
426. — *Lamarck et Decand. fl. j*
p. 104.

Carex arenaria. Leers. fl. Herb. n.



ueter a basi ad medium foliosus. *Folia* catta, glabra, margine et carina retrorsum ra. *Spica* terminalis subbiuncialis, compo-
Spiculis alternis 12-20 subdistichis, subimatis, sessilibus, inferioribus saepe remotioribus. *Spicula* infima interdum reliquis magis otata, saepius *Bractea* longa acuta foliacea otata. *Glumae* ferrugineae, margine albidae. *Sulcae* acutae, marginatae, mucronato-ciliatae. *Stigmata* duo, plumosa.

Spiculae infimae et superiores saepe fere mere feminae, intermediae masculinae.

otata. Radices praecipue in regionibus rhenanis saepe Radicibus Caricis arenariae substituantur.

15. *CAREX vulpina* spica supra decomposita, coarctato-ramosa, obtusa, spiculis superne masculis, capsulis divergentibus, culmo angulis acutissimis. *Gooden. l. c. Schkuhr. l. c. p. 321.*

Carex vulpina spica supradecomposita inferne laxiore, spiculis androgynis ovatis sessilibus glomeratis superne masculis. *Linn. Sp. pl. 2. p. 1382. — Pollich. fl. pal. n. 876.*

Carex spicis subrotundis echinatis, supremis confluentibus, capsulis rostratis bifidis. *Hall. stirp. helv. n. 1364.*

Carex palustris major radice fibrosa, caule exquisitè triangulari, spica brevi habitiori compacta. *Michel. l. c. 69. t. 33. f. 13. 14.*

Carex major spica compacta. *Mapp. fl. Als. p. 55.*

Icon. Schkuhr. l. c. t. 286. C. n. 10. — fl. Herb. t. 14. f. 5. — Oeder, fl. t. 308.

Germ. Fuchs-Segge.

Gall. Carex compact.

Hab. Utrunque in pratis udis, humidis aquarum fossas.

Flor. Majo, Junio. *Semina* Junio, Julio. *tura.* ☞

Radix dense fibrosa. *Culmi* sesquipedales et altiores, erecti, inferne crassi, exacte triquetri, angulis praeserpe scabriusculi. *Folia* latiuscula, ca. margine et carina aspera. *Spica* terminalis posita, ovato-oblonga, compacta, fasciam, ad binas uncias longa. *Spiculae* imbricatae, glomeratae, femineae, apiceae. *Bractae* ovato-setaceae, longitudinalium. *Capsulae* ovatae acutae, e ferrugineae, glumis longiores, subdivergentes. *Stylus* duo plumosa.

1426. CAREX *divulsa* spica deo-
elungata basi subramosa, spiculis
ribus remotis; summis contiguis
fulis suberectis. *Gooden. l. c. Sc.*
l. o. p. 324. Willd. Linn. Sp. pl. 4.
Lamarch et Decand. fl. franc. 3.

Carex canescens, Hudf. fl. Angl. 2. p.

Carex minor spica divulsa seu inte-
Mapp. fl. Alf. p. 56.

Carex nemorosa fibrosa radice, ca-
quifite triangulari, spica longa
seu interrupta, capitulis solitariis

MONOECIA. TRIANDRIA. 617

terquam ultimo. *Michel. l. c. 69. t. 33.*
fig. 10.

n. Schkuhr. l. c. p. 324. t. 286. D d. n. 89.
et W w. n. 89.

m. Unterbrochenährige Segge.

ll. Carex écarté.

i. In nemoribus graminosis humidis passim. Circa Muhlbürg im Lutherschen Wäldlein et alibi. In Brilgovia am Kaiserstuhl in nemoribus humidis. In Principatu superiore prope Mörsburg, Salem, Ueberlingen et alibi in nemoribus frequens. In Alsatia prope Argentinam in sylva Ostwald et alibi.

r. Majo, Junio. Semina Junio, Julio matura. 7

Radix fibrosa. Culmi pedales, cubitales, iusculi, debiles, acute triquetri, angulis iusculi. Folia culmo longiora, linearia, carinata, carina et margine aspera, saepe viridia. Spica terminalis, elongata, anthera, basi quandoque subramosa. Spiculae flores remotae, ovatae, sessiles, Bracteis eis brevibus hispidis suffultae; superiores non approximatae, contiguae, floribus masculinis terminalibus. Glumae ovatae, mucronatae, albae, dorso virides, capsula longiores. Glumulae ovatae, acutae, apice bifidae, hinc et illinc convexiusculae, suberectae. Stigmata duo.

7. *CAREX muricata* spica oblonga subdecomposita, spiculis distinctis, capsulis divergentibus, ore fisso, radice fibrosa. *Gooden. l. c. — Schkuhr. l. c. p. 324. — Willd. Linn. Sp. pl. 4. p. 234. — Roth. fl. germ. 2. p. 439.*

618 MONOECIA. TRIANDRIA

Carex intermedia. Retz. fl. Scand.

Carex canescens. Leers. fl. Herb. p

Carex spicata spiculis subrotundis
nis contiguis, capsulis ovatis acu
fl. angl. 2. p. 405.

Carex caule foliisque asperis, spici
tis subrotundis, supremis congel
stirp. helv. n. 1365.

Carex palustris media, radice fibro
exquisite triangulari, spica br
pactiori. *Michel. l. c. 69. t. 33.*

Gramen cyperoides minus, spicis
minus compactis. *Scheuchz. Aq*
488. t. 11. f. 5.

Icon. Schkuhr. l. c. t. 186. E. n. 22. -
l. c. t. 14. f. 3.

Germ. Stechende Segge.

Gall. Carex rude.

Hab. In pratis humidiusculis et sylvis
sis passim. ♀



Primo intuitu affinis *C. stellulatae* eaque saepe confunditur, a qua tamen differt, Spiculis interne non masculis, Capsulis nudo oculo observatis simpliciter mucronatis, sed apice manifeste bifidis.

Nota. *Carex canescens* Poll. fl. pal. n. 881. f. *C. loliacea*? Schkuhr. l. c. p. 326. t. E. e. n. 91. est sine dubio mera variatio, spiculis minoribus, Caricis muricatae.

428. *CAREX teretiuscula* spica supra decomposita coarctato-ramosa acutiuscula, spiculis glomeratis, capsulis patentibus, culmo teretiusculo. *Gooden. l. c.* — *Schkuhr. l. c. p. 334.*

Carex teretiuscula spicis androgynis coarctato-paniculatis, superne masculis, stigmatibus binis, fructibus ovatis bidentatis margine ciliato-ferratis, culmo inferne tereti, superne obtuse triquetro. *Willd. Linn. Sp. pl. 4. p. 244.*

Carex diandra. Schrank. fl. Bavar. p. 281. Hoppe Taschenb. 1797. p. 98.

non. Schkuhr. l. c. t. 286. D. T. n. 19. 69.

erm. Rundhalmige Segge.

all. Carex à tige ronde.

a b. In pascuis et pratis uliginosis passim frequens. In Principatu superiore prope *Härtwangen*, *Bohlingen* in turfosis et alibi ad Lacum Bodamicum passim copiose.

lor. Majo. Semina Junio matura. 7


Radix fibrosa. *Culmi* pedales, sesquipedales et altiores, erecti, curvuli, inferne teretiusculi, superne triquetri, striati, aphylli, versus radicem foliosi. *Folia* carinata, canaliculata, angusta margine et carina aspera culmis saepe breviora, lacte viridia. *Spica* terminalis,

ovato-oblonga, acutiuscula, unciam circiter
 ga, supradecomposita, densa. *Spiculae* num-
 ovatae acutae, sessiles. *Flores* masculi fu-
 res plures, *foeminei* inferiores pauciores
mae acutae, fuscae, splendentes, margini
 de-membranaceae. *Capsulae* ovatae acutae
 planiusculae, illinc convexae, gibbulae,
 ne serrato-hispidae, patentes. *Bractea* sul-
 brevissima, basi membranacea, dilatata,
 flam tenuissimam terminata. *Filamenta* tri-
therae tres, lineares, flavae. *Stigmata* d

Nota, Differt a variatione minore *Caricis* p-
 latae. Culmis tenuioribus, teretiusculis
 argute triquetris. Foliis multo angustis
 Spica densiore, saepe brevior. Bractea
 viore, Capsulis supra non dimidio majoribus
 membranaceis.

Obs. Nostra individua simillima sunt lectis in
 ab amico mihi sub *Carice* diandra communi

1429. *CAREX paniculata* spica su-
 compolita paniculato-ramosa, ra-
 ternis remotiusculis capsulis paten-



MONOECIA. TRIANDRIA. 621

Cyperoides palustre, spica laxa minus.
Scheuchz. Agrost. 501. — Hall. stirp. helv. var. β. — Pollich. fl. pal. n. 882. β. Leers. fl. Herb. n. 713. β.

l. *Schkuhr. l. c. t. 286. D. n. 20. Leers. l. t. 14. f. 4.*

m. *Rispenförmige Segge.*

l. *Carex paniculé.*

In pratis pascuisque udis paludosis cis et transrhenanis passim cum var. β.

l. Majo, Junio. *Semina* Julio matura. 7

Radix repens, fibrosa, fusca. *Culmi* ses-
 dales bipedales et altiores, erecti, a basi
 edium fere foliati, exacte triquetri, an-
 scabri. *Folia* culmo saepe breviora, quan-
 e longiora, carinata, canaliculata, 2-3
 lata, margine et carina retrorsum scabra.

terminalis paniculato-ramosa, 3-4 un-
 . *Ramuli* alterni, remotiusculi, spiculis
 que fere obtecti. *Spiculae* subovatae, fru-
 ae glomeratae. *Bractae* setaceae. *Glumae*
 -acuminatae, pallide fuscae, margine al-
 . *Capsulae* subventricosae, patentes, supra
 lio marginatae albide membranaceae, spi-
 utae, subbifidae. *Stigmata* duo, plumosa.

io minor, Spica simpliciori vix ramosa, minore,
 istructa est, caeterum quod fructificationis par-
 s simillima majori.

ta. *Schkuhr. l. c. t. 286. D. n. 20.*
 fig c. non male exprimit variationem minorem.

l. *Spicis androgynis apice femineis; Stig-
 matibus duobus.*

o. *CAREX cyperoides* spica compo-
 sita capitata, spiculis densissimis, invo-
 lucro longo. *Schkuhr. l. c. p. 342.*

MONOECIA TRIANDRIA.

Carex cyperoides capitulo terminali
tanto, floribus simplicissimis sul-
involucro longo. *Linn. Syst. veg.*
Suppl. p. 413.

Carex bohemica spica globosa conglom-
androgyna, capsulis subulatis, involu-
triphylo. *Schreb. Graef.*

Carex bohemica aquatica annua, cy-
caule molli, capitulis in glome-
spicam digestis. *Michel. gen. 70.*
Fig. 19.

Schreb. Graef. 2. l. 28. f. 2. — Schreb. l. c. t. 286. A. n. 5.

Germ. Cyperngrasartige Segge.

Gall. Carex faux Souchet.

Hab. Circa Carlsruh prope Scheibenhart
Rumice maritimo in prato udo leg
1785. et 1787.

Flor. Junio. *Semina* Julio matura. 7

Radix fibrosa. *Culmus* palmaris, pe-
erectus, triqueter, glaber, versus radice
liatus, caeterum aphyllus. *Folia* culmo
longiora, dorso carinata, retrorsum scabra
capitulum terminale, ovali-rotundum, mu-
rum, viride-luteolum, densum. *Bractea*
capitulo tres ad quatuor inaequales, foli-
longae, erectae, extrema reliquis longiore.
res in capitulo 30-50, superiores femineae
inferiores masculi. *Filamenta* tria longa. *A-*
rae tres lineari-lanceolatae, luteae. *Gla-*
lanceolatae, angustae, acuminatae, mer-
paeae, albiae. *Capsulae* ovali lanceol-
apice bifidae, acutae. *Stigmata* duo.

MONOECIA. TRIANDRIA. 609

f. *Rarissimum* Florae nostrae civem, quem ante viginti annos in loco allegato legi, ab eo tempore non amplius videre mihi contigit.

Planta in horto culta, mutationi non succubuit.

31. CAREX *leporina* spica composita, spiculis ovatis sessilibus approximatis alternis androgynis nudis. *Linn. Sp. pl. 2. p. 1381. — Poll. fl. pal. n. 878. — Leers. fl. Herb. n. 707.*

Carex ovalis spiculis subsenis ovalibus, approximatis alternis, squamis lanceolatis capsulam aequantibus, capsulis marginatis membranaceis. *Goodenough. l. c. Schkuhr. l. c. p. 343. — Willd. Linn. Sp. pl. 4. p. 229.*

Carex spicis nudis subrotundis, distinctis, in summo caule congestis. *Hall. stirp. helv. n. 1361.*

Gramen cyperoides, spica e pluribus spicis mollibus composita. *Scheuchz. Agrostr. 456. t. 10. f. 15.*

o n. Schkuhr. l. c. t. 286. B. n. 8. — Oeder. fl. Dan. t. 1115. — Leers. l. c. t. 14. 6.

r m. Hasen-Segge.

a l l. Carex nu.

a b. Utrunque in pratis et pascuis paludosis passim frequens.

o r. Majo, Junio. Semina Junio, Julio matūra. 7

Radix dense fibrosa. *Culmi* pedales sesquiduales et altiores; erecti, trigoni, striati, laeviusculi, versus radicem foliati, infra spicam rorsum scabriusculi. *Folia* linearia, uninerviata,

angusta, margine et carina scabra, culm
breviora. *Spica* terminalis, composita
lis 5-7 alternis, sessilibus, distinctis,
apice foemineis, basi masculis, mollis
ctiferis ovali-subrotundis, fuscis seu ferr
Bractea sub spicula infima ovato-oblong
terminata, longitudine spiculae, quam
multo longior. *Glumae* ovato-lanceolatae
fae, pallide ferrugineae. *Capsulae* utrinque
ginatae, membranaceae, apice bifidae. *S
ta* duo, plumosa.

Variat. Spiculis tribus ad novem.

Nota. Nomen a Linneo inditum specificum
optimum est et expressivum, non mutandum
omnino sub *C. leporina* nostram plantam
ex nota adjecta et Synonymis allegatis
patet, comprehendebat. Planta nostra in
vulgaris. Ad haec perillustres Praefes debent
hanc speciem in excursionibus botanicam
viam, hoc nomine semper salutabat.

432. CAREX Schreberi spica composita, nudiuscula, spiculis contiguis, capsulis margine membranaceis, squamis lanceolatis aequalibus. *Schuhr, f. c. p. 344.*

Carex Schreberi spiculis basi masculis confertis quinis, squamis aequalibus capsulis ovali-ovatis scutiusculis vix planis, subacutangulis, margine obtusiusculis ore emarginato, foliis angustissimis, radice repente. *Waldst. Act. Hort. bot. 1803. p. 145.*

Carex praecox spica composita distichodiuscula, spiculis contiguis androteretibus rectis, culmo superne. *Schreb. Spic. fl. Lips. 63. — Roem. germ. 2. p. 428.*

Carex spicis ternis congestis, glumis aristatis. *Hall. stirp. helv. n. 1367.*

n. Schkuhr. l. c. t. 286. B. n. 9.

m. Schreberische Segge.

l. Carex de Mr. Schreber.

i. In graminosis siccis apricis sabulosis elevatis asperis prope *Mühlburg auf den alten Schanzen*, et alibi passim. In Margraviatu superiore in monte prope *Krenzach*, nec non in sepibus gramineis circa Basileam, et ad viam versus *Riechen* ducentem, ad sepes.

r. Aprili, Majo. *Semina* Junio matura. *z*

Radix repens, articulata, fusca, fibrillosa, ni ex radice plurimi, palmares, pedales et res, erectiusculi, debiles, graciles, tri-ri, saepe curvati, versus radicem solicaeterum aphylli. *Folia* linearia, carina-perangusta, culmo saepe breviora. *Spica* ncialis composita *spiculis* alternis sessilibus, contiguis, androgynis, teretibus, utrinque is, rectis, rufo-fuscis; fructiferis ovali-acu-*Bractea* sub spicula ovali-setacea, spicula ior, basi fusca. *Glumae* ovali-lanceolatae, e. *Capulae* ovatae, convexo-planae, non branaceo-marginatae, apice acutae, bi-*Stigmata* duo, in *Stylo* longo, plumosa.

A praecedente facile distinguitur, Loco natali, Radice repente, Culmis debilioribus saepe vis, Foliolis multo angustioribus. Spiculis nica florente angustioribus, teretiusculis, utrinque s, Glumis fuscis, Capulis non membrana-marginatis.

Individua *Halleriana* circa Basileam lecta, saepissime tantum spiculis in spica tribus instructa sunt, caeterum simillima aliis, in variis germaniae regionibus lectis.

1433. *CAREX brizoides* spica comp
subdisticha nuda; spiculis oblongis
tiguīs patulis curvatis, culmo n
Schkuhr. l. c. p. 346.

Carex brizoides spica composita, disticha
da: spiculis androgynis oblongis et
guis, culmo nudo. *Linn. Sp. pl. 2p. 1*
Roth. fl. germ. 2. p. 430.

Carex brizoides spiculis basi masculis nu
rolis contiguīs sublanceolatis suber
vis, capsulis ovato-lanceolatis convexis,
fimo-subconcaviusculis attenuatis, marginibus
serrulatis ore bifido. *Wahlb. act. Holm. 1803. p. 145.*

Carex spicis distichis, teretibus, repant
Hall. stirp. helv. n. 1358.

Carex fibrata radice, angustifolia, caule erecto
site triangulari. *Michel. gen. 70. 1*
fig. 17.

Icon. Schkuhr. l. c. t. 286. C. U. n. 12.
Germ. Zittergrasartige Segge.

ilis subdistichis 5-8-10, sessilibus alternis, retiusculis, compressiusculis, inferne masculis, superne femineis, erecto-patentibus sæpe recurvis. *Bractea* sub spiculis, brevis, setacea. *Glumae* ovato seu subcordato-lanceolatae, ex albido vel luteo-virescentes. *Capitula* ovato-lanceolatae, margine membranaceae, apice bifidae, fuscae. *Stigmata* duo plures. *Antherae* lineares, longae, pallide luteae.

234. CAREX canescens spiculis subrotundis remotis sessilibus obtusis androgynis, capsulis ovatis acutiusculis. *Linn. Sp. pl.* 2. p. 1383.

Carex canescens spiculis basi masculis, subapproximatis quinis, squamis subaequantibus, capsulis subrotundo-ovatis acutiusculis convexo-subconvexiusculis, subobtusangulis, ore bidentato. *Wahlenb. Act. Holm.* 1803. p. 147.

Carex curta spiculis subsenis remotiusculis, culmis nudis, squamis ovatis acutiusculis capsula brevioribus. *Gooden. l. c. p.* 145. — *Schkuhr. l. c. p.* 347. — *Willd. Linn. Sp. pl.* 4. p. 241.

Carex cinerea spiculis oblongo-ovatis sessilibus, remotis androgynis, capsulis ex subrotundo-ovatis, convergentibus. *Poll. fl. pal. n.* 880.

Carex spicis ovatis alternis, supremis se contingentibus. Hall. stirp. helv. n. 1360.

Gramen cyperioides spicis curtis divulsis. Loefel. fl. Pruss. p. 117. t. 32.

non Schkuhr. l. c. t. 286. *C. n.* 13.

Ger m. Graulichte Segge.

Gall. Carex gris.

Hab. Utrunque in pascuis udis, aquosis, humidis cis et transrhenanis, necnon in prae-sylvaticis montosis uliginosis passim frequens.

Flo r. Majo. *Semina* Junio matura. 7

Radix fibrosa. *Culmi* pedales, sesquipedales, erecti, triquetri, versus radicem foliis angulis retrorsum scabri. *Folia* linearia, canaliculata, vix canaliculata, culmo saepe breviora, carina et margine aspera. *Spica* terminalis, compacta unciam ad sesquiunciam vix longa, composita *spiculis* alternis, sessilibus, rarius tribus saepe fine 4-5-6 ad 7 subremotiusculis, florentibus ovali-teretiusculis, fructiferis ovalibus, oblongis brevibus, exviridi canescentibus, erectis, lineas vix longis. *Bractea* infima, reliquis brevior, setacea, spicula saepissime brevior, quoadque longior. *Glumae* ovatae, acutae, cinerascinae, dorso viridulae. *Capsulae* ovatae, saepe subbifidae. *Stigmata* duo, plumosa.

MONOECIA. TRIANDRIA. 629

fulis ovali-conicis attenuatis convexo-planis acutangulis divaricatis. *Wahlenb. Act. Holm.* 1803. p. 147. — *Willd. Linn. Sp. pl.* 2. p. 236.

Carex stellulata spiculis subternis remotis, capsulis divergentibus acutis, ore integro. *Gooden. l. c.* p. 144. — *Schkuhr, l. c.* p. 349.

Carex echinata spiculis subovatis sessilibus remotis androgynis inferne masculis, superne femineis, capsulis simpliciter mucronatis. *Hoffm. fl. germ.* 326. — *Roth. fl. germ.* 2. p. 431.

Carex muricata. *Poll. fl. pal. n.* 877. — *Leers. fl. Herb. n.* 709. — *Hudsf. fl. angl.* 2. p. 406.

Carex spicis ternis echinatis, glumis lanceolatis, capsulae mucrone simplici, *Hall. stirp. helv. n.* 1366.

Stemmen cyperoides spicatum minimum, spica divulsa aculeata. *Scheuchz. Agrost.* p. 485. t. 11. f. 3.

2. *Schkuhr. l. c. t.* 286. *C. n.* 14. — *Leers. l. c. t.* 14. f. 8. — *Oeder. fl. Dan. t.* 284, m. *Sternförmige Segge.*

1. *Carex étoilée.*

Utrunque in pratis et pascuis udis, paludosis, ad piscinas passim frequens.

r. Majo, Junio. *Semina* Junio, Julio matura. ☞


Radix fibrosa, subrepens. *Culmi* ex radicibus, palmares, semipedales, pedales, et alii, erecti, obtuse triquetri, versus radicem

foliosi. *Folia* canaliculata, carinata, culmo saepe breviora. *Spica* terminalis posita *Spiculis* alternis tribus, saepius tuor ad quinque, sessilibus, remotiusculoque subapproximatis, androgynis, femineis, inferne masculis, brevibus fatis ovali subrotundis. *Bractae* f spiculis breviores, infima saepe longissimae foemineae ovato-acuminatae, vix tes, margine albae. *Capsulae* ovato-natae, apice mucronatae, divergentemata duo, plumosa.

Nota. Capsulae quammaxime divergentes stellulatas repraesentant, apice nudo c pliciter mucronatae, per lentem spec bifidae.

1436. *CAREX remota* spiculis ax solitariis remotis subsessilibus, longissimis, capsulis apice indivi den. l. c. p. 150. — Schkuhr. l. c.

Carex remota spicis ovatis subsessi motis androgynis, bracteis cul



m. Entfernte Segge.

ll. Carex espace.

b. In sylvis et nemoribus humidis ad vias, nec non in pratis paludosis inter salices et alnos passim frequens.

r. Majo, Junio. *Semina* Julio matura. γ

Radix fibrosa. *Culmi* pedales bipedales et al-
s, erecti, debiles, tenues, triquetri, angulis
orsum scabri, foliati. *Folia* angusta, subca-
a, longitudine culmi, et longiora, ex lu-
viridia, scabriuscula. *Spiculae* sessilibus;
ariae, in axillis foliorum, maxime remo-
androdynae, ovaes, inferne masculae,
ne femineae. *Bracteae* (seu folia floralia)
ae foliaceae, longissimae, culmum supe-
s, reliquae superiores sensim breviores,
stiores. *Glumae* lanceolatae, acutae, al-
ites, dorso viridulae. *Capsulae* pallide fla-
ites, ovato-lanceolatae, erectae, acutae,
oculo apice integrae, per lentem specta-
bifidae. *Stigmata* duo.

ile *Spiculis* axillaribus solitariis remotis, *Brac-*
ue foliaceis longissimis a reliquis speciebus di-
statur.

ta. Linneus olim hanc speciem etiam sub *C.*
axillari comprehendebat. Ill. Goodenough
a. c. autem sub axillari comprehendit novam
speciem, in Anglia obvenientem, spiculis subter-
natis remotis axillaribus, capsulis apice divisis
preditam.

7. CAREX *elongata* spiculis oblon-
gis sessilibus remotis androgynis, capsu-
lis ovatis acutis. *Linn. Sp. pl.* 2. p. 1383.
Poll. fl. pal. n. 879. omnino! (*excluso*
Morisoni synonymo.)

Carex spicis ovatis imis remotis, supremis congestis. Hall. stirp. helv. n. 1359. (excluso Morisoni synonymo.)

Carex multiculmis Ehrh. gram. n. 88. Hoffm. fl. germ. p. 328.

Cyperoides polystachyon spicis laxis paniculam sere componentibus. Scheuchz. Agrostr. p. 487. t. 11. fl. 4.

Icon. Schkuhr. l. c. t. 286. E. n. 25, Leers. Herb. t. 14. f. 7.

Germ. Verlängerte Segge.

Gall. Carex alongé.

Hab. In pratis humidis, paludibus et sylvis undis ad fossas et rivulos passim non inloquens.

Flor. Majo, Junio. *Semina* Julio matura. 7

Radix repens, fibrillosa. *Culmi* ex radicibus plurimi, congesti, pedales, cubitales, erecti, triquetri, angulis retrorsum scabri, versus teretem foliati, caeterum aphylli. *Folia* carinata, linearia, carina et margine retrorsum scabra, laete viridia, culmo saepe longiora. *Spica* terminalis 2-3 uncialis, erecta, composita *Spicula* alternis, sessilibus, remotiusculis 6-7-9-11, ovali cylindricis, acutiusculis, quinque ad sex lineas longis, erectis, inferne masculis superne femineis. *Bractae* setaceae, brevissimae. *Glaumae* ovatae, acutae, obtusaeque, initio viridulae, margine fuscae, demum fuscae, margine albidulae, capsula breviores. *Capsulae* ovato-acutae, erecto patentes, nervosae, subexcurvae, apice nudo oculo integrae, per lentem subbifidat. *Stigmata* duo.

Obs. Specimina in regionibus rhenanis observata, semper spiculis septem ad novem praedita sunt, rarius undecim.

MONOECIA. TRIANDRIA. 639

Individuum a Leersio exhibitum egregie nostram
lantam, a C. canescente, diversam, sistit.

ta. In C. canescente, Spiculæ sæpissime
tantum quatuor, semper multo breviores.

††† *Spicis sexu distinctis; Stigmatibus
duobus.*

8. *CAREX caespitosa* digyna, vagi-
nis nullis, spicis subseffilibus subcylind-
ricis obtusis, foliis erectis molliusculis.
Gooden. l. c. — Schkuhr. l. c. p. 361.

Carex spicis erectis cylindricis ternis sub-
seffilibus, mascula terminali, culmo tri-
quetro. *Linn. Sp. pl. 2. p. 1388. — Roth.
fl. germ. 2. p. 452.*

Carex spicis masculis duabus, femininis seffi-
libus, capsulis ovatis brevissime mucro-
natis. *Hall. stirp. helv. n. 1400. (videtur.)*


n. *Schkuhr. l. c. t. 286. Aa. Bb. n. 85. a-e.*
m. *Rafen-Segge.*

l. *Carex en gazon.*

o. In pascuis pratisque udis turfosis passim
frequens. In Principatu superiore in tur-
fosis ad Lacum Bodamicum copiose.

r. Majo, Junio, *Semina* Junio, Julio, ma-
tura. ☞

Radix cylindrica, repens, fibrillosa. Cul-
mares, pedales et longiores, erecti, tri-
ri, angulis retrorsum scabri, inferne fo-
caeterum aphylli. *Folia* linearia, cana-
ta, carinata, erecta, ex viridi-glaucula,
iuscula, carina et margine retrorsum sca-
cula, culmum aequantia. *Spicae* saepe tres,



fusculae, subcompressae, glabrae.
grae. *Stigmata* duo.

Nota. Pro diverso solo natali maxime
quoad spicarum numerum ludit. Spic
neæ et suprema mascula unica. A
quatuor quinque femineæ cum spic
quandoque altera accessoria minor
mineis quibusdam saltem aut omnib
cem masculis Spica saltem femine
unica.

1439. *CAREX gracilis* spicis fi
femineis inferioribus pedunc
ctis, bracteis longe foliaceis
bus binis, capsulis ovatis api
ore pertulis.

Carex gracilis spicis masculis
pluribus, subfiliformibus, flo
nis. *Curt. fl. Lond.*

Carex acuta spicis filiformibus, f
florescentibus suberectis, fru

Carex spicis maribus ovatis, femininis gracilibus erectis, capsula triquetra-conicis. Hall. stirp. helv. n. 1406.

n. Curtis. fl. Lond. t. 282. — Schkuhr. l. c. t. 286. E c. F. f. n. 92. a, b. — Leers. fl. Herb. t. 16. f. 1.

r m. Hagere Sumpf-Segge.

ll. Carex grêle.

b. Utrique in pratis paludosis, ad aquas stagnantes, piscinarum marginalia et ad fossas aquaticas passim abunde.

r. Aprili, Majo. Semina Junio, Julio maturata. ♀

Radix cylindrica, articulata, repens, filiosa. Culmi pedales, cubitales, tripedales altiores, erecti, foliosi, triquetri, angulis tri. Folia linearia, carinata, longa, viridia, carina et margine aspera. Spicae masculae superiores saepissime tres, graciles, e fusco nigris, subnutantes, femineae inferiores 3-4-5 sessiles vel pedunculatae, teretes, graciles, triangulares et longiores, suberecti, nigris. Glumae sub spicis femineis praelongae, foliaceae. Glumae ovato-acutae, aut obtusiusculae, e imbricatae, carinatae, fusco-nigris, viridi. Capsulae ovatae, acutiusculae, e indivisae, ore pertusae. Stigmata duo, longa, plumosa.

nota. Facile a Carice acuta riparia, et pendula Stigmatibus semper duobus, nec tribus distinguitur.

diat. Spicis femineis superioribus quandoque superne masculis.

++++ Spicis sexu distinctis, mascula unica, femineis sessilibus, seu breve peduncula-

tis, vaginis bracteatis brevibus, vagin
vel nulla; STIGMATIBUS tribus.

1440. *CAREX flava* spicis confertis sub
sessilibus subrotundis, mascula lineari
capsulis acutis recurvis. *Linn. Sp. pl.*
p. 1384. — *Poll. fl. pal. n.* 883. — *Leers*
fl. Herb. n. 714.

Carex flava vaginis brevibus subaequantibus
foliolo divaricato, spica mascula lineari
femineis subrotundis, capsulis rostrato
acuminatis. *Gooden. l. c.* — *Schkuhr. l. c.*
p. 376.

Carex flava spicis subpedunculatis ovalibus
bracteis vaginatis longe foliatis subor
proximatis, capsulis obovato-cuneatis
mibus inflatis costatis, rostro longo e
ticulari deflexo, ore bifido. *Wahlb.*
Act. Holm. 1803. *p.* 150.

Carex spicis foemineis petiolatis erectis ore
tis, echinatis. *Halt. stirp. helv. n.* 130.

β. *Carex Oederi* spicis lateralibus confertis sub
pedunculatis ovato-subrotundis andro
dynis; terminali mascula lineari, capsu
lis acumine recto terminatis. *Roth. fl.*
germ. 2. p. 438. — *Schrad. Spic. fl. germ.*
p. 51.

γ. *Carex Oederi* spicis confertis subsessilibus sub
rotundis, mascula lineari, capsulis glo
bosis rectis acuminatis. *Willd. Act. Acad.*
Berol. 1794. *p.* 44. *t.* 1. *f.* 2. — *Schkuhr.*
l. c. p. 371.

Icon. Schkuhr. l. c. t. 286. *H. n.* 36. *γ. tab.* 2
n. 26. — *Leers. fl. Herb. t.* 15. *f.* 6.

Gram. Salic. Sagge.

11. *Carex jaune.*

a b. Utrinque in pascuis arenosis udis, inundatis, passim frequens. γ. in pascuis macilentis sabulosis inundatis et ad rivulorum margines. Abunde in Principatu superiore ad Lacum Bodamicum, in turfosis.

lor. Majo, Junio. *Semina* Junio, Julio maturatione. ♀

Radix subrepens, fibrosa. *Culmi* digitales, imares, pedales erecti, quandoque decumbentes, triquetri, foliati, saepe exceptis bracteis nudi, versus radicem foliati. *Folia* culmo saepe breviora, linearia, acuta, latiuscula, subrinata, margine scabra. *Spica* terminalis mascula lineari-cylindrica, gracilis; *femineae* spicae 1-2-3 subseffiles, quandoque pedunculatae, approximatae, saepe etiam maxime remotae, subrotundae. *Bracteae* foliaceae, vaginantes, culmo saepe longiores. *Glumae* ovali lanolatae, mucronatae, pallide ferrugineae, dorso virides, capsulis breviores. *Capsulae* obovato-cuneiformes, inflatae, costatae, flavicantes, stylo longo deflexo apice bifido. *Stigmata* 2.

Nota. Pro diverso solo natali maxime polymorpha, Culmis demissioribus et longioribus, Foliis latoribus et angustioribus, brevioribus et longioribus, Bracteis erectis. patentibus et reflexis, Spicis femineis saltem unica aut 2-3-4 maxime approximatis subseffilibus, aut remotis pedunculatis, omnibus mere femineis. aut etiam sed id tamen rarius, superne masculis. Capsulis rostro reflexis, rectisque saepe in unico eodemque Individuo; in aliis omnibus reflexis, aut rectis.

Doleo Cl. Hostii gramina, opus splendidum egregium, quod citari oportebat, mihi non esse ad manus.

1441. *CAREX praecox* vaginis brevibus subæquantibus spicis subapproximatis longo-ovatis, capsulis ovatis pubescentibus, apice integris. *Schkuhr. l. c. p. 3.*

Carex praecox vaginis brevibus subæquantibus, spicis approximatis, mascula clavata, femineis ovatis, capsulis subtundis pubescentibus. *Gooden. l. c.*

Carex praecox spicis tribus, mascula terminali, foemineis subpedunculatis, squamula capsula obtusa pubescenti aequalibus. *Jacqu. fl. Austr. 5. p. 23.*

Carex montana. *Poll. fl. pal. n. 885.* (ex synonymis). — *Lightf. fl. Scot. 2. p. 3.* et plurimorum botanicorum.

Carex filiformis. *Leers. fl. Herb. n. 718.* *Roth. fl. germ. 2. p. 443.*

Carex stolonifera. *Ehrh. gram. 99.* — *Hort. fl. germ. p. 328.*

Icon. Schkuhr. l. c. t. 286. F. n. 27. — *Leers. fl. Germ. 2. p. 443.*

ata, firma, saepe reflexa, margine scabri-
a. *Spica* terminalis mascula, florens sub-
ta, obtusa, erecta, pallide ferruginea;
ineae 1-saepissime 2 ad 3, perraro 4, al-
e, subsessiles, approximatae, mascula bre-
s, erectae, ferrugineae; infima breviter
nculata. *Bracteae* vaginantes, setaceae,
a spicula propria saepe longior. *Glumae*
ulae fusco-ferrugineae imbricatae, plus
sve obtusae; foemineae, apice aristatae.
ulae ovali-subtriquetrae, obtusiusculae,
integrae, subpubescentes. *Stigmata* tria.

ta. Non confundenda cum *Carice praecoe*
Schreberi descripto.

Sine dubio *Leerfius* l. c. sub *C. filiformi*
anc speciem et sub variatione γ . *Caricem vernam*
Schkuhrii. comprehendit.

2. *CAREX tomentosa* spicis femineis
subpedunculatis, capsulis subglobosis to-
mentosis. *Linn. Mant.* p. 123. *Schkuhr.*
l. c. p. 373. — *Schreb. Spic. fl. Lips.* p.
65. — *Leers. fl. Herb. n.* 717. — *Roth.*
fl. germ. 2. p. 442.

Carex spicis subpedunculatis subcylindricis,
bracteis auriculis amplexantibus folia-
ceis subapproximatis, squamis acutis,
capsulis globosis tomentosis. *Wahlenb.*
Act. Holm. 1803.

Carex sphaerocarpa. *Ehrh. gram.* 89.

1. *Schkuhr.* l. c. t. 286. F, n. 28. — *Leers.*
l. c. t. 15. f. 7.

m. *Filzige Segge.*

1. *Carex cotonneux.*

vaginantia, linearia, acuminata, quiniunciam ad binas tres uncias longa, subtus glauco-villofiuscula. *Vagina* firratae, incanae, villosulae. *Squamulae* versus radicem purpureae. *Spica* teretiuscula, firrata, teretiuscula, subuncinatae laterales, duae, raro tres, semiduplo breviores, subseffiles, subapice erectae. *Bractea* sub spica infima, ovata, foliacea, patens, uncialis. *Capsulae* ovatae obtusiusculae, ferrugineae, margine albidae, dorso flaveminae minores, ovatae, acutae, purpureae, dorso virides. *Capsulae* sub albedo-tomentosae, apice vix prominenti. *Stigmata* tria.

Nota. Facile dignoscitur vaginis foliis villosis, Capsulis subglobosis acutatis.

1443. *CAREX montana* spiculis sessilibus subsolitariis masculinis, culmo nudo. car.

squamis oblongo - obtusis, culmo nudo, capsulis ovatis utrinque acutis pubescentibus. *Schkuhr. l. c. p. 374.* — *Wahlenb. Act. Holm. 1803. p. 159.*

arex globularis spica feminea solitaria sessili sub mascula, capsulis euneiformis-ovatis, pubescentibus, culmo subnudo filiformi. *Willd. Act. Berol. 1794. p. 45. t. 2. f. 1.*

arex collina. *Willd. Linn. Sp. pl. 4. p. 260.*

arex foliis caryophylleis, spicis sessilibus approximatis paucifloris, capsulis ovatis subhirsutis. *Hall. stirp. helv. n. 1372.*

1. *Schkuhr. l. c. t. 286. F. n. 29.*

m. Bergsegge.

L. Carex de montagne.

1. In collibus sylvaticis apricis ad sylvarum marginalia passim frequens. Prope *Berghausen, Söllingen* in eoque toto tractu collino, montoso abunde.

r. Aprili, Majo. *Semina* Majo, Junio matura. ♀

Radix dense fibrosa. *Culmi* ex radice plurimi, palmares, pedales, erecti, triquetri, densi, graciles, aphylli, versus radicem foliosi. caespiticia, ex radice longe fasciculata, tria, carinata, angusta, culmo breviora, lineae retrorsum aspera, laete ex luteo-viridi. *Spica* terminalis, mascula, solitaria, erecta, quinque ad septem lineas longa, clavato-drica, fusca, *femineae* saepissime duae, albae, sessiles, masculae subapproximatae, pauciflorae, breves, quandoque saluberrimae. *Bractea* sub spica infima, setacea, brevior, rarius duplo longior. *Glumae* nigricantes, piceae, nervo dorsali viridi.

642 MONOECIA. TRIANDRIA.

obtusae, apice saepe emarginatae, aristis
Capfulae utrinque acutae, cuneiformi-ov
triquetrae, pubescentes. *Stigmata* tria.

Variat. Spicis femineis interdum quammaxime
proximatis, quandoque infima a reliquis magis
mota.

Obs. Variationem peculiarem inter millena indi
legi, Spicis femineis tribus basi proliiferis, mu
imbricatis.

Nota. Non confundenda cum *C. praecoce*
saepissime accidit, a quo facile distinguitur,
illis angustioribus, ex radice copiosissimis,
spiticiis, fasciculatis, Glumis calycinis
obtusis emarginatis, aristatis, Capfulis
que magis acutis.

Est vera *C. montana* Linnaei secundum
Willd. praef. de Schreber sententiam. An
cos calcareo-argillaceos, quos caespites pulcherr
ornat.

1444. *CAREX ciliata* spicis tribus, ma
la terminali, foemineis oblongis sel
bus, squamis subrotundis ciliato cre
tis, capsula clavata pubescenti brevi
bus. *Willd. Act. Berol.* 1794. t. 3.
— ej. *Linn. Sp. pl.* 4. p. 261. — *Schkuhr*
l. c. p. 384.

Carex ericetorum spica mascula termi
obverse ovata obtusa, femineis sessilibus
ovatis, capsulis pubescentibus. *Pohl*
fl. pal. n. 886. — *Roth. fl. germ.* 441
Wahlenb. Act. Holm. 1803. p. 159.

Carex montana. *Leers. fl. Herb.* n. 716.

Carex spicis sessilibus, approximatis ov
Hall stirp. helv. n. 1371. an?

Icon. Schkuhr. l. c. t. 286. I. n. 42. — *L.*
l. c. t. 16. f. 6.

n. Gefranzte Segge. Heiden-Segge. Frühlings Segge.

In graminosis pascuis sabulosis aridis, ericetis gramineis et juniperitis passim, nec non in nemoribus luculentis aridis apricis.

Februario, Martio, Aprili Majo. *Semina* Majo, Junio matura. z

Radix fibrosa, multiceps, densa. *Culmi* pluridigitales, palmares et altiores, erecti, triquetri, striati, versus radicem dense striati. *Folia* linearia, carinata, latiuscula, sesquipedem ad binas lineas lata, striata, firma, rufum aspera, erecta aut deflexa, longitudo culmi, etiam multo longiora et breviora, varia. *Spica* mascula terminalis, erecta, tria, septem lineas ad unciam fere longa, ter cylindrica, obtusa, crassiuscula; *feminae* masculae approximatae, sessilibus, alterae ad tres, quandoque unica, erecta, ovata, obtusae. *Bractea* sub singula spica, ovata, brevissima, fusca. *Glumae* masculae breviores, ovatae, obtusae, fuscescentes, omnes margine albidae, per lenspectatae lacero-ciliatae. *Capsulae* pyris, seu ovali-globosae, basi subelongatae, inscentes. *Stigmata* tria.

lores apibus ceram et mel largiuntur.

5. *CAREX pilulifera* vaginis nullis, spica mascula sublineari, femineis subrotundis sessilibus confertis, culmo debili. *Gooden. l. c. Schkuhr. l. c. p. 382.*

Carex pilulifera spicis terminalibus confertis subrotundis, masculina oblonga. *Linn. Sp. pl. 2. p. 1385. — Roth. fl. germ. 2. p. 445.*

Carex pilulifera spicis subseffilibus, globosis, mascula tenui, bracteis subamplectantibus attenuatis subfoliaceis approximatis, squamis mucronatis longiusculis, capsulis subglobosis acutis pubescentibus, culmo debili ferrulato. *Wahlenb. Ac. Holm.* 1803. p. 160.

Carex filiformis. *Poll. fl. pal. n.* 887. *an?* *videtur.* (*exclusis synonymis.*)

Carex decumbens. *Ehrh. gram. n.* 70. *Hoffm. fl. germ. p.* 329.

Icon. Schkuhr. l. c. t. 286. *J. n.* 39. — *Oeder. fl. Dan. t.* 1048.

Germ. Pilen-Segge.

Gall. Carex à pilules.

Hab. In sylvaticis graminosis siccis, passim dumetosis passim, cum *Festuca ovina* *duriuscula.*

Flor. Aprili, Majo. *Semina* Majo, Junio *matura.* 7

Radix fibrosa. *Culmi* palmares, pedales, erecti, debiles, graciles, triquetri, versus radicem foliosi, caeterum aphylli; fructiferi curvi aut decumbentes. *Folia* caespiticia, linearia, carinata, margine retrorsum aspera, culmo breviora, 1-2 lineas lata. *Spica* mascula terminalis, solitaria, linearis, 3-4 lineas longa, erecta; *femineae*, masculae approximatae, alternae, sessiles, 3-4, ovali-rotundae, pauciflorae. *Bractea* sub spica infima reliquis major foliacea, lineari-fetacea, patenti-erecta, 1-2 unciam, unciam ad 16 lineas longa. *Glumae* masculae, femineis angustiores, ovali-lanceolatae, acutae, fusco-rufescentes, nervo dorsali viridi, mucrone terminato. *Capsulae* sub

obsoletae, triquetrae, acutiusculae, viridescen-
tes, pubescentes. *Stigmata* tria, plumosa.

46. CAREX *pedata* vaginis brevibus
membranaceis subaequantibus aphyllis;
spicis femineis subornithopisformibus re-
ctis, mascula subbrevis, culmo com-
presso. *Schkuhr. l. c. p. 379.* — *Allion. fl.*
Pedem. n. 2320. — *Lamarck et Decand.*
fl. franc. 3. p. 119.

Carex ornithopoda. Willd. Linn. Sp. pl. 4.
p. 255.

Carex spicis foemineis raris marem aequan-
tibus, ima petiolata. *Hall. stirp. helv. n.*
1375.

Gramen caryophyllum nemorosum spica
multiplici. Bauh. Prodr. p. 9. — *Michel.*
gen. p. 65. n. 78. t. 32. f. 14.

Gramen parvum sylvaticum paniculis forma
pedum avium. Bauh. hist. 2. p. 499.

non. Schkuhr. l. c. t. 286. H. n. 37.

erm. Vogelfuss-Segge.

all. Carex pied-d'oiseau.

a b. In rupibus praesertim calcareis sylvaticis,
humidis retro *dem Thurnberg*, retro *Grö-*
zingen, et alibi. In Margraviatu superiori
in monte prope *Crenzach*; in dumetis hu-
midis inter *Wiesam* et *Birsam t. C. Bau-*
hino; in monte *Muteto*. In Principatu su-
periore in rupibus humidis *auf dem Schien-*
berg nuper legi.

lor. Aprili, Majo. *Semina* Junio matura. **7**

Radix fibrosa, fusca. *Culmi* plures, digi-
tes, palmates, et longiores, erecti, saepe

cat, acutae, margine albidae. Stigmata
truncatae, fuscae, margine albid
subdistichae, remotiusculae, triqu
obtusae, integrae, hispidae. *Stigmata*

Affinis *C. digitatae*, a qua differ
mineis ex uno fere puncto egredientibu

Obs. In Horto Botanico culta, mutati
buit. Specimina in alpibus calcareis
lenfibus Helvetiae, nec non in
nostratibus simillima sunt.

Nota. *Carex pedata* a Linneo in
ob folia pedalia angustissima, a n
cedit. Confer Linn. Sp. pl. :
nonyma tamen allegata, ad n
pertinent.

1447. *CAREX digitata* vag
naceis dimidiatis aphyllis,
bus erectis, mascula brev
distantibus. *Schkuhr. l. c. p*

Carex digitata spicis linearibus
cula breviora inferioreq
aphyllis, capsulis distantib
pl. 2. p. 1384. — Poll. fl. 2

MONOECIA. TRIANDRIA. 607

tenuioribus inter se distantibus. *Michel.*
gen. 65. *t.* 32. *f.* 9.

Gramen caryophyllatum montanum, spica
varia. *Bauh. Prodr.* 9.

on. Schkuhr. l. c. 380. *t.* 286. *H. n.* 38. —
Leers. fl. Herb. t. 16. *f.* 4.

r m. Fingerförmige Segge.

ill. Carex digité.

ib. In montosis sylvaticis, umbrosis humidi-
usculis, sepibus et dumetis passim.

or. Aprili, Majo. *Semina* Junio matura. 7

Carex fibrosa, fusca. *Culmi* plures, palma-
res, semipedales, pedales, erecti, triquetri-
compressi, versus radicem foliati, caeterum
aphylli. *Folia* culmo breviora, linearia, striata,
marginem scabra 1-2 lineas lata. *Spica* termina-
masculina solitaria, lateralis linearis, erecta;
lineae 2 saepe 3, longiores, erectae, disti-
cae, ferrugineae, suprema supra masculam,
ultima longe saepe pedunculata, a superiore
per longius remota. *Bracteae* vaginantes, mem-
branaceae oblique truncatae, aphyllae, spadi-
formis, purpureae. *Glumae* oblongae, truncatae,
obovatae, fuscae, margine albidae; *femineae*
ovo medio viridi. *Capsulae* alternae, remo-
culae, trigonae, acutae, integrae, basi at-
tenuatae, subhirtae. *Stigmata* tria.

Dist. Culmis altioribus et demissioribus. *Spicis*
femineis quam maxime a se invicem distantibus, et
magis approximatis, (tunc cave ne pro *C. pedata*
habeas) longioribus et brevioribus. *Squamae* folia-
ceae versus radicem pulcherrime purpureae.

48. *CAREX clandestina* spicis subses-
silibus inclusis paucifloris, bracteis vagi-

nantibus superne hyalinis ampliatisaphyllis remotis, squamis magnis, capsulis subglobosis basi elongatis pilosiusculis foliis filiformibus. *Wahlenb. Act. Holm.* 1803. p. 158.

Carex clandestina bracteis membranaceis subaphyllis vaginantibus, spicis foemineis remotis paucifloris vaginam vix exsuperantibus. *Gooden. l. c. p.* 167. — *Schkuhr. l. c. p.* 385. — *Smith. fl. brit.* 3. p. 98. — *Willd. Linn. Sp. pl.* 4. p. 254.

Carex humilis spica mascula oblonga, foemineis serioribus remotis subpedunculatis paucifloris, culmo folio brevior. *Leyf. fl. Hal. n.* 952. — *Schreb. Sp. p.* 65. — *Roth. fl. germ.* 2. p. 455. — *Lamarck et Decand. fl. franc.* 3. p. 111.

Carex prostrata. *Allion. fl. Ped. n.* 2312.

Carex spica mare unica, foemineis adpressis calycinis glumis maximis, caule sentescente prostrato. *Hall. stirp. helv. n.* 1372.

Cyperoides montanum humile angustifolium culmo veluti folioso, spicis obseis. *Scheuchz. Agrost. 4.* 407. t. 10. f. 1.

Icon. Schkuhr. l. c. t. 286. K. n. 43.

Germ. Heimlichblühende Segge.

Gall. Carex clandestine.

Hab. In Margraviatu superiori in monte prope Crenzach frequens.

Flor. Martio, Aprili. *Semina* Aprili, Maio matura. ¶

Radix fibrosa. *Culmi* plures, florentes 2-3 unciales, erecti, subflexuosi, triquetri, versus radicem dense foliosi, vaginati; fructiferi decumbentes. *Folia* clumo longiora, filiformia.

ta, demum decumbentia. *Spica* mascula inalis, solitaria, erecta; *foemineae* alternatae, pauciflorae, tres ad quatuor, parvae, nculatae. *Pedunculi* cum spicis floriferis in teis vaginantibus membranaceis, fuscis ine albidis, occulti. *Glumae* ellipticae, lae, fuscae, splendidae, margine albidae branaceae, nervo dorsali viridi; foemi-, apice mucronatae. *Capsula* pyriformis, a, albida, villosula, inflata. *Stigmata* tria. facile dignoscitur Spicis foemineis floriferis in brapassum et infimis in foliis occultis.

9. *CAREX filiformis* spicis masculis oblongis subgeminis, femineis subduobus remotis ovatis, capsulis apice bifidis, sublanuginosis, bracteis longis. *Schkuhr. l. c. p. 386.* — *Gooden. l. c.* — *Willd. Linn. Sp. pl. 4. p. 303.* — *Smith. fl. brit. 3. p. 1008.*

Carex filiformis spicis subsefilibus oblongis, bracteis brevissime vaginantibus foliatis remotiusculis, capsulis ovali-ellipticis villosis rostello bifurcato, foliis convolutis. *Wahlenb. Act. Holm. 1803. p. 153.*

Carex splendida. *Willd. prodr. fl. Berol. n. 103. t. 1. f. 3.* — *Roth. fl. germ. 2. p. 446.*

Carex lasiocarpa. *Ehrh. gram. 10.*

Carex tomentosa. *Lightf. fl. Scot. 2. p. 553.*

1. *Schkuhr. t. 286. K. n. 45.*

m. *Fadenförmige Segge.*

l. *Carex filiforme.*

. In palustribus, turfosis, limosis, udis, et fossis aquaticis.



gracilis, teretiuscula, ad basin tri-
erecta, solitaria, saepe binae, ad
inferne foeminea, superne masculi
foemineae a masculis remotae, alti
scissiles, 1-2-3, breviores, ovali
crassiusculae. *Bracteae* sub spicis le
ma foliacea longissima, erecta. *Glum*
latae, fuscae, apice subciliatae, foe
statae. *Capsulae* ovali-ellipticae, feri
bifidae. *Stigmata* tria.

Variat. Spicis masculis 1-2-3, foemin
bus saepe superne masculis.

+++++ *Spicis sexu distinctis, m*
ca, seu pluribus; femineis
dunculatis, vaginis breviori
matibus tribus.

1450. *CAREX ferruginea* va
diatis, spica mascula oblonga
subternis linearibus laxis, c
distantibus oblongis, apice
ceis saepius bifidis. *Schkuhr.*

- on. Schkuhr. l. c. t. 286. M. n. 48?*
erm. Rostfarbige Segge.
ll. Carex ferrugineux.
eb. In Ditione Basileensi in saxosis auf der Kallenflue. t. Zeyhero.
or. Majo, Junio. Semina Julio matura. 7
ota. Plantam Halleri a Zeyhero loco citato lectam, hucusque non vidi, hinc illam describere non possum. Mihi adhuc dubia, magisque in loco natali indaganda videtur.

51. CAREX pilosa vaginis dimidiatis, spicis femineis gracilibus subbinis; capsulis distantibus glabris ovatis apice membranaceis bifidis, squamis longioribus, foliis margine pilosis. *Schkuhr. l. c. p. 397. — Willd. Linn. Sp. pl. 4. p. 279. — Lamarek et Decand. fl. franc. 3. p. 123.*

Carex spicis exserte pedunculatis linearibus sparsifloris, bracteis vaginantibus foliolatis remotis, capsulis subovalibus acutis ore hyalino bilobo, foliis ciliatis. *Wahlenb. Act. Holm. 1803. p. 156.*

Carex pilosa culmo triquetro, spicis erectis remotis, femineis linearibus folio proprio longioribus. *Scop. fl. Carn. 2. n. 1162. Allion. fl. ped. n. 2323.*

Carex spicis foemininis petiolatis erectis, foliis remotis. *Hall. stirp. helv. n. 1379.*

Cyperoides latifolium hirsutum, spicis tenuissimis. *Scheuchz. Agrostr. p. 412.*

- on. Schkuhr. l. c. t. 286. M. n. 49.*
erm. Haarige Segge.
ll. Carex poilu.

Hab. In Principatu superiore prope Mörsb
In sylvis montosis, caeduis abunde
vidi.

Flor. Aprili, Majo. **Semina** Junio matur

Radix repens. **Culmi** pedales, sesquipedales et altiores, erecti, triquetri, aphylli, fusca radice fusco squamati. **Folia** radicalia et pedalia et longiora, linearia, carinata, lineas lata, carina et margine, molliter ciliolatae viridia, striata. **Spica** mascula solitaria terminalis, erecta, lineari cylindrica, gracilis uncialis; **femineae** 2-3 longe remotae, pedunculatae, erectae, unciales, graciles, floribus ternis remotiusculis. **Bractae** vaginatae pedunculatas involventes, foliaceae, longitudine spicarum. **Glumae** ellipticae, acutae, ex viridi fuscae margine subalbidae, nervo dorsali viridi. **Glumulae** remotiusculae, glumis aristatis longioribus subovatae, acutae, apice bifidae, hyalinae. **Stigmata** tria, plumosa.

Obs. In horto botanico culta, mutationi non succedit.

Nota. Individuum Lacum Bodanicum in foliis

n. Schkuhr. l. c. t. 286. N. n. 53. Oeder. fl. Dan. t. 1237.

r m. Schlankährige Segge.

ll. Carex épi-grele.

b. In sylvaticis udis Principatus superioris prope Salem et alibi nuper vidi.

r. Aprili, Majo. Semina Junio matura. 7

Primo intuitu quammaxime affinis *C. sylvatica* a quo differt *Spicis* semineis gracilioribus, *ribus* remotioribus, *Pedunculis* vagina *ma-*
a ex parte involutis, brevioribus, *Glumis*
rcinis viridibus margine albis, *Capsulis* *mi-*
argute elongato-rostratis.

53. CAREX *sylvatica* vaginis longis plusquam dimidiatis brevioribus, spicis filiformibus laxis pendulis remotis pedunculo brevioribus, capsulis ovatis acuminatis, rostratis apice tenuibus bidentatis. *Schkuhr. l. c. p. 415.*

Carex sylvatica spicis pedunculatis pendulis, mascula erecta, foemineis filiformibus laxis pedunculo brevioribus, capsulis ovatis aristatis furcatis. *Huds. fl. angl. 2. p. 411. n. 28. — Schreb. spicil. fl. Lips. p. 62. — Roth. fl. germ. 2. p. 457.*

Carex Drymeja spicis sexu distinctis, mascula unica, foemineis pedunculatis remotis pendulis filiformibus, capsulis remotis apice bifidis. *Linn. Suppl. 414. — Ehrh. Beyt. 2. p. 37. — Willd. Linn. Sp. pl. 4. p. 296. — Wahlenb. Act. Holm. 1803. p. 153.*

Carex patula spicis femininis pendulis raris praelongis, capsulis rostratis bifidis. *Poll.*

Hab. In Pring. — *Scop. fl. Carn.* 2. 1
In sylvis — *et Decand. fl. fr.*
vidi.

Flor. Apr. — *Lert. fl. Herb.* n. 72

Radix — *pendulis raris, c.*
les et altis — *Hall. stirp. he*

fus radica

cta pedat — *des sylvarum tenuius f.*

lineas lat — *hiff. 3. p. 243. — Sc*

laete v — *Scheuchz. Agrost. p.*

termina — *L. r. 1. 186. L. l. n. 10*

unciall — *am. 1. 404.*

latae,

terni — *Agrost.*

los — *Carex étalé.*

rum — *graminulis passim frequen*

mar — *Junio. Semina Julio matura*

fula — *Calina cubitales, triped*

sub — *Folia radicalia li*

St — *caernata, margine repli*

O — *planiuscula. Spica termi*

gracilis; femineae 4 sae

pedunculis praelo

pendulae, graciles,

uncias longae. Bractea

longae. Glumae

margine albidae, m

longae, acutae; femineae

acronatae. Capsulae ov

ventricosae, ere

inferiores remoti

rostratae, apice

pendula vaginis inferi

mediatis superioribus

aequantibus, spicis cylindricis longissimis pendulis, capsulis ovato-oblongis punctatis, apice membranaceo-subnatis. *Schkuhr. l. c. p. 404.* — *Gooden. l. c.* — *Willd. Linn. Sp. pl. 4. p. 288.* — *Roth. fl. germ. 2. p. 458.*

arex pendula spicis subsessilibus pendulis, mascula erecta, femineis cylindricis longissimis, capsulis subrotundis acuminatis. *Huds. fl. Angl. 2. p. 411.* — *Curtis. fl. Lond.*

arex Agastachys spicis sexu distinctis, masculis pluribus, femineis pedunculatis remotis pendulis longissimis, capsulis numerosissimis, approximatis, laxis, acuminatis. *Linn. Suppl. 414.* — *Ehrh. 19.* — *Wahlenb. l. c. p. 153.*

arex maxima. *Scop. fl. Carn. 2. n. 1166.* — *Lamarck et Decand. fl. franc. 3. p. 125.* — *Allion. fl. Ped. n. 2341.*

arex spicis femininis pendulis longissimis, capsulis mucronatis ovatis. *Hall. stirp. helv. n. 1396.*

cyperoides spica pendula longiore et angustiore. *Tournef. inst. p. 529.* — *Scheuchz. Agrostr. 445.* — *Michel. gen. p. 59.* — *Morif. hist. 3. p. 242.* — *Sect. 8. t. 12. f. 4.*

ramen cyperoides latifolium typha pendula longiori. *Barrel. Icon. 45.*

n. Curtis. fl. Lond. t. 180. — *Schkuhr. l. c. t. 286. Q. n. 60.*

m. Hängende Segge. Grofse Segge.

l. Carex altier.

*fl. pal. n. 896. — Scop. fl. Carn. 2. 116.
t. 59. — Lamarck et Decand. fl. frum.
3. p. 1760.*

ex capillaris. Leers. fl. Herb. n. 720.

*ex spicis femineis pendulis raris, capi-
lis rostratis bifidis. Hall. stirp. helv.
1395.*

*men cyperoides sylvarum tenuius spica-
tum. Moris. hist. 3. p. 243. — Sect. 1.*

heuchz. Agrost. p. 404.

t. 286. L. l. n. 101. —

104.

Carex étalé.

lis passim frequens.

r. Majo, Junio. Semina Julio matura.

*Radix fibrosa. Culmi cubitales, tripedu-
erecti, triquetri, foliati. Folia radicalia line-
ria, latiuscula, carinata, margine replicata,
scabra; culmea planiuscula. Spica terminalis
mascula, teres, gracilis; femineae 4 saepe
remotissimae, alternae, pedunculis praelongis
filiformibus triquetris pendulae, graciles, ses-
quiunciam ad tres uncias longae. Bractee se-
mivaginantes, foliaceae, longae. Glumae pal-
lide ex viridi-luteae, margine albidae, nervo
dorsali viridi, oblongae, acutae; femineae ob-
longae, apice mucronatae. Capsulae ovato-
acuminatae, subtrigonae, ventricosae, erectae
virides, demum fuscae, inferiores remotiores
superiores approximatae, rostratae, apice
cuspidatae. Stigmata tria.*

1454. CAREX pendula vaginis infer-
bus longis subdimidiatis superioribus

MONOECIA. TRIANDRIA. 655

aequantibus, spicis cylindricis longissimis pendulis, capsulis ovato-oblongis punctatis, apice membranaceo-subintegris. *Schkuhr. l. c. p. 404.* — *Gooden. l. c.* — *Willd. Linn. Sp. pl. 4. p. 288.* — *Roth. fl. germ. 2. p. 458.*

Carex pendula spicis subsessilibus pendulis, mascula erecta, femineis cylindricis longissimis, capsulis subrotundis acuminatis. *Huds. fl. Angl. 2. p. 411.* — *Curtis. fl. Lond.*

Carex Agastachys spicis sexu distinctis, masculis pluribus, femineis pedunculatis remotis pendulis longissimis, capsulis numerosissimis, approximatis, laxis, acuminatis. *Linn. Suppl. 414.* — *Ehrh. 19.* — *Wahlenb. l. c. p. 153.*

Carex maxima. *Scop. fl. Carn. 2. n. 1166.* — *Lamarck et Decand. fl. franc. 3. p. 125.* — *Allion. fl. Ped. n. 2341.*

Carex spicis femininis pendulis longissimis, capsulis mucronatis ovatis. *Hall. stirp. helv. n. 1396.*

Cyperoides spica pendula longiore et angustiore. *Tournef. inst. p. 529.* — *Scheuchz. Agrostr. 445.* — *Michel. gen. p. 59.* — *Morif. hist. 3. p. 242.* — *Sect. 8. t. 12. f. 4.*

Gramen cyperoides latifolium typha pendula longiori. *Barrel. Icon. 45.*

2. *Curtis. fl. Lond. t. 180.* — *Schkuhr. l. c. t. 286. Q. n. 60.*

m. *Hängende Segge. Grosse Segge.*

l. *Carex altier.*



tura. 7

Radix fibrosa. Culmus 3-4-5
ctus, triqueter, foliatus, striatus
lia linearia, carinata, striata, virid
et carina aspera, femiunciam lata
ciosissima, omnes pendulae; sup
la, basi attenuata, versus apicem in
rius androgyna, secunda et saepe
neae graciles, apice incrassatae, n
feriores 3-4 mere femineae long
lindricae, 4-5-6-7 uncias longa
praelongae, foliaceae. **Glumae** ov
latae, acuminatae, e fusco-purpu
viae. **Capsulae** ovali-oblongae, pun
membranaceo-subintegrae, acumi
mata tria.

Nota. Omnium speciosissima. Variat p
Spica saltem suprema mere mascul
nibus femineis, aut suprema uni
perne mascula.

Carex mutabilis. Willd. prod

squamis sublinearibus-aristae formibus,
capsulis oblongis longo rostratis retro-
versis, apice subfurcatis. *Schkuhr. l. c.*
p. 417. — *Willd. Linn. Sp. pl. 2. p. 295.*
— *Gooden. t. c. p. 188.* — *Wahlenb. t.*
c. p. 161.

rex *Pseudo-Cyperus* spicis pendulis, pe-
dunculis geminatis. *Linn. Sp. pl. 2. p.*
1387. — *Pollich. fl. pal. n. 891.* — *Röth.*
fl. germ. 2. p. 451. — *Lamarck et De-*
cand. fl. franc. 3. p. 128.

rex spicis femineis pendulis asperis, cap-
sulis longe rostratis retroversis. *Hall.*
stirp. helv. n. 1397.

yperoides spica pendula brevior. *Scheuchz.*
Agrost. 440. — *Morif. hist. 3. p. 242.*
Sect. 8. t. 12. f. 5.

n. Schkuhr. l. c. t. M. m. n. 102. — *Oeder.*
fl. Dan. t. 1117.

m. Cyper-Segge.

l. Carex Souchet.

b. In sylvis ad fossas aquaticas. Inter Rufs-
heim et Graben in sylva ad fossas. In di-
ctione Basileensi in udis sylvaticis Michel-
feldae t. Lachenalio; circa Basileam inter
Kluben et Klein-Huningen sub Friedlin-
gen t. C. Bauhino. In Palatinatu prope Lau-
tern ad rivi marginem in sylva Halgrund
inter populos et alnos; in piscina prope
molam imperialem; circa Germersheim,
Ladenburg etc. t. Pollichio.

r. Majo, Junio. Semina Junio, Julio ma-
tura. 7

Radix repens, fibrillosa. Culmi cubitales,
scissiles, erecti, superne antrorsum nutantes,

triquetri, angulis asperi, versus radicem foliat.
Folia linearia, longa, latiuscula, carinata, carina et margine scabra. *Spica* terminalis macula, cylindrica, acuminata, erecta; *feminae* 3-4-5 subapproximatae, alternae, longe pendunculatae, cylindricae, fructiferae pendulae, unciam ad tres uncias longae. *Bracteae* vaginifere destitutae, foliaceae, longae. *Glumae* neari-lanceolatae, angustae, longe aristatae, ferrugineae, margine albidae, nervo dorsali viridi. *Arista* setacea, ciliata. *Capsulae* oblongae, triquetrae, acutae, longe rostratae, retroversae, apice subfurcatae. *Stigmata* tria.

Facile glumis angustis, longe setaceo-aristatis dignoscitur.

1456. *CAREX fulva* vaginis inferioribus subdimidiatis, superioribus subaequantibus; spicis femineis fructiferis remissis ovato-oblongis; capsulis ovatis rostratis. *Schkuhr. l. c. p. 405.* — *Gooden. l. c. p. 177.* — *Willd. Linn. Sp. pl. 4. p. 270.* — *Lamarck et Decand. fl. franc. 3. p. 146.* — *Smith. fl. brit. 3. p. 991.*

Carex fulva spicis incluse pedunculatis ovato-oblongis, bracteis vaginantibus foliorum altitudine culmi distantibus, capsulis ovatis convexis ancipitibus subinflatis rostris longiusculo ancipiti, ore diviso, culmo ferrulato. *Wahlenb. l. c. p. 150.*

Carex culmo foliisque firmis erectis, spicis femininis quatuor breviter petiolatis erectis. *Hall. stirp. helv. n. 1382. B.*

Icon. Schkuhr. l. c. t. 286. T. n. 67.

Germ. Braungelbe Segge.

Gall. Carex fauve.

Tab. In pascuis pratisque udis, aquaticis. In ditione Basileensi prope *Michelfelden* in paludolis *t. Hallero*. In Principatu superiore ad Lacum Bodamicum prope *Constanz*, *Petershausen* et alibi in pascuis turfosis udis, nuper vidi.

Flor. Junio, Julio. *Semina* Julio, Augusto matura. 7

Maxime affinis *C. distanti* Linn. a quo distinguitur, Radice repente, Vaginis basi bractae truncatis, non albide membranaceis. Spicis semineis brevioribus minus longe remotis. Antheris magis obtusis, fuscis.

457. *CAREX distans* vaginis inferioribus subdimidiatis, superioribus subaequantibus, spicis oblongis remotissimis, capsulis ovatis acutis. *Gooden. l. c. p. 178.* — *Schkuhr. l. c. p. 406.* — *Smith. fl. brit. 3. p. 992.* — *Willd. Linn. Sp. pl. 4. p. 271.*

Carex distans spicis remotissimis subsessilibus bractea vaginante, capsulis angulatis mucronatis. *Linn. Sp. pl. 2. p. 1387.* — *Poll. fl. pal. n. 892.* — *Roth. fl. germ. 2. p. 454.*

Carex spicis brevissime petiolatis diffitis. *Hall. stirp. helv. n. 1382.*


Carex spicis parvis longe distantibus. *Scheuchz. Agrofl. p. 431.*

Non. *Schkuhr. l. c. t. 286. T. Y. y. n. 68.*

erm. *Entlegenährige Segge.*

all. *Carex distant.*

Tab. In pascuis pratisque udis aquosis passim frequens. Copiose in Principatu superiore ad Lacum Bodamicum.



tae, internodi 2-3 breviter pedunculati
ctae ovali-cylindricae, semiunciam c
gae. *Bracteae* vaginantes, foliaceae
culae, erectae, basi albidae membrana
mae ovali-acutae, fuscae, margine all
vo dorsali viridi. *Capulae* ovatae, ac
tricolae, triquetrae. *Stigmata* tria.

1458. *CAREX limosa* vaginis
subnullis; spicis femineis ovatis
sublonge pedunculatis pendu
lis ovatis apice integris. *Schku*
409. — *Gooden. l. c.*

Carex limosa spicis ovatis pendul
la longiore erectiore, radice
Linn. Sp. pl. 2. p. 1386. —
pal. n. 888. — Roth. fl. germ. 2.
Lamarck et Decand. fl. franc.

Carex elegans. Willd. prodr. fl. Bo
t. 1. f. 1.

Carex spicis femininis pendulis

vidi. In Principatu superiore ad lacum Bodamicum in turfosis, limosis, sphagno repletis. In Palatinatu prope *Lautern* in paludibus muscolis circa piscinas et rivulos, nec non circa *Germersheim* et *Neustatt. t.* Pollichio.

or. Majo, Junio. *Semina* Junio, Julio maturantur. ☞

Radix repens, fibrosa, fusca. *Culmi* palres, pedales, erecti, triquetri, tenues, alice ad medium fere foliosi. *Folia* culmiora, carinata, striata, rigidiuscula, ex virido-viridia. *Spica* mascula terminalis, erecta, cylindrica, semiunciam a unciam longa; *perianthiae* duae, alternae, remotiusculae, penniculatae, ovato-lanceolatae, obtusiusculae, demum pendulae. *Bractae* brevissime vaginantes, foliaceae, longiusculae. *Glume* ovatae plus minusve acutae, fusco-ferrugineae, nervo dorsali viridi. *Capulae* ovatae, quatuor, depressae, obtusiusculae. *Stigmata*

Nota. Non confundenda cum sequente specie a qua differt, Spicis masculis non 2-3; femineis brevioribus non cylindricis.

59. *CAREX flacca* spicis masculis pluribus erectis; femineis pendulis, capsulis ovatis obtusis bractea brevioribus. *Schreb. Spic. fl. Lips. app. p. 150. ad p. 62.*

Carex flacca vaginis brevissimis vel nullis spicis cylindricis, femineis laxis fructiferis pendulis, capsulis optuso-ovatis, apice integris. *Schkuhr. l. c. p. 421. — Wahlenb. Act. Holm. 1803. p. 160.*

Carex glauca spicis maribus una, pluribus, femininis pendulis, capsulis ovato tri-

nigra recurva. *Morif. hist.*
Sect. 8. t. 12. f. 14.

Icon. Schkuhr. l. c. t. 286. O. P. n.
Oeder. fl. Dan. t. 1051. — Lee
fig. 3.

Germ. Schlaffe Segge.

Gall. Carex flaccide. Carex glau

Hab. Utrinq. in pratis pascuisqu
folijs, aquosis, passim frequen

Flor. Majo, Junio. Semina Junio
tura. ☞

Radix repens, fibrosa. Culmi
quipedales, cubitales, erecti, obtu
glaucovirides, versus radicem foli
Folia linearia, carinata, latiuscul
retrorsum aspera, ex viridi glauca,
viora. Spicae masculae superiores
tertia saepe androgyna (rarius unica
erectae, graciles; femineae inferio
2-3-4 pedunculatae, flaccae, de
vae, pendulae, cylindricae. Bracte
nantes, foliaceae, longiusculae.

460. *CAREX panicea* vagina infima subdimidiata, superioribus subaequantibus, spicis erectis remotis; squamis capsulisque ovatis subacutiusculis, subdistantibus, capsulis apice integris. *Schkuhr. l. c. p. 414.*

Carex panicea spicis pedunculatis erectis remotis: semineis linearibus, capsulis obtusiusculis inflatis. *Linn. Sp. pl. 2. p. 1387.* — *Poll. fl. pal. n. 890.* — *Leers. Herb. n. 722.* — *Roth. fl. germ. 2. p. 450.*

Carex spica mare una pluribusque, femininis petiolatis, erectis, capsulis raris maximis ovato-triquetris bidentatis. *Hall. stirp. helv. n. 1405, an?*

Cyperoides foliis caryophylleis spicis e rarioribus et tumidioribus vesicis compositis. *Michel. gen. 61. t. 32. f. 11.*

On. Schkuhr. l. c. 286. L. l. n. 100. — *Oeder. fl. Dan. t. 261.* — *Leers. l. c. t. 15. f. 4.*

erm. Hirsen-Segge.

all. Carex panio.

a b. Utrisque in pascuis pratisque udis turfosis passim abunde.

For. Majo, Junio. Semina Junio, Julio, maturata. 7

Radix repens. *Culmi* palmares, pedales et superiores, erecti, subcurvati, obtuse triquetri, ferne foliati, laeves ex glauco-viridescentes. *Stemata* culmo breviora, latiuscula, subcarinata, glauco-viridia, margine scabra. *Spica* terminalis mascula, cylindrica; *femineae* inferiores ternae, pedunculatae duae, saepe unica, erectae, remotae, subcylindricae. *Bractea* infima



atur, a qua distinguitur, spicis
bus paucioribus remotioribus infra

Variat. Spica feminea infima, longissima
pedunculo ex basi culmi prodeunte,

1461. *CAREX pallescens* va
viatissimis, spicis femineis sul
fructiferis pendulis, squamis
liatis, capsulis oblongis obt
integris. *Gooden. l.c. Schkuh*

Carex pallescens spicis penduli
erecta, femineis ovatis imbr
fulis confertis obtusis. *Linn.*
1386. — *Poll. fl. pal. n. 88*
fl. germ. 2. p. 449. — Leer
n. 721.

Carex spicis femineis pendulis, c
to-conicis. *Hall. stirp. helv.*

Gramen cyperoides polystachio
spicis brevibus prope sumu

b. Utrunque in pratis pasouisque udis, arenosis sylvisque humidis passim frequens.

r. Aprili, Majo, Junio. *Semina* Junio, Julio matura. ☞

Radix fibrosa. *Culmi* palmares, pedales et caules, erecti inferne foliati, triquetri, villosuli, vel laeves. *Folia* linearia, latiuscula, uninerviata, lutescente-viridia, villosiuscula, culmibreviora. *Spica* terminalis mascula, teres, albido-ferruginea, erecta; *femineae* inferiores, alternae, pedunculatae, subremotae 2 3 o 4, florentes ovali cylindricae, aut ovales, compactae; fructiferae pendulae. *Bracteae* vaginabreviatis, foliaceae, longae, basi undulato-plicatae, infima reliquis longior. *Glumae* pallide virescentes, margine pallidiores ovali-lanceolatae, masculae margine superne ciliato-nervo dorsali viridi. *Capsulae* oblongae, compressissimae, integrae, turgidae, pallide virentes, demum ex viridi-fuscae. *Stigmata* tria.

riat. Spicis femineis cylindricis, ovalibus et fere globosis, longioribus et brevioribus. Spicis masculis duabus, altera minore,

ota. Ne confundas variationem Spicis subglobosis praeditam, cum *C. pilulifera*, a qua differt Spicis remotis, pedunculatis,

62. CAREX *alba* vaginis membranaceis plusquam dimidiatis brevioribus subaphyllis, spicis femineis linearibus laxis paucifloris; capsulis obtuse-ovatis mucronatis, apice membraneis subvaginantibus integris. *Schkuhr. l. c. p. 399.*

Carex alba spicis longiuscule exsertis pedunculatis sparsifloris paucifloris, bracteis vaginantibus hyalinis aphyllis remotius-

culis, squamis brevioribus capsula
bosa cum acumine, foliis setaceis
lenb. l. c. p. 158.

Carex alba culmo tereti nudo, spicis
culatis erectis albidis: feminea su-
rem elata, capsulis laevibus. *S.*
Carn. 2. n. 1148. — Haenke in
Coll. 2. p. 81.

Carex nemorosa. Schrank. fl. Bav. p.

Carex spicis femineis raris petiolatis
bus acutis marem aequantibus. *Ha-*
helv. n. 1377.

Cyperoides foliis tenuissimis caule sul-
tro, spicis exiguis albicantibus,
lis longis infidentibus, capsulis
tis rotundo-triquetris lutescent
apicem tenuissimum una praecipue
apertum coarctatis. *Michel. gen. 6*

Gramen sylvaticum angustifolium spi-
Bauh. prodr. p. 8. n. 21. — Sc
Agrostr. p. 410. t. 10. f. 4. 5.

Icon. Schkuhr. l. c. t. 286. O. n. 55.

Germ. Weisse Segge.

Gall. Carex blanc.

Hab. In Marggrayiati superiore in sylv-
aker supra *Vögesheim t. Vulpio.* In
Basileensi in sylvia Huningensi *t. C. B.*
in monte *Muteto, Wasserfallen et*
berg, nec non prope *Arlesheim.* In
cipatus superioris pinetis arenosis
bus passim, prope *Hohenbottmann*
nec non in Principatu *Fürstenberg*
per legi.

Flor. Aprili, Majo. *Semina* Junio ma-

Radix repens. Culmi palmares, pedales et oeres, erecti, obtuse triquetri, versus radium foliosi. *Folia* radicalia fasciculata, perantia, lineari-setacea, laete viridia, culmioriora, erecta. *Spica* mascula terminalis, erecta, lineari-oblonga, gracilis, circiter semiumm longa, albida; *femineae* infra masculam, alternae, remotae, longe pedunculatae, compactae, florentes culmo arcte appressae, parvae, laxae, pauciflorae. *Bractae* subaphyllae, persistentes, membranaceae, scariosae, pellulae, albae, splendentes, pedunculos dimittim vestientes. *Glumae* ovatae, latae, albae, membranaceae, obtusae; *femineae* mucrone terminatae. *Capsulae* in singula spica paucae, laxae, ovatae, obtuse triquetrae, tumidae, acute acutae, integrae, membranaceae. *Stigmata* tria, quandoque sed rarius duo.

Pulchella ob vaginas et glumas argenteas plantariat. Spicis femineis 1-2-3.

f. *Capsulae* saepissime utrilagini obnoxiae.

Ob radices longe lateque horizontaliter repentes, alium in rupibus ex arena congestis figit.

63. *CAREX paludosa* spicis oblongis subcylindricis vaginis nullis; mascularum squamis oblongis obtusis superioribus saepe aristatis; capsulis oblongo-ovatis apice brevissime bidentatis. *Schkuhr. l. c. p. 425.* — *Gooden. l. c. Willd. Linn. Sp. pl. 4. p. 305.* — *Wahlenb. l. c. p. 163.* — *Lamarck et Decand. fl. franc. 3. p. 130.*

Carex acuta spicis masculis pluribus obtusis, squamis obtusiusculis caule acutangulo. *Curtis. fl. Lond.*



Germ. Sumpf-Segge.

Gall. Carex des marais.

Hab. Utrinque in paludibus, ad
cas passim frequens.

Flor. Aprili, Majo, Junio, *Sem-*
lio matura. ♀

Radix repens. *Culmi* sesq-
aquosis saepe bipedales, tripeda-
lioli, angulis acutis asperis trique-
lia linearia, carinata, margine c-
ra, tres ad quatuor lineas lata,
ridia. *Spicae* masculae plerumqu
oblongae, remotiusculae, suprem
quiunciali; *femineae* inferiores
pedunculatae, suberectae, longio
apice saepe mascula. *Bracteae* fi-
rioribus longae, foliaceae, cul-
tes, sequentes sensim minores. *G-*
lae arcte imbricatae, ovali-oblon-
fusco-purpureae, nervo dorsali

ota. Sine dubio sub Carice acuta Linnaei militat.

34. *CAREX riparia* spicis masculis oblongis; femineis subcylindricis, vaginis nullis; squamis lanceolatis aristatis, capsulis oblongis, apice furdato bidentatis. Schkuhr. l. c. p. 426. — Gooden. l. c. — Willd. Linn. Sp. pl. 4. p. 306. — Wahlenb. l. c. p. 163.

Carex riparia spicis masculis pluribus, triquetris nigricantibus, acutis. Squamis aristato-acuminatis, capsulis subinflatis bicornibus. Curtis. fl. Lond.

Carex acuta spicis masculis pluribus, femineis subpedunculatis, erectis, capsulis ovato lanceolatis aristato acuminatis furcatis. Hudsf. fl. Angl. 2. p. 413.

Carex crassa. Ehrh. Beitr. 4. p. 43. — Roth. fl. germ. 2. p. 462.

Carex vesicaria. Leers. fl. Herb. n. 724.

Carex spicis masculis ternis, femininis numerosis erectis brevissime petiolatis capsulis bicornibus. Hall. stirp. helv. n. 1404.

Cyperoides aquaticum maximum foliis vix unciam latis, caute exquisite triangulari, spicis habitioribus erectis, squamis in aristam longius productis, capsulis oblongis bifidis. Michel. gen. 57. t. 32. f. 6.

n. Schkuhr. l. c. t. 286. Qq. Rr. n. 105. — Curtis. fl. Lond. t. 281. — Leers l. c. t. 16. f. 2. l. — Oeder. fl. Dan. t. 1118.

r m. Ufer-Segge.

41. *Carex des rives*.

reticulata. *Spicae* masculinae saepe
que, erectae, fusco-nigricantes,
acutae, congestae, bracteatae, fi-
ciali, inferioribus minoribus inae-
mineae 3-5 pedunculatae, remo-
ternae, ovato-acutae, erectae,
quando pendulae, supremae sessil.
Bractea sub spica infima longe se-
longior. *Glumae* imbricatae, lan-
flatae, ex nigro-purpureae, foem-
latiores, superne magis aristatae.
longae, striatae, tumidae, apice
cae. *Stigmata* tria plumulosa.

Variat. Culmis demissioribus et longi-
femineis sessilibus, breviter aut lon-
erectis, patentiusculis, pendulis.

Nota. Cum praecedente sub *Carice*
militat.

1465. *CAREX vesicaria* sp.
cylindricis, femineis oblon-
gatis, fructiferis patentibus
longis acutis, capsulis infla-

tis acuminatis biaristatis, culmo triquetro acuto. *Willd. Act. Berol.* 1794. p. 55.
— *Linn. Sp. pl.* 2. p. 1388. — *Poll. fl. pal.* n. 895. β. — *Leers. fl. Herb.* n. 724. γ.

Carex inflata spicis remotis supedunculatis erectiusculis, mascula erecta lanceolata, capsulis ovatis acuminatis calyce duplo longioribus. *Hudsf. fl. Angl.* 2. p. 412. — *Roth. fl. germ.* 2. p. 466.

Carex spicis maribus una pluribus, cylindricis, femininis teretibus pendulis, capsulis conicis rostratis bicornibus. *Hall. stirp. helv.* n. 1409.

Cyperoides vesicarium spicis viridantibus et subfuscis. *Scheuchz. Agrostr.* p. 470.

non. *Schkuhr. l. c. t.* 286. S. s. n. 106, — *Oeder. fl. Dan.* t. 647. — *Leers. l. c. t.* 16. f. 2. III.

non m. *Blasen-Segge.*

non ll. *Carex en vessie.*

non b. Utrinque in locis aquosis, udis, turfosis, limpis ad fossas aquaticas, paludes, piscinas passim frequens.

non r. Aprili, Majo. *Semina* Junio matura. ¶

Radix repens, fibrosa. *Culmi* cubitales, pedales, erecti, foliati, argute triquetri, an-
tis asperi. *Folia* linearia, carinata, latiuscula, una et margine hispida, culmo saepe longiora.
Stylae masculae 2-3 terminales, remotiusculae, ailes, erectae, cylindricae, graciles; *femi-*
lae 2 rarius 3, alternae, pedunculatae, remo-
lo, florentes graciles, erectae, fructiferae
Stiores, erecto-patientiusculae. *Bractea* sub
sa infima, foliacea, culmo longior. *Glumae*
longo-lanceolatae; acutae, subfuscae, nervo

tis subglobosis ampullaceiformis apice patente-bifidis; ca-
quetro. *Schkuhr. l. c. p. 429*
l. c. — Willd. Linn. Sp. pl.
Wahlenb. l. c. p. 163. — S.
3. p. 1006.

Carex ampullacea spicis masculinis
femineis pedunculatis, capitulis
globosis, acuminatis bifidibus,
culmo obsolete triquetro.
Act. Berol. 1794. p. 55.

Carex vesicaria. *Poll. fl. pal. n. 1*
fl. Herb. n. 724. a. — Roth
p. 464.

Carex obtusangula. *Ehrh. gra.*

Carex spicis maribus una plu-
vinis erectis teretibus, capitulis
latis rostratis. *Hall. stirp. helv.*

Icon. Schkuhr. l. c. t. 286. T t. n. 1
l. c. t. 16. f. 2. II.

Germ. Flaschen-Segge.

Jor. Aprili, Majo. *Semina* Junio matura. \mathfrak{z}

Affinis praecedenti, cum qua saepe confunditur, differt autem ab illa, *Culmis* non ariste sed obsolete triquetris, *Foliis* angustioribus, *Spicis* femineis fructiferis pendulis, *Capulis* ampullaceis, divergentibus, pallide ex viridi-luteolis, demum fulcis.

ariat. Spicis masculis 2-3-4, femineis 2-3 pedunculatis aut saltem infima pedunculata, reliquis sessilibus.

467. CAREX *hirta* pilosa spicis omnibus oblongis, femineis remotis, vaginatis, vaginis hinc lanato-villosis, capsulis hirtis. *Gooden. l. c. p.* 208. — *Schkuhr. l. c. p.* 431. — *Willd. Act. Berol.* 1794. *p.* 56. — *Smith. fl. brit.* 3. *p.* 1007.

Carex hirta spicis remotis, masculis pluribus, femineis subpedunculatis erectis, capsulis hirtis. *Linn. Sp. pl.* 2. *p.* 2389. — *Poll. fl. pal. n.* 897. — *Leers. fl. Herb. n.* 725. — *Roth. fl. germ.* 2. *p.* 467.

Carex caule foliis capsulisque hirsutis. *Hall. stirp. helv. n.* 1403.

Cyperoides polytachion lanuginosum. *Morif. hist.* 3. *p.* 243. *Sect.* 8. *t.* 12. *f.* 10. — *Scheuchz. Agrostr. p.* 478.

non. *Schkuhr. l. c. t.* 286. *Uu. n.* 108. — *Leers. l. c. t.* 16. *f.* 3.

erm. Borstige Segge.

all. *Carex hérissé.*

ab. In pascuis pratisque aggeribus praesertim sabulosis aridis, apricis, ad fossas et viarum marginalia passim frequens. Inter *Carlsruhe* et *Durlach* ad viam publicam.

Flor. Majo, Junio. *Semina* Junio, tura. 7

Radix longe lateque repens, articulata. *Culmi* palmares, pedales erecti, triquetri, inferne foliati, *vaginatae* villosae-hirtae. *Folia* latiusculi carinata, hirta. *Spicae* terminales duae saepe tres, sessiles, subapproximate inferiores, alternae, remotae viter pedunculatae, erectae, ovales rae crassiusculae. *Bractae* vaginante foliaceae, longae. *Glumae* masculae obtusae, *feminae* apice acutae, longiusculae, fusco-purpureae, nervo dorsali viridi hirtae. *Capsulae* oblongo-ovatae, ac apice furcatae, utrinque hirtae. *Stigma* plumulosa.

Variationem sane peculiarem Vaginis bractearum glabriusculis, Spicis femineis spiculis compositis, quasi proliferis nuper Bodamicum in fabulosis siccis graminosis Schkuhr. l. c. t. U u. n. 108. B.

Pharm. CARICIS HIRTAE Radix.



ifidum, f. quatripartitum: laciniis ovatis, ob-
 lis. *Corolla* nulla. *Feminei* flores in eadem
 anta. *Calyx Amentum* cylindraceum f. subro-
 ndum imbricatum, squamis bifloris. *Corolla*
 alla. *Perianthium* nullum. *Amentum* sub sin-
 da spuama semina duorum flosculorum fovens.
 mina solitaria, ovata.

5f. BETULA T. Stamina 10-12. Fructus in
 amentum cylindricum digesti. Squa-
 mae trifurcae. Sem. ala duplici late-
 rali.

ENUS T. Fructus in Strobilum subrotun-
 dum digesti. Squamae subrotundae. Se-
 mina angulata aptera.

468. BETULA *alba* foliis ovatis acumina-
 tis ferratis. *Linn. Sp. pl.* 2. p. 1393. —
Pollich. fl. pal. n. 898.

Betula foliis cordato-lanceolatis ferratis.
Hall. stirp. helv. n. 1628.

Betula. *Mapp. fl. Alf.* p. 38. — *Bauh. pin.*
 427. — *Tabernæmont.* p. 1398. *Icon.*

on. *Pfleger. arb. et fruct. Badens. Blackw.*
 t. 240. — *Regnault. bot.*

=rm. Gemeine Birke. Weisse Birke.

211. Le Bouleau. Bouleau blanc.

2b. Utrique in sylvis et nemoribus passim
 frequens.

or. Majo. Semina Septembri, Octobri ma-
 tura. 5.

Rami in arboribus junioribus erecti, in
 altioribus penduli, filiformes.

676 MONOECIA. TETRANDRIA.

Nota. BETULA pendula vix peculiarem speciem constituit.

Pharm. BETULAE Folia; BETULAE liquor Lympha.

Ex Betulae trunco vulnerato, verno tempore quor copiosus limpidissimus, dulciusculus, pergratissimus effluit qui vires resolventes, aperitivas exercet, ex quo Saccharum Mannae analogum, vinum et cerevisia parari possunt. Lignum durum (in borealibus durius tenax, album, circa nodos rubello colore tinctum, variegatum, ad rotas, axes, trahas et alia iusfrumenta nec non ad carbones praestantes utile.

Rami tenaces flexiles ad doliorum circulos adhibentur.

Virgae ad scopas, corbes et ad funes contorquendos inferviunt.

Cortex sic dictus intermedius, adustus, per se, cum Ledo palustri mixtus, destillatus, prodest oleum sic dictum Birkentheer, Russis aliisque partibus borealibus, coriis inungendis aptum, ab Europa Juchten dicta odorem habent. Bellon Voyag. III. p. 25.

Folia cum aqua et aluminae cocta flavo tingunt colore; eadem ovibus et capris per hyemem salubre pabulum.

Flores melliferi.

1469. BETULA *pubescens* ramulis pubescentibus foliis ovatis acutis duplicato-ferratis pubescentibus, strobilis cylindricis pedunculatis, squamarum lobis aequalibus. Ehrh. Beitr. 6. p. 38. Hoffm. fl. germ. 335. — Willd. Linn. Sp. p. 462.

Betula pumila Broccembergenfis. Thal. Horticyn. 20. — Bauh. pin. 427. — DuRoi Harbke 1. p. 92.

non. Pfleger arb. et frut. Badens.

erm. Weichhaarige Birke.

all. Buleau pubescent.

ab. In montosis udis turfosis auf dem Kaltenbrunn, auf der Herrnwiese, auf dem Kniebis et alibi passim. In Alsatiae montosis turfosis auf dem Hoehgebürg, Hochfeld, auf dem Ottilienberg prope Barr et alibi.

lor. Majo. Semina Septembri, Octobri matuta. b

Differt a praecedente specie affini, Statura multo minore, *Ramulis Foliisque* albide pubescentibus.

¶. In horto culta, mutationi ita succubuit, ut altior evaderat et pubem omnem fere deponeret hinc vix pro peculiari specie declaranda.

470. *BETULA Alnus* foliis subrotundocuneatis inaequaliter dentato-subretusis, glutinosis, subtus ad nervorum angulos villosis, pedunculis ramosis.

Betula Alnus pedunculis ramosis. *Linn. Sp. pl. 2. p. 1394. — Pollich. fl. pal. n. 899.*

Alnus glutinosa foliis subrotundo-cuneatis subretusis glutinosis, axillis venarum subtus villosis. *Willd. Linn. Sp. pl. 4. p. 334.*

Alnus foliis glabris rotundis, rotunde crenatis, spongiolis ad nervorum angulos, *Hall. stirp. helv. n. 1630.*

Alnus rotundifolia glutinosa viridis. *Bauh. pin. 428. — Mapp. fl. Als. p. 15.*

non. Pfleger arb. et frut. Badens,

erm. Gemeine Erle. Else. Eller.

678 MONOECIA. TETRANDRIA.

Gall. Aulne commune. Aulne glutineux.

Hab. Utrunque in humidis sylvis, ad fossa aquaticas et rivos passim frequens.

Flor. Martio. *Semina* Septembri, Octobri maturata. **h**

In hortis Carlsruhanis aliisque prostat hujus arboris variatio, foliis laciniatis.

Pharm. ALNI Cortex; Folia.

Cortex junioris arboris a nonnullis cortici Peruviano substituitur, adstringendi certe vires habet. Pro coriis parandis aptus. Piscatorum retia reddit fusca, et masticatus Lapponum coria rubra. Linn. fl. Suec. p. 337.

Folia recentia praestantissima ad lac puerperarum lactationem respuentium dissipandum, dum postquam concisa, in disco incaluerunt et sudarunt bis vel pluries quotidie calide imponuntur. Etiam hyeme applicari possunt, modo antea aqua calida conspergantur. Mart. app. med. I. ed. auct. p. 121.

Flores apibus mel et caeram largiuntur.

Radices sustinent ripas, et littora defendunt. Lignum molle, sub aqua quammaxime indurescit, ad siphones, urbium fundamenta, rotas coli, patinas, vasa lactis, manubria sclopetorum, pectinumque pratorum nec non ad ligneos calceos in quibusdam regionibus adhibetur.

1471. *BETULA ovata* foliis ovatis bilobatis, dentibus acutis, ramis angulosis, amentorum pedunculis ramosis. *Schrank. fl. Salisb. n. 159. — Fl. Bavar. 1. p. 434. Willd. Linn. Sp. pl. 4. p. 465.*

Alnus folio oblongo viridi. *Bauh. pin. 426.*

Alnus alpina minor. *Mapp. fl. Als. an?*

Germ. Eyblättrige kleine Ehrle. (Geringe) oder Ringe Ehrle in Brisgovia.

111. *Aulne à feuilles ovales.*

b. In Brisgovia prope *St. Mariam*, vulgo *St. Mergen* nec non am *Zwerrenbacher Wasserfall* nuper cum *Cl. Eckero* florentem abunde vidi. In ditione Basileenfi inter *Muttenz* et *Münchenstein* t. *C. Pauhino*. In Alsatiae sylvis erga *Hochfeld in dem Hochgebürg* copiose t. *Mappo*.

r. Majo. *Semina* Septembri, Octobri matura. *h*

Medium quasi tenet locum inter praecedentem et sequentem speciem, a quibus differt *tura* semper multo minore 3-4-5 pedali, *mis* angulosis, *Foliis* multo minoribus, utrinque viridibus, glabris, argute duplicato-ferratis, ovali-subrotundis, *Pedunculis* longioribus nosis.

Nota. Non confundenda cum *Betula fruticosa*. *Pallasii* aut *B. humili*, *Schrankii* quae differt a nostrate, Statura 2-3 pedali, Ramis teretibus, Foliis minoribus subrotundis, aequaliter et simpliciter ferratis, ferraturis grossiusculis, *Pedunculis* simplicibus.

72. *BETULA incana* pedunculis ramosis, foliis subrotundo-ellipticis acutis subtus pubescentibus, axillis venarum nudis, stipulis lanceolatis, *Act. hort. Pew. 3. p. 33.*

Betula incana foliis ovatis acutis biserratis incanis, stipulis lanceolatis, amentis spicatis. *Linn. Suppl. 417.*

Alnus foliis mucronatis acute ferratis, subtus lanuginosis. *Hall. stirp. helv. n. 1631.*

Alnus folio incano. *Bauh. pin. 428. — Mapp. fl. Alf. p. 15.*

Alnus incana et hirsuta. Bauh. hist. 1. p. 134.

Icon. Pfleger arb. et frut. Badens.

Germ. Graue Eller. Graue Ehrle.

Gall. Aulne blanchâtre.

Hab. In sylvaticis passim rhenanis prope *Alneuburgweyher* et alibi. Prope *Basileam* in sylva *Huningensi*; inter *Muttenz* et *Münchenstein*; inter *Klüben* et *Klein-Huningen*. In Principatu superiore ad *Lacum Bodamicum* in sylvis passim copiose, necnon in Principatu *Fürstenbergensi*.

Flor. Majo. *Semina* Septembri, Octobri maturata. **B**

Differt a *Betula Alno*, *Foliis* ovatis acutius inaequaliter ferratis, superne non glabris, sed hirsutis, inferne incanis pubescentibus.

Variat. *Foliis* plus minusve profunde inciso-ferratis.

Haec species amat praecipue montosa humidiora ubi citissime crescit, omnino magis propaganda.

Obs. In Hortis *Carlsruhanis* praesertim adhuc habentur *Betula populifolia* Ait. h. kew. *B. excelsa* Ait. *B. nigra*, papyracea, lenta, nana, fruticosa, pumila, *Alnus* β . *lacinia*, et *ferrulata* Ait.

CCCCXXIV. B U X U S.

Linn. Gen. pl. ed. Schreb. n. 1420.

Masculi flores. *Calyx* triphyllus: foliolis subrotundis concavis, patentibus. *Corolla* *Petalata* duo, subrotunda, concava simillima *calyci*, at majora. *Stamina* quatuor. *Antherae* didymae. *Germinis* rudimentum. *Foeminei* flores in eadem gemma cum masculis. *Calyx* 16

MONOECIA. TETRANDRIA. 685

thyllus: *Corolla* petala tria simillima calyci, majora. *Styli* tres brevissimi persistentes. *Stigma* obtusa, hispida. *Capsula* subrotunda, trirris, trilocularis, trifariam elastice dehiscens. *Semina* gemina, oblonga, hinc rotundata, inde ca.

73. *BUXUS sempervirens* caule fruticoso, foliis ovali-oblongis, petiolatis, antheris ovatis.

Buxus sempervirens β. *arborescens*. Linn. *Sp. pl.* 2 p. 1395. — *Hall. stirp. helv. n.* 1610.

Buxus arborescens foliis ovato-oblongis, versus apicem attenuatis antheris ovatis, caule arborescente. *Lamarch. Encyc. p.* 505.

Buxus arborescens. *Bauh. pin.* 471. — *Tabernæmont. p.* 1464. *Icon.*

Don. Pfleger arb. et frut. Badens. — *Bulliard. herb. t.* 263. — *Blackw. t.* 196.

Arm. Buchs.

11. *Buis.*

12. In Margraviatu superiore in adscensu montis prope *Crentzach*, in colle prope *Friedlingen* et *Höllstein* abunde. In *Sponhemia* prope *Winterburg*, nec non in rupibus schistosis ad *Mosellam* prope *Trarbach*, versus *Winnungen* frequens.

13. Martio, Aprili. *Semina* Julio matura. 5

Frutex in regionibus citatis magnitudinem 4-5 pedum habet, laete ibidem vegetat, et, et semina matura fert.

Arm. Buxi Folia.

682 MONOECIA. TETRANDRIA.

Lignum durum flavum, ponderosum, exqui

Folia nauseosi odoris et saporis, amara.

Nota. Variationes sequentes Foliis margine
Foliis lanceolatis, angustioribus etc. in B
Carlsruhanis cum Buxo balearica, foliis
ciosis, ovalibus, latis, firmis laeteviridibus,
mentis praelongis, hospitantur.

Obs. Buxus balearica hyemes nostras duriores
fert.

Nota. Ingens Buxi sempervirentis quantitas,
olim in horto aulico Carlsruhano et Duris
pro circumvallationibus viarum adhibebatur
Marggraviatu superiore, et nominatim ex
prope Crenzach apportabatur.

AUCUPA japonica ad hancce classem Mo
ciae tetrandriae pertinens, caule arboreo, foliis
fitis petiolatis, oblongis, remote ferratis, manu
sempervirentibus, sub dio in hortis Carlsruhanis
tatur.

CCCCXXV. URTICA.

Linn. Gen. pl. ed. Schreb. n. 1422.

Masculi flores. Calyx tetraphyllus: foli



MONOECIA. TETRANDRIA. 683

Urtica sexubus sede disjunctis, foliis ovato-lanceolatis, serratis, julis oblongis. *Hall. stirp. helv. n. 1615.*

Urtica urens minor. *Bauh. pin. p. 232. — Mapp. fl. Al. p. 333.*

n. Bulliard, herb. t. 230. — Oeder. fl. Dan. t. 739.

m. Kleine Brenneffel.

l. Ortie mineure.

Utrisque ad sepes, parietes, muros, et in cultis passim frequens.

n. Julio, Augusto. Semina Septembri, Octobri matura. ☉

Caulis pedalis, sesquipedalis et altior, erectus, tetragonus, viridis s. rubellus, stimulis dentibus obsessus. *Rami* oppositi. *Folia* opposita, petiolata, ovalia, obtusa, serrata, stimulis obsessa. *Racemi* axillares, ramosissimi. *Flores* sciculati, mares et feminei in eadem planta, sede disjuncti.

rm. URTICÆ MINORIS Herba.

5. *URTICA dioica* foliis oppositis, cordatis, racemis geminis. *Linn. Sp. pl. 2. p. 1396. — Poll. fl. pal. n. 901.*

Urtica stirpibus sexu distinctis, foliis serratis ovato-lanceolatis, productis. *Hall. stirp. helv. n. 1614.*

Urtica urens maxima. *Bauh. pin. 232. — Mapp. fl. Alf. p. 332.*

n. Oeder. fl. Dan. t. 746. — Blackw. t. 12. (mas.)

m. Grofse Brenneffel.

Gall. Ortie grande. Ortie dioique.

Hab. Utrunque ad sepes, ruderata, in nemoribus et sylvis humidiusculis passim abunde.

Flor. Julio, Augusto. *Semina* Septembri, Octobri matura. ♀

Statura altiore, *Foliis* cordatis, *Floribus* in distincta planta dioicis, facile a praecedente specie distinguitur.

Rharm. URticae MAJORIS Radix; Herba, Semina.

Vires habent Urticae resolventes, diureticas. Herba recens caesionibus in membris paralyticis jam ab antiquissimis temporibus inservit.

Turiones cocti, pergratum cibum praebent.

Radices cum alumine in aqua coctae, fila limbo colore tingunt.

Caules macerati, corticem deponunt, qui in tenuissima duci se patitur, ex quo contexta linteamina pulcherrima, sub nomine Nesseltuch nota. Necnon ex confusa cortice, chartae sat bonae parantur.

CCCCXXVI. M O R U S.

Linn. Gen. pl. ed. Schreb. n. 1424.

Masculi flores in amenta digesti. *Calyx* quadripartitus: *foliolis* ovatis concavis. *Corolla* nulla. *Stamina* quatuor. *Feminei* flores congesti vel in eadem, vel diversa planta a masculis. *Calyx* tetraphyllus. *Corolla* nulla. *Calyx* maximus, carnosus, succulentus factus. (*Bacca*) *Semen* unicum, ovatum, acutum.

476. *MORUS alba* foliis oblique cordatis laevibus. *Linn. Sp. pl. 2. p. 1398.*

Morus foliis scabris cordatis et semitrilobis, *Hall. stirp. helv. n. 1611.*

Morus fructu albo. *Bauh. pin. 459.*

Mon. Pfleger arb. et frut. — Schkuhr. bot. Handb. t. 290.

erm. Weisser Maulbeerbaum,

all. Morier a fruits blancs.

Lab. In China et Persia, Colebatur praecipue in variis Marchionatus Badensis regionibus circa *Durlach, Carlsruhe, Rastadt* et alibi.

lor. Majo, Junio. *Fructus* Julio maturi. **♂**

Arbor variat fructibus albis, violaceo-rubellis et nigris.

Nota. Ante triginta circiter annos maxima in copia ad loca citata colebatur, hodie quammaxime diminuitur.

Ufus foliorum notissimus.

Lignum durum, brunneo-flavum, exquisitum.

h. In regionibus nostris, nec non in Gallia saepe vidi individua mere mascula, alia autem feminina.

Nota. In hortis Carlsruhanis aliisque praecipue adhuc hospitantur *MORUS nigra, rubra, et papyracea* s. *BROUSONETIA papyracea, ♂. ♀.*

MONOECIA. PENTANDRIA.

CCCCXXVII. XANTHIUM.

Linn. Gen. pl. ed. Schreb. n. 1426.

Masculi flores compositi. *Calyx* communis plurium flosculorum, polyphyllus, imbricatus.

catus squamis tenuibus longitudine flosculi aequalibus. *Corolla* composita uniformis, tubulosa, aequalis, in hemisphaerium digesta. *Pria* monopetala, tubulosa, infundibuliformis, erecta, quinquefida. *Receptaculum* communis vix ullum, paleis distinguens flosculos. *Enei* flores infra masculos, in eadem planta plicati. *Calyx* *Involucrum* biflorum, diphyllum, foliolis oppositis trilobis, aculeis uncinatis vallatum. *Corolla* nulla. *Styli* duo paria, pillaria. *Stigmata* simplicia. *Drupa* sicca, ovata, oblonga, undique aculeis uncinatis tecta, apice bifida. *Semen* *Nux* bilocularis.

1477. *XANTHIUM Strumarium* *tenax* L.
inermi, foliis cordatis trinervatis. L.
Sp. pl. 2. p. 1400. — *Poll. fl. pal.* n.

Xanthium foliis semitrilobatis, aculeis uncinatis. *Hall. stirp. helv.* n. 1621.

Lappa minor f. *Xanthium* dioscoridis. B.
pin. 198. — *Tabernæmont. p.* 1157.
— *Mapp. fl. Als.* p. 334.

Icon. *Oeder. fl. Dan. t.* 970. — *Blackw. t.*
— *Schkuhr. bot. Handb. t.* 291.

Ger m. Spitzkletten. Kleine Kletten. Klette.

Gall. Lampourde glouteron.

Hab. In ruderatis et ad vias passim communis.
Circa Carlsruhe, Beurtheim, Ripp,
alibi frequens.

Flor. Julio, Augusto. *Semina* Septembri
tobri matura. ☉

MONOECIA. PENTANDRIA. 687

Caulis sesquipedalis, cubitalis, erectus, es, scabriusculus, ramosus. *Rami* alterni patentes. *Folia* alterna, petiolata, cordata, suboblonga, dentata, trinervata, scabra, pallide viridia.

ar m. LAPPÆ minoris Herba; Semen.

Planta in officinis nostris hodie vix cognita in Galienus sanguinis massam depurandam et contra scrophulosis adhibetur.

Pulvis seminum ad unciam dimidiam datus in Suecia erysipelate arcanum fuit. Linn. fl. Suec. n. 864.

78. XANTHIUM *spinosum* spinis ternatis. Linn. Sp. pl. 2. p. 1401.

Xanthium spinosum atriplicis folio. Moris. hist. 3. p. 604. — Sect. 15. t. 2. f. 3.

Xanthium lusitanicum spinosum. Pluck. alm. 206. t. 239. f. 1. — Herm. parad. bat. 246. Icon.

an. Volkhamm. Norimb. 149. t. 149.

ar m. Dornige Spitzklette.

all. Glouteron ou Lampourde épineuse.

ab. In Gallia meridionali et Hispania. Circa Carlsruhe am Holzmagazin, nec non ad vias publicas urbis passim, sine dubio ex horto botanico emissa, nunc quasi spontanea.

or. Julio, Augusto. Semina Septembri, Octobri matura. ☉

Caulis 2-3 pedalis, erectus, valde ramosus. *Folia* alterna, pedunculata, triloba, glabra, laete superne viridia, subtus incana. *Spines* in caule et ramis ad petiolorum basin ternatis, longae, pungentes, albidæ, splendentes.

CCCCXXVIII. A M A R A N T H U

Linn. Gen. plant. ed. Schreb. n. 1431.

Masculi flores in eadem planta cum *neis*. *Calyx* (pentaphyllus) vel triphyllus, *ctus*, (in exoticis speciebus pluribus colorat persistens: foliolis lanceolatis, acutis. *Corolla* nulla. *Stamina* (quinque) vel tria. *Femine* *res*. *Calyx* ut in *mare*. *Corolla* nulla. *Stylis* breves, subulati. *Stigmata* simplicia, persistia. *Capsula* ovata, subcompressa, trilocularis, circumscissa. *Semen* unicum, *hosum*, compressum.

Nota. Plures species *A. triandri* for

1479. *AMARANTHUS Blitum* glom
lis lateralibus trifidis, triandris, foliis
tis retusis, caule diffuso. *Linn. Sp.*
p. 1405.

Amaranthus procumbens, foliis ovatis

MONOECIA. TETRANDRIA. 669

or. Junio, Julio. *Semina* Augusto, Septembri matura. ☉

Caulis pedalis, sesquipedalis, diffusus, quandoque adscendens, fulcatus, glaber, viridis aut bellus. *Folia* alterna, petiolata, ex rhomboidata, integriuscula, retuso-emarginata, obtusa, viridia cum rubore mixta, venis albidis purpureis picta, glabra. *Flores* axillares racinalesque, spicato-glomerati, triphylli, quadri, virides, femineis quinquephyllis mixti. *Fructus* splendens, nigrum.

80. AMARANTHUS prostratus glomerulis triandris spicatis, floribus triphyllis, foliis rhomboido-ovatis, obtusiusculis retusis, caule prostrato ramoso. *Willd. Linn. Sp. pl.* 4. p. 387.

Amaranthus prostratus glomerulis triandris, floribus trifidis, foliis rhomboido-ovatis, caule procumbente. *Balbis Misc. bot.* p. 44. t. 10.

Amaranthus viridis. *Vill. fl. Delph.* 2. p. 567. — *Poll. fl. pal. n.* 903.

Blitum sylvestre spicatum. *Mapp. fl. Als.*

Blitum album minus. *Joh. Bauh. hist.* 2. p. 967.

Blitum rubrum minus. *Camerar. epitom.* 235. *Icon.*

erm. Liegender Amaranth.

ill. Amaranthe à épi.

b. In ruderatis, viis hortorum nec non in plateis urbium et pagorum neglectis passim frequens. Circa *Carlsruhe* et alibi cum *Chenopodiis* passim.

lora. Bad. Tom. III.

X x

690 MONOECIA. TETRANDRIA.

Flor. Julio, Augusto. *Semina* Septembri tura. ☉

Caulis sesquipedalis, cubitalis, prostratus, quandoque erectiusculus, basi tantum decussatus, obsolete angulosus, sulcatus, glaber, teretis, quandoque rubellus, ramosus. *Folia* ternata, petiolata, ex rhombeo-ovata, obtusiuscula, integra, viridia, striata, glabra. *Inflorescentiae* spicatae, cylindricae, axillares, terminalesque, ex albido-virides. *Calyx* in floribus masculis triphyllus, in femineis quinquelobus. *Stamina* tria. *Capsula* viridis, rugosula, calyce cincta. *Semen* orbiculatum, compressum, nigrum, splendens.

Nota. Sub A. viridi Linnei militabat.

MONOECIA. POLYANDRIA.

CCCCXXIX. CERATOPHYLLUM.

Linn. Gen. pl. ed Schreb. n. 1439.

Masculi flores. *Calyx* multipartitus: laciniis subulatis, aequalibus. *Corolla* nulla. *Stamina* 16-20. *Feminei* flores in eadem planta cum masculis. *Calyx* multipartitus, laciniis subulatis aequalibus. *Corolla* nulla. *Stylus* et *Stigma* simplex. *Drupa* ovata, acuminata, cortice tenui. *Semen Nux* unilocularis.

1481. *CERATOPHYLLUM demersum* foliis dichotomo bigeminis, fructibus spinosis. *Linn. Sp. pl. 2. p. 1409.* — *Pl. fl. pal. n. 904.*

Ceratophyllum foliis subspinosis, capsula corni. *Hall. stirp. helv. n. 1599.*

MONOECIA. POLYANDRIA. 691

eratophyllum asperum aquis immersum.

Mapp. fl. Als. p. 65.

quisetum sub aqua repens foliis bifurcis.

Loesel. prussic. p. 67. t. 12.

m. Spitzfrüchtiges Wasserhorn.

l. Cornifle äpre.

. In stagnis, aquis lente fluentibus, et foliis aquaticis praesertim in Rheni vicinia passim abunde, sub aqua.

. Julio, Augusto. *Semina* Septembri maturatione. ♀

Caules longi, copiosi, filiformes, debiles, i, saepe rubelli, dichotomi, sub aqua de-

. *Folia* verticillata, in quolibet verticillo

3, linearia, angusta, semibifida, scabra,

entis uno latere tuberculis dentibus ob-

. *Flores* ad foliorum verticillos sessiles, sub-

rii, parvi.

at. Capsulis basi sub muticis, et 1-2-3-4 spinosis.

sub aqua integras sylvulas quasi format.

2. CERATOPHYLLUM *submersum*

foliis dichotomo trigeminis, fructibus mu-

ticis. *Linn. Sp. pl. 2. p. 1409. — Leers.*

fl. Herb. n. 735. — Roth. fl. germ. 2.

p. 485.

eratophyllum laeve, aquis immersum. *Mapp.*

fl. Als. p. 66.

1. Oeder. fl. Dan. t. 510.

m. Stumpffrüchtiges Wasserhorn.

l. Cornifle submerge.

. et *Flor.* in iisdem cum praecedente regionibus, sed rarius.

Affinis praecedenti, a qua differt, *Folia* profundius divisibilibus glabris, *Calycis* foliolis apice dentatis, *Fructibus* ovalibus basi muticis.

CCCCXXX. MYRIOPHYLLUM.

Linn. Gen. pl. ed. Schreb. n. 1440.

Masculi flores. *Calyx* tetraphyllus: foliis oblongis, erectis: horum extimum majus, intermedium minus. *Corolla* Petala quatuor, caduca. *Stamina* octo. *Feminei* flores infra masculos. *Calyx* ut in mare. *Corolla* ut in mare. *Germina* quatuor oblonga. *Styli* nulli. *Stigmata* pubescentia. *Capsulae* quatuor tenues, monospermae. *Semina* quatuor, oblonga.

Obs. Interdum floribus hermaphroditis praeditum est.

1483. MYRIOPHYLLUM *spicatum* L.

ribus masculis interrupte spicatis. Linn. Sp. pl. 2. p. 1409. — Poll. fl. pal. n. 905.

Myriophyllum *spicatum* foliis omnibus pinnatis capillaceis, spica terminali verticillata nuda. Willd. Linn. Sp. pl. 4. p. 466.

Pentapteris spica nuda. Hall. stirp. helv. 2. 993.

Millesfolium aquaticum pennatum spicatum. Bauh. pin. 141. — Prodr. 73. t. 73.

Icon. Oeder. fl. Dan. t. 681.

Germ. Aehrenförmiges Federkraut.

Gall. Volant-d'eau à épi.

Hab. Utrinque in aquarum fossis stagnantibus praesertim in Rheno vicinia. Prope Dachs-

land, Knielingen, Eckenstein, Linkenheim
et alibi passim non infrequens.

lor. Julio, Augusto. *Semina* Septembri ma-
tura. ¶

Caules pedales et longiores, sub aqua rep-
ntes, demum adscendentes, ramosi, articu-
ti. *Folia* verticillata, quaterna, quina, sena
plura, pinnata, uncialia, glabra. *Flores* spi-
to verticillati, interrupti, supra aquae super-
iem eminentes, superne masculi, nudi, in-
ne feminei, parvi; infimi foliosi.

84. *MYRIOPHYLLUM verticillatum*
floribus omnibus verticillatis. *Linn. Sp.*
pl. 2. p. 1410. — *Poll. fl. pal.* n. 906.

Pentapteris floribus alaribus. Hall. stirp. helv.
n. 992,

Millefolium aquaticum, flosculis ad folio-
rum nodos. *Bauh. pin.* 141.

On. Oeder. fl. Dan. t. 1046.

r m. Quirlförmiges Federkraut,

l t. Volant d'eau verticillé,

z b. Cum praecedente in fossis et aquis flag-
nantibus passim.

or. Julio. *Semina* Augusto matura. ¶

Differt a praecedente, *Caule* subsimplici,
seto. *Foliis* brevioribus, saepe, quaternis,
tribus omnibus verticillatis, rarius superne
rnamphroditis et inferne femineis,

CCCCXXXI. SAGITTARIA.

Linn. Gen. pl. ed. Schreb. n. 1441.

Masculi flores plures. *Calyx* triphyllus: fo-
lis ovatis, concavis persistentibus. *Corolla*

Pimpinella minor sanguisorba hirsuta. Bauh.
pin. 160. — *Mapp. fl. Alf. p.* 240.

Sanguisorba minor. *Tabern. p.* 316. *Icon.*

β. *Pimpinella Sanguisorba minor laevis.* Bauh.
pin. 160. — *Mapp. fl. Alf. p.* 240.

Icon. Curtis. fl. Lond. — *Blackw. t.* 413.

Germ. Biebernelle. Becherblume.

Gall. Petite Pimpinelle.

Hab. Utrinque in pratis siccioribus apicibus
collibus graminolis asperis, nec non adve-
rum marginalia passim frequens.

Flor. Majo, Junio. *Semina* Julio, Augusto,
matura. ☞

Caulis palmaris, pedalis, sesquipedalis et
altior, erectiusculus, vel adscendens, angu-
tus, striatus, plus minusve villosus, quavis
que glabriusculus, ramosus. *Folia* alterna,
pari-pinnata; *foliola* ovato-orbiculata, dente-
ta, venulosa, villosa, interdum glabriuscula.
Stipulae denticulatae. *Flores* spicato-capitatae,
terminales. *Spicae* erectae, ovatae, obtusae,
densae, ex rubello virides, demum fusco-pur-
purascentes. *Flores* inferne masculi et hemi-
phroditi steriles, imperfecti, superne semina-
fertilis. *Calyces* 2-3-4 phylli.

Pharm. PIMPINELLÆ MINORIS s. PIMPINELLÆ
ITALICÆ Herba.

Leniter adstringentes vires habet.

Planta in hortis culta, villositatem fere exuit, in
aria et iuscula ingreditur. Egregium in pratis pabulum
in Anglia sub nomine Burnet colitur.

Non confundenda cum *Sanguisorba officinali*
Linnei.

CCCCXXXIII. Q U E R C U S.

Linn. Gen. pl. ed. Schreb. n. 1447.

Masculi flores. *Amentum* filiforme, longum, laxum. *Calyx* monophyllus, subquinque-lobatus: *laciniis* acutis saepe bifidis. *Corolla* nulla. *Stamina* 5-10. *Antherae* didymae. *Femineae* semina sessiles in eadem planta cum masculis. *Involucrum* squamis plurimis, imbricatis, in cupulam coriaceam coalitis, uniflorum, persistens. *Corolla* nulla. *Nux* ovali-cylindrica, carnosa, glabra, basi dearsa, unilocularis, recepta *Cupula* hemisphaerica, brevi, extus serrulata.

37. *QUERCUS Robur* foliis deciduis oblongis, superne latioribus: sinibus acutioribus, angulis obtusis. *Linn. Sp. pl.* 2. p. 1414. — *Poll. fl. pal.* n. 909.

Quercus sessiliflora foliis petiolatis deciduis oblongis, sinibus acutioribus oppositis, fructibus sessilibus. *Smith. fl. brit.* 3. p. 1026.

Quercus foliis glabris semipinnatis, pinnis obtusis, superioribus majoribus. *Hall. stirp. helv.* n. 1626.

Quercus latifolia mas, quae brevi pediculo est. *Bauh. pin.* 419. — *Mapp. fl. Alf.* p. 255.

var. *Pfleger arb. Badens.*

var. *Steineiche. Klebeiche.*

698 MONOECIA, POLYANDR

Gall. Chêne commune.

Hab. Utrique passim in sylvis frequ
Flor. Majo. *Fructus* Octobri, Nove
turi. 6


Pharm. QUERCUS Cortex; Folia.

Arbor maxime longaeva, perpulchra ult
tibus conservanda, ad aedificialia et navalia
Lignum durum in annosioribus durius, sub a
indurescit,

Cortex ad coria paranda adhibetur. ac
vires habet. Gallae spongiosae, cynipis op
saepè in abundantia ad quercuum ramos cre
coria paranda exquisitae. cum Vitriolo Marti
mentum scriptorium ut gallae Turcicae (a Qu
ris venientes) adhiberi possunt. Glandes sui
sumum pabulum, adstringentes.

Nota. Quercus folia saepe destruuntur et
tur à Phalaena processionaria.
arborum damno,

1488. *QUERCUS pedunculata*
deciduis oblongis subsessilibus,
superne latioribus, sinuatis, ne



11. *Chêne pedunculé.*

b. Cum praecedente passim, praesertim ad sylvarum marginalia.

or. Aprili, Majo. *Fructus* Septembri, Octobri maturi. **B**

Differt a praecedente, *Fructibus* longioribus, majoribus, longe pedunculatis.

arm. *QUERCUS Glandes.*

Glandes decorticatae, siccatae, tostae, Caeae in um paratae in Phthysi pulmonali a quibusdam com-
dantur.

89. *QUERCUS pyramidalis* foliis deciduis oblongis, superne latioribus, sinuatis, ramis omnibus erectis a basi trunci ad apicem exuntibus.

Quercus fastigiata. Lamarck *Encycl.* 1. p. 725.

arm. *Pyramiden-Eiche.*

11. *Chêne pyramidal.* *Chêne cyprès.*

b. Circa Carlsruhe im Hardtwald passim rarius.

Facile dignoscitur Statura stricta pyramidalis cherrima, habitum Populi italicae pyramidis repraesentans. *Fructus* pedunculati.

f. Arborea altae hujus speciei in horto Carlsruhano conspicuae, pulcherrimae.

Nota. In Hortis Carlsruhanis sub dio adhuc praecipue hospitantur, *QUERCUS Phellos*, — *Prinus*, — *aquatica*. Ait. tinctoria Mich. — *rubra* — *coccinea* Mich. *Catesbaei* Mich. — *palustris* Mich. *Aegilops*. *nigra*, *alba* — *Esculus* — *Cerris*.

CCCCXXXIV. JUGLANS.

Linn. Gen. pl. ed. Schreb. n. 1446.

Masculi flores. Amentum cylindraceum undique imbricato-sparsum, squamis unilobis extrorsum versis. *Calyx* ellipticus, planus, lacinie partitus: lacinie erecto-concavis, obtusis. *Gerula* nulla. *Stamina* 18-24. *Feminei flores* coacti. *Calyx* monophyllus, companulatus, quadrifidus, erectus, brevissimus, uniflorus. *Gerula* monopetala, quadrifida, erecta, acuta, calyce paulo major. *Germen* inferum. *Stigmata* duo, magna, reflexa, superne lacera. *Drupe* sicca, ovalis, magna, unilocularis. *Nux* maxima subrotunda, sulcis reticulata, semiquadrilobis: *nucleo* quadrilobo, varie sulcato.

1490. *JUGLANS regia* foliolis ovalibus glabris subserratis, subaequalibus. *Linn.*

Sp. pl. 2. p. 1415.

Juglans foliis septenis, ovato-lanceolatis integerrimis. *Hall. stirp. helv. n. 101.*

Nux juglans seu regia vulgaris. *Bauh. pin. 417.*

β. *Jux juglans* fructu maximo. *Bauh. pin. 417.*

γ. *Nux juglans* fructu tenero et fragili putrescente. *Bauh. pin. 417.*

δ. *Nux juglans* bifera. *Bauh. pin. 417.*

ε. *Nux juglans* fructu serotino. *Bauh. pin. 417.*

Icon. Pfleger arb. Regnault. bot. Bl. 6. 247.

MONOECIA. POLYANDRIA. 701

m. Gemeine Wallnuss.

11. Noyer commun.

b. In Persia. Colitur utrinque passim frequens.

r. Aprili, Majo. Fructus Septembri, Octobri matura. B

1 r m. JUGLANDIS Folia. Fructus immaturi, Cortex exterior fructus viridis.

Fructus immaturi saccharo et aromatibus conditi stolidi. Ex illis ope spiritus vini gallici, sacchari et nomi vel alius aromatis spiritus (Eau de Noix) s acceptissimus paratur.

Cortex exterior fructus viridis, amarus, acerbus, gum anthelminticum

Nuclei maturi recentes dulces, sapidi. Oleum nujuglandium expressum flavicans, dulce, egregium, re non coit, pictoribus acceptum, facile enitescit. Lignum praesertim in radicibus obscurum, pulchre variegatum, durum, pro elegantioribus scriptorum operibus acceptum. Folia et cortex fructus nior colore luteo-fusco tingunt.

In hortis Carlsruhanis et Schwezingentibus aliisque passim sub dio hospitantur JUGLANS alba, cinerea, nigra, et angustifolia. Ait. cuius arbor pulcherrima in Horto Schwezingensi.

CCCCXXXV. FAGUS.

Linn. Gen. pl. ed. Schreb. n. 1448.

Masculi flores. Amentum subrotundum vel draceum longum. Calyx monophyllus, cambratus, subsexfidus. Corolla nulla. Feminei in gemma ejusdem plantae. Involucrum monophyllum, quadrifidum, erectum, acutum, persistens. Corolla nulla. Germen triquetrum

702 MONOECIA. POLYANDRI

3-6 loculare, rudimentis seminum genus brevissimus 3-6 partitus. *Stigmata*. *Capsula* maxima, subrotunda spirulocularis, quadrivalvis. *Semina* N. f. tres ovatae, triquetrae f. hinc converplaniusculae.

1491. *FAGUS Castanea* foliis
tis acuminato-ferratis, subtus nu
Sp. pl. 2. p. 1416.

Castanea foliis ovato-lanceolatis
dentibus aduncis. *Hall. stirp.*
1623.

Castanea sylvestris. *Bauh. pin.* 419.
bernaemont. p. 1385. *Icon.* —
Alf. p. 60.

Icon. Pfleger arb. Badens.

Germ. Castanienbaum. Zahme Casta
Gall. Chataigner.

Hab. Prope Gernsbach, Baden, Bü
satz, Durrbach, Oberkirch et in



t, sic vidi ande viginti adhuc annos in regionibus ci-
is arbores in diametro ultra novem ad decem pedes
illas. Exemplum arboris insignis proceritatis et cras-
lei exstat prope Aetnam in Sicilia Castagnaro di-
nto Cavalli ibidem dictae, peripheria trunci ae-
at centum et sexaginta pedum.

492. *FAGUS sylvatica* foliis ovatis ob-
solete ferratis. Linn. *Sp. pl.* 2. p. 1416. —
Poll. *fl. pal.* n. 910.

Fagus foliis ovato-lanceolatis, oris undu-
latis. Hall. *stirp. helv.* n. 1622.

Fagus. Bauh. *pin.* 419. — Tabernaemont.
p. 1388. — Mapp. *fl. Alf.* p. 104.

non. *Pfleger arb. Badens.*

erm. Buche. Rothbuchen.

all. Hêtre.

ab. Utrunque passim integras sylvas constituit.

lor. Aprili, Majo. Fructus Octobri maturi. B

Arbor procera, longaeva. Lignum durum ad alen-
s focos et ad carbones praestans.

Nuclei decorticati dulces, oleosi, sapidi. Oleum
pressum praestans, vix oleo olivarum cedit.

Fructus in suum praecipue faginem cedunt, lar-
na tamen exinde non adeo firmæ est consistentiae,
am si sues glandibus vescantur.

Nota. In hortis Carlsruhanis praecipue hospitat
Fagus atropurpurea pro peculiari specie, ut
cultura ejus ex semine demonstravit, habenda.

CCCCXXXVI. C A R P I N U S .

Linn. *Gen. pl. ed. Schreb.* n. 1449.

Masculi flores. Amentum cylindraceum,
num, imbricatum: squamis ovatis, concavis,

704 MONOECIA. POLYANDRI

acutis, ciliatis, unifloris. *Corolla* nul
nei flores in eadem planta. *Amentum*
 bricatum, squamis lanceolatis, villosi
 reflexis, unifloris. *Involucrum* mono
 ovatum, persistens, sexfidum: *laciniis*
 libus. *Amentum* fructiferum maximu
 singulae squamae semen continens. Nu
 compressa, recta, involucrio persistente
 compresso, costato, ore sexfido: denti
 bus oppositis majoribus.

1493. *CARPINUS Betulus* squa
 hilorum planis. *Linn. Sp. pl.* 4.
Poll. fl. pal. n. 911.

Carpinus. *Hall. stirp. helv.* n. 1627.
fl. Alf. p. 57. — *Tabernaemont*
Icon.

Fagus sepium vulgo. *Bauh. hist.* 1
Icon. Pfleger arb. Badensf.

Germ. Hagenbuche, Hainbuche, W



MONOECIA, POLYANDRIA. 705

Lignum tenacissimum, durum, ad axes, molendi-
rotas, trochleas, pistilla exquisitum.

Cortex sic dictus interior aut Liber lanas luteo co-
nfit. Act. Stockh. 1742. p. 26.

Folia pecoribus pabulum sistunt.

ta. CARPINUS Ostrya et C. americana Mich.
in hortis Carlsruhanis et Schwezingensibus ho-
spitantur.

CCCCXXXVII. C O R Y L U S.

Linn. Gen. pl. ed. Schreb. n. 1450.

Masculi flores in amentum longum cylin-
m, imbricatum digesti, constans squamis
pris trifidis. Corolla nulla. Stamina octo.
nei flores remoti a masculis, in eadem
a, sessiles, gemma inclusi. Involucrum di-
um, coriaceum, margine lacerum, sub
centia exiguum. Corolla nulla. Styli duo
ei. Stigmata subulata. Nux ovata, basi
a.

4. CORYLUS *Avellana* stipulis ova-
tis obtusis. Linn. Sp. pl. 2. p. 1417. —
Pollich. fl. pal. n. 912.

orylus sepium. Hall. stirp. helv. n. 1625.

orylus sylvestris. Bauh. pin. 418. — Mapp.
fl. Alf. p. 82.

1. Pfleger arb. Badensf.

m. Haselftaude. Haselnussbaum.

l. Noisetier sauvage. Coudrier noisetier.

1. Utrunque in sepibus et sylvis passim fre-
quens.

706 MONOECIA. POLYANDR

Flor. Februario, Martio. *Fructu*
Septembri maturat. 5

Frutex, quandoque arbor medi
gae juniores villosae.

Multae prostant in hortis quoad nuci
calycem longiorem seu breviorē variation

Virgae tenaces, flexiles ad dolioru
cribra, corbes etc. adhibentur. Carbones
bus accepti. Nuclei dulces, nutrientes, sap
Oleum praestans, pictoribus acceptum.


Flores apibus ceram et mel praebent.

Nota. In hortis Carlsruhanis adhuc hos
RYLUS Colurna et rostrata. Ai

Obs. Ex hac Classe et ordine porro in hor
hanis sub dio hospitantur GINGHO bilc
seu SALISBURIA adiantifolia Smit
PLATANUS orientalis et occide
quibus pulcherrima ambulacrea circa u
non Platanus orientalis acerifol
LIQUIDAMBAR styraciflua. Linn.

MONOECIA. MONADELI

CCCCXXXVIII. PLIN. II.



Folius geminis seu pluribus ex eadem basi vaginali.

195. *PINUS sylvestris* foliis geminis: primordialibus folitariis glabris. *Linn. Sp. pl. 2. p. 1418. — Poll. fl. pal. n. 913.*

Pinus foliis geminis rigidis, conis ovato conicis longitudine foliorum subgeminis basi rotundatis. *Ait. hort. Kew. 3. p. 366.*

Pinus foliis binis convexo-concavis, conis masculis folitariis alaribus. *Hall. stirp. helv. n. 1660.*

Pinus sylvestris. Bauh. pin. 491. — Mapp. fl. Alf. p. 241.

n. *Pfleger arb. Badenf.*

rm. *Die Forc. Gemeine Fichte.*

ll. *Pin sauvage.*

b. Utrique praesertim in regionibus fabulosis passim frequens ubi integras sylvas format.

r. Majo. *Semina* Octobri, Novembri matura. *h*

rm. *Pini Turiones, Liber, Refina communis, alba, Pix liquida, atra, Terebinthina communis, Colophonium.*

96. *PINUS Pumilio* foliis geminis, truncis adscendentibus, strobilis ovatis erectis. *Waldst. et Kitaibl. pl. rar. Hung. 2. p. 160.*

Pinus foliis geminis abbreviatis, strictis, strobilis abbreviatis obtusis minimis, junioribus sessilibus erectis. *Lambert. monogr.*

Pinus sylvestris Mugho live *Crein. Bauh. hist. 1. p. 246. — Mapp. fl. Alf. p. 241.*

*Pinus Mugo. Tabernaem. p. 1343. Icon.
Icon. Pfleger arb. Badenf.*

*Germ. Krummholz. Krummholz-Fichte.
Gall. Pin mugho.*

Hab. In montosis turfosis auf dem Kalbsbrunn, am wilden Hornsee, auf der Herrnwiese am ausgetrockneten See, et in monte Kniebis in turfosis. In Alsatiae montosis turfosis, auf dem Hochgebürg, retro Perisverlus dem Schwarzen See, nec non in monte St. Ottiliae.

Flor. Majo. Semina Septembri, Octobri maturat. 5

Affinis P. sylvestri, a quo differt, Statura multo demissiore, Ramis a basi trunci ad terram decumbentibus vel adscendentibus, densis lobis nigro cortice tectis. Foliis ex singula vaginamineis, multo densioribus, crassioribus, hirsutioribus, magis saturate viridibus, Conis subtundis-ovatis, erectis, minoribus.

1497. PINUS *Larix* foliis fasciculatis obtusis. Linn. Sp. pl. 2. p. 1420.

Pinus foliis fasciculatis mollibus obtusifoliis, bracteis extra squamas strobilorum exsertantibus. Ait. hort. Kew. 3. p. 369.

Pinus foliis fasciculatis deciduis. Hall. stirp. helv. n. 1658.

Larix folio deciduo, conifera. Bauh. hist. 1. p. 265. — Tabernaem. p. 1348. Icon.

Icon. Pfleger arb. Badenf.

Germ. Die Lerche. Lerchenbaum.

Gall. Meleze d'Europe.

26. In alpinis helveticis aliisque. Indigena passim circa Carlsruhe et praesertim in montosis prope *Ebersteinburg*, *Baden* et alibi, nec non in Principatu superiore, et Principatu Fürstienbergenli, ubi hodie pulcherrimas jam sylvas integras constituit.

27. Aprili, Majo. *Semina* Octobri matura. 5

Arbor praesertim montosa subalpina amat, ubi cetera et longaeva est. Lignum exquisitum, firmum, tubellum, putredini resistens.

ARM. TEREBINTHINA VENETA. AGARICUS LARYCIS.

Ut aliae quaedam acerosae arbores, ita et haec quum folia eruperint Mannae speciem exsodat, graecoriandri semini similia, condensata, quae Manna picea s. Brigantina (Manne de Briançon) vocantur.

Cortex pro coriis condensandis adhibetur.

†† *Foliis solitariis et basi distinctis.*

98. PINUS *Picea* foliis solitariis emarginatis. *Linn. Sp. pl.* 4. p. 1420. — *Poll. fl. pal. n.* 914.

Pinus foliis solitariis planis, pectinatis, emarginatis. *Hall. stirp. helv. n.* 1657.

Pinus *Abies* foliis solitariis emarginatis: conis oblongis erectis, squamis subrotundis planis, basi acuminatis. *Du Roi Harbke. 2. p.* 95.

Pinus *alba* foliis subtus argenteis, emarginatis, conis erectis. *Mill. Dict. n.* 1.

Abies taxi folio, fructu sursum spectante. *Mapp. fl. Als. p.* 1.

Abies conis sursum spectantibus sive mas. *Bauh. pin.* 505.

Abies femina f. *Elate teleja*. *Rauh. hist.* 1.
p. 231.

Abies. Tabernaem. p. 1347. Icon.

Icon. Pfleger arb. Badenf.

Germ. Weifstanne Edeltanne.

Gall. Sapin en peigne. Pine blanc.

Hab. Utrique in montosis altioribus ubi integras sylvas constituit.

Flor. Majo. *Semina* Septembri, Octobri maturantur. **B**

Arbor speciosa, longaeva. Lignum album, elasticum, instrumentis musicis aptum.

Pharm. TEREBINTHINA COMMUNIS, ARGENTORATENSIS.

1499. *PINUS Abies* foliis solitariis fimbriatis mucronatis laevibus bifariam versis.
Linn. Sp. pl. 2. p. 1421.

Pinus foliis solitariis tetragonis mucronatis.
Hall. stirp. helv. n. 1656.

Pinus Picea foliis solitariis subulatis, bifariam versis: conis oblongis pendulis, squamis ovalibus planis marginibus undulatis et laceris. *Du Roi Harbke. 2. p. 110.*

Pinus excelsa. Lamarck fl. franc. 2. p. 201.

Abies tenuiore folio, fructu deorsum inflexo.
Mäpp. fl. Alf. p. 2.

Picea major prima sive Abies rubra. Banks. pin. 493.

Picea. Tabernaem. p. 1347. Icon.

β. *Pinus viminalis* ramulis elongatis pendentibus, flexuosis, laxis. *Alstroem. in act. Holm. 1777. p. 297. t. 8. 9.*

*non. Pfleger. arb. Badens. — Regnault. bot.
erm. Rothe Tanne. Schwarztanne. Fichte.
Kiefe.*

all. Sapin élevée. Pine femelle.

*ab. Utrinque passim in montosis cum prae-
cedente.*

For. Majo, Junio. Semina Novembri matura. 5

Arbor excelsa, speciosa, longaeva. Lignum opi-
tis variis, instrumentis musicis etc. inservit. Cortex
monae caritati in septentrionalibus regionibus subve-
nens.

*arm. ABIETIS Turiones; Coni; Ramus-
culi; Thus; Refina; Pix burgundica.*

Refina pro colore vario, alba, flava, bruna
cta.

*5f. In sylvis circa Carlsruhe et passimalibi Magni Ducis
Badensis CAROLI FRIDERICI (qui arbores
inter primarias suas delicias habet) auspiciis intro-
ductae sunt. PINUS sylvestris rubra et ta-
tarica, P. Strobilus (Weimouths Fichte)
pulcherrimae arbores, quae circa Carlsruhe jam ex
feminibus propagantur, P. balsamea et Cana-
densis. In hortis porro hospitantur sub dio P.
Pineaster. P. rigida. Du Roi. taeda, pa-
lustis, P. Cembra. P. Cedrus quae hyemes
apud nos aegre fert. P. laricina Du Roi. P.
balsamea. P. mariana f. nigra. P. mariana
alba.*

Nota. THUJA occidentalis ad hanc Classem
pertinens hospitatur in parte sylvatica im Pha-
sanengarten, am Baad Steinbach nec
non in hortis, ubi ex feminibus propagatur, ve-
rum lucorum ornamentum, pro sepibus semper-
virentibus densis pulchris apta. Frondes in dolori-
bus rheumaticis, et febribus intermittentibus nec
non in hydrope omnino efficaces. Canadensibus
sub ARBORE VITAE nota.

Porro in hortis Carlsruhanis aliisque sub dio
hospitantur THUJA orientalis, CUPRES-
sus disticha et Thyoides. Linn.

CCCCXXXIX. BRYONIA.

Linn. Gen. pl. ed. Schreb. n. 1480.

Masculi flores. *Calyx* monophyllus, campanulatus, quinquedentatus: denticulis subulatis. *Corolla* quinquepartita, campanulata, calyci adnata. *Antherae* quinque, quarum duas in singulo filamento connatae, unica in tertio filamento. *Feminei* in eadem planta cum masculis, aut dioici. *Calyx* et *Corolla* ut in masculis. *Stylus* trifidus. *Stigmata* emarginata, petiula. *Bacca* subglobosa, laevis. *Semina* nulla, subovata.

1500. *BRYONIA alba* foliis quinquelobis utrinque calloso scabris, baccis nigris.

Bryonia foliis palmatis utrinque calloso scabris, Linn. Sp. pl. 2. p. 1438.

Bryonia alba baccis nigris. Bauh. pin. 247. Mapp. fl. Alf. p. 43.

Bryonia. Tabernaem. p. 1291. Icon.

Icon. Bulliard. herb. t. 55. — Oeder. fl. Da t. 813. — Blackw. t. 533. — Regnault. bot.

Germ. Zaurrüben mit schwarzen Beeren.

Gall. Bryone.

Hab. Ad sepes passim rarius.

Flor. Majo, Junio, Julio. Baccae Augusto, Septembri maturae. ¶

Radix fusiformis, pedalis, et major, saepe ramosa, ex lutescente albida, transversim rugosa, interne alba, carnosa, succulenta post sectionem transversalem a centro ad peripheriam

diſque punctata. *Caules* plures, ſcandentes, xuofi, ſæpes obducentes, anguloſi, calloſoſabri, debiles. *Folia* alterna, petiolata, palmato-quinqueloba, ſinuata, calloſoſcabra. *Fructi* oppoſitifolii, ſpirales. *Flores* axillares, ſemoli. *Corollae* pallide ex luteo virescentes. *ſecae* maturae nigrae, globoſae. *Semina* ovata, compreſſa, ſubrugosa, ſulca.

arm. *BRYONIAE Radix*; *Turiones*; *Buccae*; *Semina*.

Radix recens lacteſcens, acriſcula, naſeoſa, ſtica, emetica, hydragoga, in Hydrope et obſtrictionibus inteſtinalibus ſæpe ſummo cum ſucceſſu adhibetur.

Amylum dat album friabile, in aqua decoctum ſtinoſum, ſubſatuum, mite.

501. BRYONIA dioica foliis palmato-quinquelobis, calloſoſcabis, floribus dioicis, baccis rubris.

Bryonia dioica, foliis palmatis utrinque calloſoſcabis, floribus dioicis. *Lin. Syſt. veg. ed. Murr. p. 810.*

Bryonia foliis palmatis ſcabis. *Hall. ſtirp. helv. n. 574.*

Bryonia alba. *Poll. fl. pal. n. 915.*

Bryonia aspera ſ. *alba* baccis rubris. *Bauh. pin. 397. — Mapp. fl. Alf. p. 43.*

On. Jacqu. fl. Auſtr. t. 199. — Blackw. t. 37.

arm. *Zaunwurz mit rothen Beeren.*

all. *Bryone dioique.*

ab. Utrinque ad hortorum ſepes in pagis et urbibus paſſim abunde.

cucurbitis nonnullis vago recipitur.

Nota. Non confundenda cum Rad
nigrae, quae a Tamo com

Obf. Ex hac Classe et ordine in hortis
passim adhuc coluntur. CUCUR
CUCUMIS Melo et fativus. L

CLASSIS XXII.

I O E C I A.

DIANDRIA.

- XL. SALIX. ♂ *Ament.* squama. *Cor.* o.
Stam. 2. raro 5.
♀ *Ament.* squama. *Cor.* o.
Stigm. 2. *Capf.* 2-valvis. *Semina* papposa.

TRIANDRIA.

- XLI. EMPETRUM. ♂ *Cal.* 3-partitus.
Cor. 3-petala.
♀ *Cal.* 3-partitus. *Cor.* 3-petala. *Styl.* 9. *Bacca* 9-sperma.

TETRANDRIA.


- XLIII. HIPPOPHAE. ♂ *Cal.* 2-partitus. *Cor.* o.
♀ *Cal.* 2-fidus. *Cor.* o. *Pist.* 1. *Bacca* 1-sperma: arillo truncato.
XLII. VISCUM. ♂ *Cal.* 4-partitus. *Cor.* o.
♀ *Cal.* 4-phyllus. *Cor.* o.
Stigm. obtusum. *Bacca* 1-sperma, infera.
* *Urtica dioica*,

DIOECIA.

PENTANDRIA.

CCCCXLV. CANNABIS. ♂ *Cal.* 5.*Cor.* nulla.♀ *Cal.* 1-phyllus. *Co*2. *Nux.*CCCCXLVI. HUMULUS. ♂ *Cal.* 5.*Cor.* nulla.♀ *Cal.* 1-phyllus. *Co*2. *Semen* calyce alCCCCXLIV. SPINACIA. ♂ *Cal.* 5.*Cor.* nulla.♀ *Cal.* 4-fidus. *Co*4. *Semen* 1, calyci* *Salix pentandra.*

HEXANDRIA.

CCCCXLVII. TAMUS. ♂ *Cal.* sexpartit♀ *Cal.* 6-partitus.

DIOECIA.

717

ENNEANDRIA.

- XLIX. MERCURIALIS. ♂ *Cal.* 3-phyl-
lus. *Cor.* nulla. *Stamina* 9-12.
♀ *Cal.* 3-phyll. *Cor.* o. *Styl.*
2. *Capsf.* 2 - cocca.

- L. HYDROCHARIS. ♂ *Cal.* 3 - phyll.
Cor. 3 - petala.
♀ *Cal.* 3 - phyll. *Cor.* 3 - pe-
tala. *Styl.* 6. *Capsf.* infera, 6-
locularis.

DECANDRIA.

- Lychnis dioica.* * *Cucubalus Otites.*

ICOSANDRIA.

- * *Spiraea Aruncus.*

MONADELPHIA.

- LI. JUNIPERUS. ♂ *Ament.* *Cor.* nulla.
Stam. 3.
♀ *Cal.* 3-partitus. *Cor.* 3 pet.
Styl. 3. *Bacca* infera, 3 sper-
ma calycina.

- LII. TAXUS. ♂ *Cal.* 4 - phyllus. *Cor.* o.
Antherae 8 - fidae.
♀ *Cal.* 4-phyll. *Cor.* o. *Stigm.*
1. *Bacca* monosperma, re-
cutita.

SYNGENESIA.

- naphalium dioicum.* * *Bryonia dioica.*
-

Salix pentandra foliis ellipticis acuminatis
ferratis glabris, petiolis superne glandu-
lofis, amentis serotinis pentandris, ge-
minibus lanceolatis glabris. *Willd. Linn.*
Sp. pl. 2. p. 658.

Salix foliis glabris ovato-lanceolatis, pe-
tis glandulosis, floribus hexastemonibus
Hall. stirp. helv. n. 1639.

Salix folio laureo f. lato glabro odorato. *Rap.*
hist. 1420.

Icon. Pfleger arb. et frut. Badensf. — Oeder.
fl. Dan. t. 943.

Germ. Fünfmännige Weide. Lorbeerweide.
Baumwollenweide.

Gall. Saule à cinq étamines.

Hab. In Principatu superiore ad Lacum Bo-
micum passim frequens, nec non in Prin-
cipatu *Fürstenbergensi* nuper legi.

Flor. Aprili, Majo. *Semina* Junio, Julio
tura. 5

Frutex 4-6 9 pedalis, quandoque in ar-
buscam adscendens. *Rami* erecti, fusci. *Folia*
alterna, petiolata, ovata, acuta, quandoque
obtusa, glaberrima, crassiuscula, (laurina) for-
ma, superne splendentia, laete viridia margi-
na cartilagineo-ferrata. *Petoli* superne glandu-
lofi. *Stipulae* saepe nullae. *Stamina* plerum-
que quinque, rarius sex ad septem. *Antherae*
pulcherrime luteae, magnae. *Capsula* glabra
Pappus longus, gossypinus.

Pulchra omnino species, folia trita odorata.

Pharm. SALICIS LAUREAE f. PENTANDRAE
Cortex.

Pappus in Gossypii modum tractari potest.

Flores apibus mel et ceram praestant.

4. *SALIX vitellina* foliis ferratis ovatis acutis, ferraturis cartilagineis, petiolis calloso-punctatis. *Linn. Sp. pl.* 2. p. 1442. — *Roth. fl. germ.* 3. p. 503.

alix vitellina foliis lanceolatis acuminatis glabris subtus glaucescentibus ferratis, petiolis pilosiusculis, amentis coaetaneis, germinibus sessilibus lanceolatis glabris, stigmatibus sessilibus bilobis. *Willd. Linn. Sp. pl.* 1. p. 668.

alix foliis ferratis, ovato-lanceolatis acutis, supra glabris: ferraturis cartilagineis. *Hoffm. Salic. I.* p. 57.

alix sativa lutea, folio crenato. *Bauh. pin.* 473.

alix lutea, tenuior, sativa, viminea. *Bauh. hist.* 1. p. 214.

n. Pfleger arb. et frut. Badens. — *Hoffm. l. c. t.* 11. 12. f. 1. 2. 3. et T. 24. f. 1.

m. Gelbe Weide, Eydotter-Weide, Gold-Weide, Band-Weide.

l. Saule jaune. Saule d'oziere.

o. Utrunque passim ad fossas aquaticas. Copiose in regionibus rhenanis aliisque passim colitur.

r. Aprili, -Majo. Semina Junio matura. B Vulgatissima, facile colore virgarum amoene luteo, ino dignoscitur. Virgae tenues flexiles.

rm. SALICIS VITELLINAE Cortex.

Adstringentes roborantes et antisepticas vires habet.

a. Cortices salicis verno tempore ex arborum solo modice humido crescentium ramis junioribus colligendi, et caute siccandi sunt.

Plantatur ad rivulorum et fossarum oras. Ad ligamenta et corbes exquisita. Flores apibus accepti.

Ex semine culta, mutationi non succubuit.

a Bad. Tom. III.

Zz

110. *Baum. p. 475.*

Salix spontanea, fragilis, auriculata et non auriculata.

1. p. 214.

Germ. Mandelweide.

Gall. Saule Amandier.

Hab. Passim in humidis ad aquas.

Flor.

Maxime affinis *S. triandra* videtur, *Statura* minore, *Folia* subangustioribus, *Stipulis* subdentatis, *Staminibus* duobus?

Florentem hucusque frustra duas observavit Pollichius.

Rami tenaces flexiles, fusco-rubri et corbes conficiendas utiles.

Nota. *Salix amygdalina*, triobliquis serratis glabris, geratis, stipulis maximis Smith p. 1045. — Willd. Linn. Vix ullibi in Germania, sed in liae palustribus crescit.

1506. *SALIX fragilis* foliis

amentis coaetaneis, nectario masculorum gemino, germinibus subseffilibus lanceolatis glabris. *Willd. Linn. Sp. pl.* 4. p. 669. alix foliis lanceolatis acuminatis undique ferratis glaberrimis, petiolis dentato-glandulosis, nectario masculinorum gemino. *Smith. fl. brit.* 3. p. 1051.

alix decipiens. *Hoffm. fl. germ.* p. 343. ej. *Salic.* 2. p. 9.

alix folio longo latoque splendente, fragilis. *Raj. angl.* 3. p. 448.

n. *Hoffm. Salices.* t. 31. — *Schkuhr. bot. Handb.* t. 317. b. n. 6. (folium).

m. *Bruchweide. Glasweide.*

l. *Saule fragile.*

b. In *Brisgovia* prope *Villingen*. In Principatu superiore passim. In Principatu Fürstenbergenli prope *Engen* et alibi passim non infrequens. In Palatinatu circa *Lautern* prope molam imperialem ad aquas sepes et in pratis paludosis passim. t. *Pollichio.*

r. Aprili, Majo. *Semina* Junio matura. 5 *Arbor* adulta sat procera, erecta. *Cortex* reus, rimosus. *Rami* patenti-erectiusculi, ni, subcinerei, juniores pallide fusco-rubri, laxissime inarticulati, ut digito concussi, iant et decidant. *Folia* alterna, petiolata, lanceolata, acuminata, collose ferrata, utrinque glabra, laete viridia, duriuscula. *Petiolus* rne dentato-glandulosus, glandulis apice micantibus. *Stipulae* saepe nullae aut duae latae, crenatae, saepe deciduae. *Amenta* cylindrica, obtusa. *Flores* diandri. *Nectaria* parvae flavescentes. *Nectaria* gemina foeminea cylindrica. *Germina* coepta, subseffilia, glabra.

Obs. Salicem nostram nullibi hucusque in den-
rhenanis et mosellanis aliisve regionibus,
per in fridioribus humidis montosis plagis

Nota. SALIX ANDROGYNA Rothii f.
2. p. 523. est annotante Cl. Willd.
monstruosa variatio S. fragilis.

Pharm. SALICIS FRAGILIS Cortex.

Roborantes febrifugas et antisepticas vir-
exercet. Hinc cortici peruviano a quibusdam
tur, praesertim in febre tertiana, et viscera
adhibetur. conf. ROSENBLAD de usu c
salicis in febribus intermittentibus
AKERBERG. 1782. GUNZ. comment d
Salicis 1787.

1507. SALIX praecox foliis lato-
latis acuminatis glanduloso-ferru-
bris subtus glaucis, amentis praeco-
germinibus sessilibus ovatis glabri-
lis elongatis. *Willd. Linn. Sp. pl.* 4

Salix praecox. Hoppe.

Salix bigemmis foliis ferratis glabris
latis petiolatis, gemmis duabus



o r. Februario, Martio. *Semina* Majo matura. **5**

Differt a praecedente specie, cum qua saepe confunditur. *Statura* humiliore. *Ramis* fusco-fis. *Foliis* non utrinque virentibus, sed sub- semper glaucis, supra viridibus, demum gricantibus, nitidis. *Amentis* praecocioribus. *Arminibus* sessilibus.

Rami huius etiam speciei fragiles sunt.

508. *SALIX babylonica* foliis ferratis glabris lineari-lanceolatis, ramis pendulis. *Linn. Sp. pl.* 2. p. 1443.

Salix orientalis, flagellis deorsum pulchre pendentibus. *Tournef. Cor.* 41. *Du Ham. arb.* 20.

Salix arabica folio atriplicis. *Bauh. pin.* p. 475.

Carb arabum. *Rauw. itin.* 183. *Dalech. app.* 130.

erm. Babylonische Weide. Trauer - Weide. Hängweide. Tränenweide.

all. Saule de Babylon. Saule du Levant. Saule pleureuse.

ab. In Oriente, Barbaria. Hospitatur apud nos in multis hortis, aliisque locis.

o r. Martio, Aprili. **5**

Arbor pulcherrima, saepe et id praesertim locis humidiusculis sat procera. *Virgae* praenigae, tenues pendulae. *Folia* lineari-lanceolata, glabra, ferrata, subtus glaucescentia, supra laete viridia. *Stipulae* minimae.

b f. Speciosissima individua procera sane pulcherrima vidi ad Aranjuez in Castilia.

1509. *SALIX purpurea* monandra decumbens, foliis ovato-lanceolatis serratis glabris, stigmatibus brevissimis ovatis subsessilibus. *Smith. fl. brit.* 3. p. 100.

Salix purpurea foliis serratis glabris lanceolatis: inferioribus oppositis. *Linn. Sp. pl.* 2. p. 1444. — *Huds. fl. Angl.* 2. p. 427. — *Poll. fl. pal.* n. 919.

Salix vulgaris nigricans folio non serrato. *Bauh. pin.* 473. — *Mapp. fl. Alf.* p. 174.

Salix folio longo lubluteo non auriculato, viminibus rubris. *Raj. Synopf.* 450.

Icon. *Curtis. fl. Lond.* (sub *S. monandra*.)

Germ. Purpur-Weide. *Rothe Korbweide.*

Gall. Saule pourprée.

Hab. Utrunque passim in pascuis paludosis, ad aquarum ripas cis et transrhenum cum Helice.

Flor. Martio, Aprili. *Semina* Majo matura.

Frutex decumbens, 2-4 pedalis et altior. Virgae adscendentes, lucidæ, saepissime purpureae aut corallinae, tenacissimae. *Folia* alterna inferiora saepe opposita, petiolata, obovato-lanceolata, subacuminata, versus apicem praesertim serrulata, glabra, subtus glauca. *Stipulae* nullae. *Amenta* mascula subsessilia, basi bracteolata, cylindrica, semiunciam ad unciam longa, gracillima. *Nectaria* solitaria, squamis opposita. *Stamen* unicum in singulo flore. *Anthera* fulva. *Amenta* foeminea masculis subsessilia. *Germen* subsessile, ovato-ellipticum, sericeum. *Stylus* subnullus. *Stigmata* subsessilia, subovata, integra, superne sulco exarata. *Capitula* sericea.

Distinguitur a *Salice Helice* cum qua saepe confunditur, Caulibus decumbentibus Virgis multo tenuioribus, magis purpureis, Foliis maxime amaris, superioribus saepissime oppositis, Amentis femineis tenuioribus et gracilioribus, subseffilibus.

Rami saepe sanguineo-corallini. Gemmae unilobae promunt Folia 3, transversim cauli applicata (contra in plerisque) ex 2 exterioribus transversim positis. Rami tenacissimi omnium. Linn.

Egregia ad vincendum, nec non ad corbes mires et alia utensilia conficienda.

Nota. Diligentius in regionibus rhenanis inundationes saepe patientibus expositis propaganda, et colenda.

510. *SALIX Helix* monandra erecta, foliis lanceolatis acuminatis ferrulatis glabris, stylo elongato filiformi, stigmatibus linearibus. *Smith. fl. brit.* 3. p. 1040.

Salix Helix foliis ferratis glabris lanceolato-linearibus, superioribus oppositis obliquis. *Linn. Sp. pl.* 2. p. 1444. — *Poll. fl. pal.* n. 920.

Salix monandra. *Hoffm. Salic.* 1. p. 18. *Vill. fl. Delph.* 3. p. 767. — *Roth. fl. germ.* 2. p. 507.

Salix monandra foliis glabris lineari-lanceolatis ferratis superne conjugatis, julis tomentosis. *Hall. stirp. helv.* n. 1640.

Salix tenuior folio minore utrinque glabro fragilis. *Joh. Bauh. hist.* I. p. 213.

Salix humilis, capitulo squamoso. *Bauh. pin.* 474. — *Mapp. fl. Als.* p. 275.

non. *Hoffm. l. c. t.* 1. f. 1. 3. t. 5. f. 1. t. 23. f. 1. *Curtis. fl. Lond.* (sub *S. monandra*).



Rami alterni, erecti, cinerei, o
minusve fusco - purpurascens.
superiora saepe opposita, petiola
ceolata, utrinque attenuata, a me
ferrata, glabra, subtus glauca. *Amenta* mascula opposita, altern
terminalia, cylindrica, gracilia, c
ctiuscula, saepe curvata. *Filame*
flore unicum, erectum, pilosun
plo longius. *Anthera* quadrangul
primo rubello - aurantiaca, deini

Nota. III. Smith, l. c. Hanc spec
maxime affini distinguit *Styl* o
mi. Sed Characteres a *Styl* o e
desumpti circa *Salicum* species
laces et inconstantes sunt. Sic sae
que individuo *Stylus* mox n
vissimus, et mox filiformis elong

III. Curtis l. c. Sub *S. mor*
praecedentem speciem combinat

Obf. Pro diverso loco natali demissior
ciori variet. foliis plus minusve et

511. SALIX rubra floribus diandris, foliis lineari-lanceolatis acutis ferrulatis, glabris concoloribus, filamentis basi subconnatis.

■ **Salix rubra monadelpha?** foliis lineari-lanceolatis elongatis acutis denticulatis glabris subtus concoloribus. *Smith. fl. brit.* 3. p. 1042.

■ **Salix fissa** floribus diandris, filamentis basi cohaerentibus, foliis lineari-lanceolatis glabris subintegerrimis. *Willd. arb.* 337.

■ **Salix fissa** foliis integris oblongo-lanceolatis acuminatis glabris. *Hoffm. Salic.* 1. p. 61. — *Roth. fl. germ.* 2. p. 510.

■ **Salix rubra** foliis integerrimis glabris lineari-lanceolatis acutis. *Huds. fl. angl.* 2. p. 428.

Salix folio angusto longo acuto non ferrato glabro, ex utraque parte atro viridi. *Mapp. fl. Als.* p. 277. an?

Salix minime fragilis foliis longissimis utrinque virentibus non ferratis. *Raj. Synops.* 449.

con. *Hoffm. l. c. I. t.* 13. 14.

erm. *Rothe Weide.*

all. *Saule rouge.*

ab. In Principatu superiore ad Lacum Bodamicum passim rarius. In Principatu Fürstenbergensi prope *Donaueschingen* in aquosis. In Alsatia prope *Eschau* in locis uliginosis. t. *Mappo.*

lor. Martio, Aprili. *Semina* Majo matura. 5

Frutex 4-6 pedalis, saepe arbuscula, corice cinereo glabriusculo vestita. *Rami* viminei,

marcula interna, ovar-cylindrica,
 cta, demum patentia, breviter
 uncialia et longiora, ante anthesin
 lutea. *Squamae* densae pilosae.
Filamenta basi plus minusve co-
 subrotundae, didymae, luteae,
 cae. *Nectarium* subcylindricum
 ventricolum. *Amenta* foemina
 erecta, demum reflexa, masculina
 oblongum, basi ventricosum.
Capsula ovato-subulata,
 pressa.

Obs. Folia juniora ut fere in omnibus
 pubescentia.

1512. *SALIX arbuscula*
 glabris subdiaphanis subtus
 suffruticosa. *Linn. Sp. p.*
Salix foliis ferratis glabris
fl. Lapp. 352. t. 8. f. c.
Salix foliis ovato-lanceolatis
 stipulis lanceolatis. *Ha.*
 1647.

duarum Leucarum diffito a *Donaueschingen* ubi nuper eam legi.

Cor. Martio, Aprili, Majo. *Semina* matura. 5

Fruticulus radice longe lateque reptante. *caules* lignosi, palmares, semipedales, vix al-
res, ramosi, diffusi, fusci, glabri, tenues. *Folia* alterna, sparsa, breviter petiolata, ovali-
ptica, apice acutiuscula, margine obsolete
glandulo-ferrulata, utrinque glabra, supra
viridia, subsiriata, splendentia, subtus
glauca, nervis a costa media ad mar-
em elevatis percussa, unciam longa, semi-
unciam lata, tenuia, subdiaphana, juniora ob-
scura, complicata, minora, villosula. *Amenta*
minora crassiuscula, erecta, semiuncialia, vix
matura. *Germina* ovato-lanceolata, glabra.
Stipulus longus, sericeus, albus.

Mascula individua non vidi.

Nota. Planta nostra perfecte cum caractere a Lin-
naeo dato convenit. Folia quoad formam op-
time respondent folio a Linnaeo in flora
Lapponica exhibito, recedunt saltem, quod iusto
minora sint.

Salix arbuscula. Jacqu. fl. Austr. t. 408.
Host. Synopf. 527. vix recedit a fruticulo no-
stro; sculptor ibi, ut ipse Jacquinus monet,
folia nimis manifeste ferrata exprimebat.

Salix myrsinites. Schk. bot. Handb.
t. 317. a. f. A. maxime affinis *S. arbusculae*
Jacquini, et sine dubio eadem cum nostra spe-
cies.

Recedit nostra ab Halleri planta stipulis
nullis.

III. Willdenow in Sp. pl. 4. p. 681. sine
dubio sub *S. coruscante* nostram speciem
comprehendebat.

†† *Foliis villosis.*

1513. *SALIX argentea* foliis integerrimis ellipticis subrevolutis mucronulatis acutis, supra subvillosis subtus ramulis sericeo - argenteis. *Smith. fl. brit.* 3. 1059.

Salix argentea foliis ellipticis integerrimis recurvato - acutis, supra pubescentibus subtus sericeo - pilosis argenteis, gemminibus ovato - lanceolatis villosis. *Willd. Linn. Sp. pl.* 4. p. 693.

Salix repens. var. γ. Hudf. fl. Angl. p. 489.

Salix arenaria *Lightf. fl. Scot.* 2. p. 604.

Germ. Silberweide.

Gall. Saule argentée.

Hab. In pascuis arenosis udis montosis et infertioribus passim. In uliginosis sabulosis *auf dem Kaltenbrunn, auf dem Kniebis* et alibi in Sylvae nigrae altioribus sabulosis filiceis cum Vaccinio uliginoso.

Flor. Aprili, Majo, Junio. **Semina** Julio maturantur. **5**

Frutex 2-3-4 pedalis. **Rami** copiosi, diffusifusi, adscendentes, teretes, quandoque obsolete angulati, fusci, pubescentes, in summitate ut et juniores sericei, argentei. **Folia** alterna, breviter petiolata, elliptica, integra, denticulis obsolete quandoque raris a medio ad apicem praedita, crassiuscula, plana, margine subrevoluta, supra fusco-viridia, pubescentia, subtus argentea, sericea, semiunciam ad unciam longa, quatuor ad octo lineas lata, junior saepe majora. **Stipulae** in inferiore parterami

im saepe nullae, aut caducae, in superiore epe ad basin petioli binae, oblongae, acutiusculae, integrae, erectae, argenteo-sericeae, ses lineas longae. *Amenta* alterna, sparsa, breviter pedunculata, ovata, obtusa brevia, *squamis* oblongis, obtulis, villosis, nigricantibus imbricata. *Stamina* duo. *Germen* pedicellatum, inereun, basi ventricosum, superne attenuatum, villosum. *Stigmata* duo, bifida. *Capsula* valiacuminata, fusco-ferruginea, glabra.

b C. Folia in hac specie plana, firma, crassiuscula. Rami juniores semper sericei, argentei.

Nota. Non confundenda cum *S. arenaria* Linnaei, quae est planta mere alpina, vix unquam in Germania obveniens.

Differt a *Salice fusca*, foliis non elliptico-lanceolatis, supra non glabris, ramis junioribus semper argenteo-sericeis, stipulis majoribus.

514. *SALIX repens* foliis integerrimis elliptico-lanceolatis submucronulatis supra nudiusculis, subtus glaucis sericeisve, caule depresso. *Smith. fl. brit.* 3. p. 1061.

Salix repens foliis elliptico-lanceolatis integerrimis acutis subtus subsericeis, capsulis demum glabris, caule depresso exstipulato. *Willd. Linn. Sp. pl.* 4. p. 693.

Salix repens foliis integerrimis lanceolatis utrinque subpilosis, caule repente. *Linn. Sp. pl.* 2. p. 1447.

Salix arenaria. *Poll. fl. pal. n.* 922. (omnino).

Salix depressa foliis integerrimis ovato oblongis supra glabris subtus sericeis. *Hoffm. germ. p.* 341. ej. *Salic. I. p.* 63. — *Roth. fl. germ.* 2. p. 516.

Salix humilis repens. Lobel. Icon. vol. 1. 138. (bona).

Icon. Hoffm. Salic. I. t. 15. 16.

Germ. Kriechende Weide.

Gall. Saule déprimé.

Hab. In montosis praesertim pascuis saluberrimis uliginosis. Auf dem Kaltenbrunn, auf der Herrnwiese, auf dem Kniebis. In Bregenzia in monte Feldberg ad Stagnum Scheuchzeria palustris nuper vidi. In Principatu superiori in pratis udis prope Wangen. In Principatu Fürstbergensi in pratis udis.

Flor. Martio, Aprili, Maio. Semina laeva matura. h

Radix lignosa, crassiuscula, longe lateralis repens, fusco nigricans. Fruticulus depressus. Rami sparsi, procumbentes, quandoque acutiusculi, teretes, fusci, vel ex viridi-flavescentes, digitales, palmares, vix altiores, tenuiter glabri, juniores subpubescentes. Gemmae laeves, rotundae, fusco nigricantes. Folia sparsa, breviter ovata, longe petiolata, vix altiora, tenuiter glabra, juniores subpubescentes.

um attenuatum. *Stigmata* duo, simplicia, a, subrubella. *Capsulae* maturae lutescent rubellae.

ta. Descriptionem dedi ad individua spontanea, in altioribus uliginosis montosis lecta.

Planta pro aetatis et diversi foli natalis ratione maxime polymorpha, praesertim magnitudine folii et forma.

In horto culta in omnibus partibus multo major. ciebibus affinis S. sericea, fusca, incubacea ambigua et rosmarinifolia praesertim ab stipularum dignoscitur.

Hic sine dubio pertinet *Salix incubacea* Roth. erm. 2. p. 514. et quidem ob stipularum absentiam. Vera *S. incubacea* Linnaei stipulis ovatis instructa est.

Salix prostrata foliis subdentatis ellipticis acutius glaucis sericeis, caule prostrato. Smith. fl. p. 3. 1060. in montosis angliae humidis obveniens. sentiam stipularum sine dubio etiam non a *S. repens* Linn. removenda est.

Salix fusca. Schkuhr. bot. Handb. p. 448. tab. 317. c. n. 17. sine dubio ad *S. repens* Linn. pertinet.

5. *SALIX fusca* foliis elliptico-lanceolatis subintegerrimis acutis, supra glabris, subtus sericeo-tomentosis, stipulis lanceolato-subulatis, capsulis lanceolatis pedicellatis sericeis. Willd. Linn. Sp. pl. 4. p. 694.

Salix fusca foliis subdenticulatis elliptico-oblongis acutis supra glabris subtus glaucis sericeis, petiolis attenuatis. Smith. fl. brit. 3. p. 1060.

Salix fusca foliis integerrimis ovatis subtus villosis nitidis. Linn. Sp. pl. 2. p. 1449. — Flor. Lapp. p. 365. t. 8. f. R.



longis, elliptico-lanceolatis, acutatis, supra saturate viridibus, glabris subtus argenteo-sericeis, *Stipulis* lanceolatis acutis.

Obs. Simillima saepe praecedenti specie praesentia stipularum distinguitur.

1516. *SALIX incubacea* foliis convexiusculis integris supra villosis nitidis, stipulis deciduis.

Salix incubacea foliis integerrimis subtus villosis nitidis, fimbriatis acutis. *Linn. Sp. pl. 2. p. 14*

Salix incubacea foliis lanceolatis attenuatis integerrimis coriatis glabris subtus incanis subsericeis lanceolatis sericeis, fimbriatis. *Willd. Linn. Sp. pl. 4*

Salix angustifolia suberecta, f

alix pumila linifolia incana. Zanichell. hist. hist. lit. venet. p. 253. n. 3.

alis pumila angustifolia prima. Clus. hist. plant. p. 85. 86.

n. Zannich. l. c. t. 44.

m. Mattenweide. Schmalblättrige kleine Weide.

l. Saule nicheur.

l. In pascuis pratisque sabulosis humidis altioribus et demissioribus cis et trans Rhenum passim.

r. Martio, Aprili. Semina Majo matura. 5

Radix repens. Frutex sesquipedalis, tripedum, decumbens. Rami adscendentes, erectius- maxime ramulosi. Ramuli fusci aut luteo- urei, glabri. Folia alterna brevissime petio- quandoque subseffilia, lanceolata, integera, subconvexa, angusta 2-3 lineas lata, un- circiter et vix ne longa, supra dilute viridia, ubescentia, nitentia, subtus pubescentia, se- , albida, demum pube fere destituta, incana. nta macula alterna, sparsa, seffilia, initio stunda, demum ex ovali-cylindrica, erecta, unciam vix longa, squamis imbricatis, line- bovatis, atro rubentibus, demum nigrican- , dense glauco-villosis. Stamina duo, squa- ongiora. Antherae didymae, flavae. Amenta linea subseffilia, ovata, demum ovali-cylin- , semiunciam longa. Germina brevissime cellata obverse pyriformia, pubescenti feri- incana. Stylus minutus, incanus. Stigmata crassiuscula, obsolete bifida, saturate rubra. Larium in utroque sexu ovatum, breve, ses- rubellum, bali exteriori germinis sive fila- torum annexum.

tegriusculis lineari lance
tus incanis rariter serice
ari-fetaceis erectis.

Salix rosmarinifolia foliis
cœolato linearibus strict
tus tomentosis. *Linn. Sp.*

Salix rosmarinifolia foliis
latis subintegerrimis pla
centibus subtus sericeis,
cœolatis villosis, stylis e
Linn. Sp. pl. 4. p. 697.

Salix rosmarinifolia procum
sessilibus, lineari-lance
mis, supra saturate virid
dentibus subtus ex decur
ricea albescentibus. *Wulfe*
lect. 3. p. 45.

Salix rosmarinifolia foliis in
ri-lanceolatis strictis sub
erecto, stipulis erectis
brit. 3. p. 1062.

Salix humilis repens angustifolia

monte *Feldberg* ad Stagnum in turfosis; In Principatu superiore ad Lacum Bodamicum in turfosis prope *Bohlingen* et alibi passim.

o r. Majo, Aprili. *Semina* Junio maturæ: 5

Frutex 1-2-3 pedalis, decumbens. *Rami* teretes, fusci, glabri, juniores subvillosi. *Folia* ovata, sparsa, subsessilia; erecta, lineari-lanceolata, integruscula, margine rariter subdenticulata, supra saturate viridia, glabra, subtus incana, pilis adpressis raris munita, sesquiunciam ad binas uncias longa, 3-4 lineas lata, nervata, siccitate margine reflexa. *Stipulae* ad basem petiolorum lineari-setaceae, villosiusculae. *Florae* utriusque sexus simillima *S. incubae*, *antherae* primum rubellae, demum flavae. *Gemmae* obverse pyriformia, brevissime pedicellata, sericea, incana. *Styli* brevissimi, exilissimi. *Stigmata* bina, obsolete bifida, saturate rubra.

f. Planta in horto culta major quatuor pedalis. *Rami* ascendentes, virgati, simpliciusculi. *Folia* breviter petiolata, erecta, lineari-lanceolata, longiora, biuncialia, integruscula; per lentem spectata margine remote obsolete denticulata; supra saturate viridia, glabriuscula, subtus incana, pilis sericeis raris arcte adpressis instructa. *Stipulae* lineari-subulatae, erectae, incano-villosulae. *Styli* in uno eodemque individuo mox brevissimi, mox longiusculi.

18. SALIX aurita foliis obovatis recurvato-acutis apice undulato-ferratis supra viridibus pubescentibus subtus cano-villosis rugoso-venosis, stipulis subsemicordatis dentatis, capsulis lanceolatis pedicellatis, stigmatibus subsessilibus. *Willd.*
Linn. Sp. pl. 4. p. 700.

Salix aurita foliis integerrimis utrinque villo-
sis obovatis appendiculatis. Linn. *Sp.*
pl. 2. p. 1446. — *Poll. fl. pal.* n. 91.

Salix aurita foliis subserratis obovatis vil-
lis, mucronulo adunco, utrinque vil-
lis reticulato-venosis, stipulis polymor-
phis. *Smith. fl. brit.* 3. p. 1064.

Salix foliis rugosis reticulatis integerrimis
inferne tomentosis, julis ovatis. *Hall. stirp.*
helv. n. 1652.

Salix folio rotundo minore. *Dillen. app.* 2.
Icon. Pfleger arb. et frut. Badens. — *Hoffm.*
Salic. t. 22. f. 1. — *Schkuhr. bot. Handb.*
t. 317. a. n. 2.

Germ. Geöhrte Salbeyweide. Kleine Weis-
weide.

Gall. Saule à oreillettes.

Hab. Utrinque ad ripas, prata fossas aquarum
piscinas, et in sylvaticis udis passim fre-
quens.

Flor. Martio, Aprili, Majo. *Semina* Junio
maturat. **5**

Frutex seu arbor 4-6 pedalis, et altior,
erecta, *Salici Capreae* affinis. *Cortex* cinereus
glaber, in adultioribus rimosus. *Rami* copiosi,
patentiusculi, novelli fusci, fusco-rubelli, saepe
subtomentosi. *Folia* alterna, breviter petiolu-
ta, obovata, subrotunda, obtusa vel breviter
acuminata, acumine inflexo, subserrata, sub-
undulata, supra viridia, subpubescentia, rupe-
sula, subtus reticulato-venosa pallidiora, vil-
losa, sesquiunciam ad binas uncias longa, un-
ciam circiter lata. *Stipulae* in basi petiolorum
oppositae, variae, saepe reniformi-semicorda-
tae, repando-dentato-crenulatae. *Amenta* mas-

ula alterna, breviter pedunculata, ovata, ob-
 usa, semiunciam ad unciam longa, squamis im-
 ricato ovato - lanceolatis, obtusis, apice ni-
 ricantibus, villosis. *Stamina* duo. *Filamenta*
 asi subconnata. *Antherae* luteae didymae, sub-
 undae. *Germen* cylindrico-lanceolatum, pe-
 icellatum, sericeum. *Stylus* subnullus. *Stig-*
ata duo, conniventia. *Nectarium* utriusque
 us cylindricum, obtusum, breve.

519. *SALIX aquatica* foliis obovato-el-
 lipticis subserratis acutis supra viridibus
 pubescentibus subtus glaucis pilosiuscu-
 lis, stipulis rotundatis dentatis, germi-
 nibus pedicellatis lanceolatis villosis,
 stigmatibus sessilibus. *Willd. Linn. Sp.*
pl. 4. p. 701.

Salix aquatica foliis subserratis obovato-el-
 lipticis pubescentibus planis subtus glau-
 cescentibus, stipulis rotundatis dentatis.
Smith. fl. brit. 3. p. 1065.

Salix cinerea Wetterauische fl. 3. p. 414. an?

Salix aurita. Hoffm. Salic. 1. t. 5. f. 3.

Salix folio ex rotunditate acuminato. Bauh.
pin. 474. — Raj. Synops. 449.

erm. Wasserweide.

all. Saule aquatique.

a b. Cum praecedente in humidis ad sepes et
 dumeta.

Hor. Cum praecedente. 5

Nimis omnino affinis praecedenti speciei a
 a mere differre videtur, *Statura* humiliori,
 amis copiosioribus erectis, *Foliis* obovatis,

pubescentibus subtus glauco-toment
celatis, stipulis obsoletis, capsulis l
latis pubescentibus, stigmatibus sub
Linn. Sp. pl. 4. p. 702. — Smit
1068 a *S. caprea* vix separanda est
Salic. 1. p. 28. t. 5. f. 4. et t. 21. f
celatam Willd. et Smithii sub
preae comprehendebat.

1520. *SALIX caprea* foliis
subtus tomentosis undatis
culatis. Linn. Sp. pl. 2. p.
fl. pal. n. 923.

Salix caprea foliis ovatis ac
tis undulatis subtus toment
sublunatis, capsulis ven
fl. brit. 3. p. 1067.

Salix foliis ovatis rugosis sub
tomentosis, julis ovatis, l
caulibus ferratis. Hall. fl.

Salix latifolia rotunda. Bau
Mapp. fl. Alf. p. 276.

Salix caprea latifolia Tabern

11. *Saule Marceau.*

12. Utrunque in dumetis humidiusculis sylvisque passim abunde.

13. Aprili, Majo. *Semina* Junio maturat. 14

Frutex quandoque *arbor* mediocris. *Corticinereus*, glaber, in adultis rimosus. *Rami* erni patenti-erecti, cinereo-virides, subflavescentes, aut fusci, glabri, juniores villosi. *Folia* alterna, petiolata, ovata, ovali-subrotunda, oblonga, quandoque ovato lanceolata, undulata, crenato-dentata, quandoque integruscula, supra saturate viridia, glabriuscula, pubescentia, subtus villosa-tomentosa, glau-incana, rugoso-reticulata. *Stipulae* in raris praesertim junioribus ad petiolorum basin, junctae, subrotundae, subulnatae, cordatae, semicordatae, crenulatae, subtus villosae, glabrae. *Amenta mascula* alterna, pedunculata, oblongo cylindrica, sesquiuncialia et longiora, integruscula, erecta. *Squamae* imbricatae, oblongae, nigricantes, villosae. *Nectarium* cylindricum, viridulum. *Stamina* duo. *Antherae* ovatae, didymae, luteae. *Amenta feminina* alterna, petiolata, cylindrica, duas uncias longa, glabra. *Squamae* lanceolatae, obtusae, nigricantes, villosae. *Germen* pedicellatum, lanceolatum, pubescens. *Styli* brevissimi. *Stigmata* bina, lutescentia.

Nota. Pro diversa aetate et diverso solo quammaxime polymorpha ratione magnitudinis, et foliorum forma. Folia ovali-elliptica, basi plus minusve rotundata, superne plus minusve acuta, a basi ad medium margine plus minusve undulata, a medio ad apicem subsinuato-denticulata, quandoque integruscula, supra saturate viridia, glabra, saepe etiam plus minusve pubescentia, subtus venis reticulata, villosa-tomentosa, incana 1½-2-3-5 uncias longa, 1-3 uncias lata,

omnium facile latissima. Stipulae in ramis senosioribus saepe nullae, in junioribus oppositae, fere semicordatae, rotundae, aut semilunatae, plus minusve acutae, denticulatae, longitudine super petiolorum.

Amenta masculina florifera grate odorata, apice acceptissima.

1521. SALIX acuminata foliis oblongo-ovatis acuminatis apice obtuse dentatis subtus rugosis tomentosis. Willd. arb. p. 346.

Salix acuminata foliis lanceolato-oblongis, acuminatis undulatis, dentatis subtus tomentosis, stipulis reniformibus, capsulis ovato-fusculatis. *Smith. fl. brit.* 3. p. 106.

Salix acuminata foliis ovato-oblongis subtus tomentosis superioribus integris, inferioribus crenatis. *Hoffm. Salic.* 1. p. 47.
— *Roth. fl. germ.* 2. p. 519.

Salix foliis oblongo-ovatis acuminatis rugosis subtus tomentosis. *Du Roi Harb.* 2. p. 407.

Salix caprea acuto longoque folio. *Rej. Synops.* 450.

Icon. Hoffm. Salic. 1. t. 6. et 22. f. 2. *Schkuhr. bot. Handb.* t. 317. c. n. 12.

Germ. Gemeine Werstweide. Grofse Werstweide.

Gall. Saule pointu.

Hab. Utrinque passim cum praecedente specie praesertim in regionibus rhenanis frequens.

Flor. et Semina cum praecedente maturat h

Affinis praecedenti, a qua autem differt, *Foliis* ovato-oblongis, acuminatis, angustior-

is, a basi ad medium integris, dehinc ad apicem
 m dendato - ferrulatis, *Stipulis* subreniformi-
 micordatis, falcatis, *Amentis* masculis sublon-
 oribus, *femineis* masculis subfimilibus, *Ger-*
ine longius pedicellato, conico, albide feri-
 to, *Stylo* brevi.

Flores quammaxime melliferi.

Lignum hoc cum praecedente carbonibus aptum
 pulvere pyrio miscendo, nec non pictorium in
 sm.

522. *SALIX viminalis* foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis, ramis virgatis. *Linn. Sp. pl.* 2. p. 1448. — *Poll. fl. pal. n.* 624.

Salix viminalis foliis subintegerrimis lineari-lanceolatis longissimis subtus albo-sericeis, pilis squamarum amenti feminei germinae longioribus. *Willd. Linn. Sp. pl.* 4. p. 706.

Salix viminalis foliis lanceolato-linearibus longissimis acuminatis integerrimis subtus sericeis, ramis virgatis, stylo elongato. *Smith. fl. brit.* 3. p. 1070.

Salix foliis praelongis obscure dentatis subtus tomentosis et albicantibus. *Hall. stirp. helv. n.* 1641.

Salix angustis et longissimis foliis crispis subtus albicantibus. *Bauh. hist.* 1. p. 212.

con. Pfleger arb. et frut. Badens. — *Hoffm. Salic.* 1. t. 2. 5. et 21. f. c. f. g. — *Schkuhr. bot. Handb.* t. 317. b. n. 4.

erm. Bandweide. Korbweide.

all. Saule à longues feuilles.

Hab. Utrinque passim ad fossas aquatiles, in pratis et sylvis aquosis frequens.

Flor. Aprili, Majo. **Semina** Junio matura.

Frutex saepe arbor sat alta. **Rami** virgati, longi, fusco-flavicantes, virides, glabri, novelli pubescentes. **Folia** alternata, linearilanceolata, acuminata, integra, aut subserrulata, praelonga, breviora et longiora, subundulata, supra viridia, glabriuscula, subtus sericea, tenella seu juniora margine revoluta et villosa. **Stipulae** praesertim in ramis junioribus ad petiolorum basin, linearilanceolatae, integrae. **Amenta** alterna, pedunculata, erecta, oblonga sesquiunciam fere longiora, bracteata. **Squamis** obovatis, obtusis, pilosis. **Stamina** duo. **Filamenta** basi coalita. **Anthodymae**, subrotundae, luteae, apice sessiles. **Germen** sessile, ovalilanceolatum, mucosum. **Stylus** filiformis, elongatus, bifidus. **Capsula** ovali-subulata.



presso - pilosis viridibus, pilis squamarum amenti feminei longitudine stigmatatis. *Willd. Linn. Sp. pl. 4. p. 707.*

Salix mollissima foliis lanceolato-linearibus acuminatis subintegerrimis subtus nudiusculis viridibus, amenti feminei squamis pilosis, pilis longitudine pistilli, germine sericeo, stylis supra medium conglutinatis indivisis. *Ehrh. Beitr. 6. p. 101.*
Salix foliis lineari-lanceolatis utrinque tenuissime tomentosis mollissimis. *Hoffm. germ. p. 343.*

Fr m. Weichblättrige Weide.

Fl l. Saule molle.

Fr b. In dumetis et sepibus humidis ad aquarum ripas. In Principatu superiore ad Lacum Bodamicum. In Principatu Fürstembergenensi ad fontes Danubii et alibi.

Fr o r. Martio, Aprili. *Semina* Junio matura. *Fr*

Affinis praecedenti, a qua differt, *Virgis* mollissime pubescentibus tenuioribus, *Foliis* cinque viridibus, mollissime pilosis, brevioribus, et *Pilis* squamarum Stylorum longitudine.

524. *SALIX alba* foliis lanceolatis acuminatis ferratis utrinque pubescentibus: ferraturis infimis glandulosis. *Linn. Sp. pl. 2. p. 1049. — Poll. fl. pal. n. 925.*

Salix alba foliis lanceolatis acuminatis ferratis utrinque sericeis, ferraturis infimis glandulosis, stigmatibus bipartitis. *Smith. fl. brit. 3. p. 1071.*

Salix arborea foliis ellipticis lanceolatis subtus sericeis, dentibus crassescens. *Hall. stirp. helv. n. 1635.*

Salix vulgaris alba arborescens. Bauh. pin.
473. — *Mapp. fl. Alf. p.* 274.

Icon. Pfleger arb. et frut. Badenf. — Hofm.

Salic. 1. t. 7. 8. et 24. f. 3. — *Schkuhrht.*

Handb. t. 317. b. n. 7. — *Blackw. herb.*

327.

Germ. Weisse Weide.

Gall. Saule blanc.

Hab. Utrunque passim ad pagos et urbes, in
pascuis paludosis rhenanis frequens.

Flor. Martio, Aprili. *Semina* Majo, Junio ma-
tura. h

Arbor procera. *Cortex* rimosus, cinereus.
Rami alterni, patenti-erecti, fusco-virentes,
superne subpubescentes. *Folia* alterna, petio-
lata, acuminata, argute ferrulata: *ferraturae*
glandulosae, fuscae, supra pubescentia, subtus
sericea, albida. *Stipulae* nullae, aut minutae.

Amenta alterna, sparsa, pedunculata cylindrica,
sesquiunicam ad binas uncias longa. *Squamae*
ovali-lanceolatae, acutae, virides, pubescentes.
Nectarium in masculis duplex; *anterior*
minimum, ovatum, obtusum; *posterior* linearis,
tenue, anteriori duplo longius, filamentis pos-
sice adnatum; in femineis unicum subcylindri-
cum, compressum, emarginatum ad basin squa-
mulae. *Stamina* duo. *Filamenta* basi connata.
Antherae subrotundae, didymae, luteae. *Germen*
sessile, ovali-oblongum. *Stylus* brevis, bifidus. *Stigmata* duo bipartita.

Pharm. SALICIS ALBAE Cortex.

Arbor inter salices maxima pulchra, magis in pa-
cuis rheno vicinis inundationibus expositis propaganda.
Pascua ad pagos et urbes mire condecorat.

In annosioribus truncis saepe *Boletus sat-
veolens*.

iger, Sanddorn.

vide.

superiore in sabulosis
ken; nec non prope
 cipatu superiore in
adamicum passim
 um prope *Oth-*
Rhenum prae-

is lai.

mo, spicis a ae. 5

1451. — *Poll. ju.* pedalis

.. *stirp. helv. n. 1609.* filia,

baccis albis. *Bauh. pin. 423.* — *Ma* pl. 75

Als. p. 329.

um. *Tabernaem. 1376. Icon.*

Pfleger arb. et frut. Bad. Blackw. t. 184.

• *Mistel. Kleister.*

• *Gui commun.*

Utrunque in variis arboribus parasitica.

• Martio, Aprili. *Baccae* Novembri, De-
 cembri maturae. 5

Fruticulus elegans, sempervirens, in ra-
 rborum omnis generis crescens, a basi sta-
 amosus, dichotomus. *Folia* sessilia, oppo-
 lanceolato-cuneiformia, integerrima, cras-
 sa, ex luteo virescentia. *Flores* lutescen-
 parvi. *Baccae* axillares, sessiles, albae, glo-
 nitidae, massa tenaci, glutinosa turgidae.

rm. Visci Lignum; Baccae.

in Epilepsia aliisque morbis convulsivis, a qui-
 n ita commendatur ut Viscum in quercubus nas-
 raeferatur.

Erica baccifera procumbens nigra
pin. 436.

Erica baccifera tenuifolia. Tabern.
Icon.

Icon. Oeder. fl. Dan. t. 975. — Pfl.
Badenf.

Germ. Schwarze Raufschbeere.

Gall. Camarine à fruits noirs.

Hab. In montosis uliginosis turfosis
repletis auf dem Kaltenbrunn bei
Hornsee et den Oelachen frequ
der Herrnwiese in turfosis am a
neten See; in monte Kniebis an de
denschanze.

Flor. Aprili, Majo. Baccae Julio,
maturae. h

Fruticulus pedalis, sesquipedalis
procumbens. Folia numerosa, subpetio
quina, linearia, margine reflexa, int
ma, patentia, glabra, superne saturati
pallide viridia. Flores axillares, sess
tarii. Corollae purpureae. Filamenta p



Opposita. *Calyx* tetraphyllus. *Corolla* nulla. *Ger-*
men oblongum, trigonum, margine obsolete
quadrifido coronatum, inferum. *Stylus* nullus.
Stigma obtusum, vix marginatum. *Bacca* glo-
 -osa, unilocularis, laevis. *Semen* unicum, cor-
 -atum, compressum, obtusum, carnosum.

526. VISCUM foliis lanceolatis, obtusis,
 caule dichotomo, spicis axillaribus. *Linn.*
Sp. pl. 2. p. 1451. — *Poll. fl. pal. n.* 926.
 — *Hall. stirp. helv. n.* 1609.

Viscum baccis albis. *Bauh. pin.* 423. — *Mapp.*
fl. Als. p. 329.

Viscum. *Tabernaem.* 1376. *Icon.*

Icon. *Pfleger arb. et frut. Bad. Blackw. t.* 184.

Ger. Mistel. *Kleister.*

Hall. *Gui commun.*

Tab. Utrunque in variis arboribus parasitica.

Flor. Martio, Aprili. *Bacca* Novembri, De-
 cembri maturae. *h*

Fruticulus elegans, sempervirens, in ra-
 -mis arborum omnis generis crescens, a basi sta-
 -tim ramosus, dichotomus. *Folia* sessilia, oppo-
 -sita, lanceolato-cuneiformia, integerrima, cras-
 -sifuscula, ex luteo virescentia. *Flores* lutescen-
 -tes, parvi. *Bacca* axillares, sessiles, albae, glo-
 -bosae, nitidae, massa tenaci, glutinosa turgidae.

Pharm. Visci Lignum; Baccae.

In Epilepsia aliisque morbis convulsivis, a qui-
 busdam ita commendatur ut *Viscum* in quercubus nas-
 -cens praefertur.

Linn. Gen. pl. ed. Schreb.

Mas. Calyx monophyllus, l
niis subrotundis, concavis, apic
tibus, lateribus dehiscentibus.

Femina Calyx monophyllus, ov
tubulatus, clavatus, ore bifido,
rolla nulla. Stigma crassiusculu
erectum, calyce duplo longius.
bosa, unilocularis. Semen unicu

1527. HIPPOPHAE *rhama*
lauceolatis. Linn. Sp. pl. 2

Hippophaë foliis linearibus,
nolis. Hall. stirp. helv. n.

Osyris rhamnoides. Scop. Carn

Rhamnoides fructifera, foliis
leviter flavescens. Mo
268.

Rhamnus falicis folio fruct
Bauh. pin. 477.

erm. Weidenartiger Sanddorn.

all. Griset rhamnoide.

ab. In Margraviatu superiore in fabulosis ad Rhenum prope *Zienken*; nec non prope *Kleinhüningen*. In Principatu superiore in fabulosis ad Lacum *Bodamicum* passim abunde. In Alsatia ad Rhenum prope *Othmarsheim* et alibi passim ad Rhenum praefertim superiorem.

or. Majo. *Bacca* Novembri maturae. *B*

Arbor seu *Frutex* elegans, 3-4-9 pedalis altior, erectus. *Folia* alterna, subseffilia, ceari-lanceolata, integerrima, superne palliviridia, inferne argentea, squamulis villosis latis nitentibus, pulcherrimis adspersa. *Spiculae* rectae, nitidae, acutae, ferrugineae. *Flores* laterales, axillares, seffiles, fasciculati, rudinosi. *Bacca* subglobosae, flavescentes, acidae, ingratae.

Nota. Inter flores masculos interdum hermaphroditos observavit Cel. Viborg.

Frutex pulcher ad littora fabulosa humida condensatus.

f. In Hortis Carlsruhanis adhuc hospitatur *HIPPOPHAE canadensis* Linn. foliis latioribus et brevioribus instructa.

Porro *MYRICA* Gale, et cerifera Linn. ad eam Classem pertinentia; nec non *XANTHOXYLUM* a *va* Herculis.

DIOECIA PENTANDRIA.

CCCCXLIV. SPINACIA.

Linn. Gen. pl. ed. Schreb. n. 1520.

Mas. Calyx quinquepartitus: *laciniis* convivis, oblongis, obtusis. *Corolla* nulla. *Femina*
Flora Bed. Tom. III, B b b

Calyx monophyllus, quadrifidus acutus: laciniis duabus oppositis minimis, persistens. Corolla nulla. *Styli* quatuor, capillares. *Stigmata* simplicia. *Calyx* seminiferus induratus, coalitus. *Men* unicum, subrotundum, calycē tectum.

1528. *SPINACIA spinosa* foliis sagittatis, fructibus sessilibus, spinosis.

Spinacia oleracea. Linn. *Sp. pl.* 2. p. 1456. var. α.

Spinacia foliis sagittatis, seminibus aculeatis. Mill. *dict. n.* 1.

Lapathum hortense f. *Spinacia* semine spinoso. Bauh. *pin.* 114.

Spinachia. Tabernaem. p. 809. Icon.

Icon. Blackw. t. 49. — Regnault. bot.

Germ. Spinat mit gehörntem Saamen. Winter-Spinat.

Gall. Epinard cornu.

Hab. Colitur utrinque in oleraceis, agris et vineis passim.

Flor. Majo, Junio. Semina Julio matura. ♂

Ufus plantae culinaris notissimus.

1529. *SPINACIA inermis* foliis oblongo-ovatis, seminibus sessilibus inermibus.

Spinacia oleracea. Linn. β.

Spinacia glabra foliis oblongo ovatis, seminibus glabris. Mill. *Dict. n.* 2.

Lapathum hortense, f. *Spinacia* semine non spinoso. Bauh. *pin.* 115.

Spinacia nobilis. Tabernaemont. p. 809. Icon.

erm. Dornlofer Spinat. Sommerspinat. Holländer Spinat.

all. Epinard sans cornes. Epinard de Hollande.

ab. Colitur cum praecedente.

lor. Majo, Junio. Semina Augusto matura. ♂

Usus ut in praecedente specie culinaris, sed magis omnino tenera planta, specie diversa.

CCCCXLV. C A N N A B I S.

Linn Gen. plant. ed. Schreb. n. 1522.

Mas. Calyx quinquepartitus: foliis oblongis acuminato-obtusis, concavis. Corolla nulla. Antherae tetragonae. Femina. Calyx monophyllus, oblongus, acuminatus, latere altero longitudinaliter dehiscens, persistens. Corolla nulla. Styli duo, subulati, longi. Stigmata acuta. Capsula seminiferus arcte clausus. Semen nux globose-depressa, bivalvis.

530. CANNABIS *fativa* foliis digitatis.

Linn. Sp. pl. 2. p. 1457.

Cannabis fativa. Bauh. pin. 326. ♀

Cannabis erratica. Bauh. pin. 320. ♂

Cannabis. Tabernaem. p. 936. Icon.

Leon. Blackw. t. 322. — Regnault. bot.

Serm. Hanf. planta mascula nostratibus vulgo Fimmel. foeminea Maskel. (vox corrupta contraria).

Gall. Chanvre cultivée.

Hak. In Persia. Colitur utrinque passim abunde.

Flor. Majo, Junio, Julio. **Semina** Octobri
tura. ☉

In regionibus rhenanis praesertim Hassi-
sibus, et Hochbergensibus cultura dicitur
cannabis ita ut singuli caules altitudinem 10-12 pes
atingant, quin superent.

Obs. Annis pluviosis praesertim individua feminina
re saepe emaciantur planta parasitica, Oroban-
chae ramosa Linn. Hausto d. nostratibus, quae
dices Cannabis occupat.

Pharm. CANNABIS Semina.

Oleum viride ex feminibus expressum obtinetur
ex quo praecipue Sapo viridis paratur.

Plantae masculae teneriores tribus circiter hepti-
matibus prius evelluntur femineis. Collectis plantis
in fasces viuctis curandum, ut gluten, quod fibras
tenuis jungit, destruat, quod maceratione (in pluri-
regionibus,) in aqua stagnante fit. Stagna cannabi-
plenda, omnino a pagis et viis publicis remove-
re ob effluvia foetidissima.

Usum Cannabis multifarium, quidem sed no-
tum praetermitto.

Planta recens narcotica.



uctiferus basi semen includens. Semen subrotundum, tunica obtectum.

531. **HUMULUS** *Lupulus* caule volubili, anguloso scabro, foliis oppositis petiolatis cordatis tri-quinquelobis, serratis, scabris.

Humulus Lupulus. Linn. Sp. pl. 2. p. 1457. — Poll. fl. pal. n. 927.

Lupulus. Hall. stirp. helv. n. 1618.

Lupulus mas et foemina. Bauh. pin. 298. — Tabernaem. p. 1293. Icon. — Mapp. fl. Alf. p. 181.

con. Regnault. bot. — Blackw. t. 536.

verm. Hopfen. Gemeiner Hopfen.

Wall. Houblon.

Tab. Utrisque passim ad sepes et viarum marginalia dumetosa.

Flor. Junio, Julio, Augusto. Semina Novembri, Octobri matura. 7

Planta notissima passim, cum primis in Palatinatu colitur.

harm. LUPULI Coni Turiones.

Turiones primo vere loco Asparagi comeduntur.

Coni s. Strobili Lupuli Cerevisiae gratam amaritudinem conferunt. hisdem ab acore defenditur. Tonicas, anarcoticas et diureticas vires habent.

Flores apibus mel praebent.

DIOECIA HEXANDRIA.

CCCCXLVII. T A M U S.

Linn. Gen. plant. ed. Schreb. n. 1527.

Mas. Calyx sexpartitus: foliolis ovato-lanceolatis, superne patentioribus. Corolla nulla.

Femina. Calyx monophyllus, sexpartitus, campanulatus, patens: laciniis lanceolatis; superus, deciduus. Corolla nulla. Nectaria: Punctum oblongum, singulae lacinae calycinae ad basin interne adnatum. Germen inferum, ovali-oblongum, magnum, glabrum. Stylus cylindraceus. Stigmata tria, reflexa, emarginata, acuta. Baccae ovatae, trilocularis. Semina bina, globosa.

1532. TAMUS communis foliis cordatis indivisis. Linn. Sp. pl. 2. p. 1458. — Hall. stirp. helv. n. 1620.

Bryonia laevis f. nigra racemosa. ♂ Bauh. pin. 297.

Bryonia laevis f. nigra baccifera. ♀ Bauh. pin. 297.

Vitis nigra l. Tabernaem. p. 1288. Icon.

Icon. Miller. illustr. — Blackw. t. 457.

Germ. Glatte Wald-Bryonie.

Gall. Tamme commune. Sceau de notre Dame.

Hab. In Rheni vicinia ad sylvarum marginalia, in dumetis passim prope Eckenstein, Liehenheim et alibi rarius. In Margraviatu superiori in sylva Kündel et Wolsaker prope Mullheim. In Ditione Basileensi, in monte Muteto, in sylva die Hardt, et in dumetis circa Friedlingen. In Alsatia prope Argentinam passim in sylvis versus Rhenum.

Flor. Majo, Junio. Baccae Augusto, Septembri maturae. ♀

Radix subrotunda crassa, extus nigra, intus alba, viscida. Caules ex radice plures, 4-5-6 pedales, et altiores, volubiles, scandentes, debiles, glaberrimi, laete virides. Folia alterna, petiolata, cordata, acuminata, integra, glaberrima,

ida, tenera, ad basin glandulis duobus pellulis instructa. *Flores* mares laterales, racemosi, *ninei* ramosi, parvuli, omnes ex viridi flavicantes. *Baccae* succulentae, subrotundae, rubrae.

arm. TAMNI f. *BRYONIAE NIGRAE Radix*.

Radix suspecta, drastica, acris, hodie in medicina exoleta.

Nota. Ad rariores Florae cives pertinet, a germaniae Floristis hucusque omissa.

DIOECIA OCTANDRIA.

CCCCXLVIII. P O P U L U S.

Linn. Gen. pl. ed. Schreb. n. 1531.

Mas. Amentum oblongum, laxè imbricatum, cylindricum, constans squamis unisloris, longis, planis, margine laceris. *Corolla* Petala nulla. *Nectarium* monophyllum, inferne tubulatum, turbinatum, superne oblique in ovatum limbum terminatum. *Femina. Amentum* ut in mare. *Squamae* ut in mare. *Corolla* Petala nulla. *Nectaria* ut in mare. *Stigma* quadrifidum. *Spula* ovata, monocularis, bivalvis: *valvulis* reflexis. *Semina* numerosa, ovata, *Pappo* pillari volatilia.

533. *POPULUS alba* foliis subrotundis dentato-angulatis, subtus tomentosis. *Lin. Sp. pl. 2. p. 1463.* — *Poll. fl. pal. n. 928.* *Hall. stirp. helv. n. 1634.*

Populus alba foliis cordato-subrotundis lobatis dentatis subtus tomentoso-niveis, amentis ovatis. *Smith. fl. brit. 3. p. 1079.*

Populus alba majoribus foliis. *Bauh. pin.* 439.
Mapp. fl. Als. p. 247.

Populus alba. *Tabernaemont. p.* 1392. *Icon.*

Icon. Psleger arb. et frut. Badenf. — Blam.
t. 548.

Germ. Silber-Pappelbaum.

Gall. Peuplier blanc.

Hab. Utrique in dumetis et sylvis humidis
et transrhenanis, ad rivos et pagos passim
frequens.

Flor. Martio, Aprili. *Semina* Majo, Junio
tura. 5

Frutex seu arbor variae magnitudinis. *Folia*
alterna, petiolata, cordata, inordinate tri vel
quineloba, angulata, dentata, inferne tomentis
niveo obducta, superne obscure viridia hirsuta
seu glabriuscula, *Amentum* ovatum. *Stamina*
octo.

Arbor pulcherrima, saepe procera cujus folia venti
agitata omnia eminus argentum mentiuntur. Pulcherrime
hujus arbores in Horto Carlsruhano.

Gemmae balsamicae, hinc inde sub nomine *Opobalsami germanici* prostrant.

1534 *POPULUS canescens* foliis subrotundis angulato-repandis dentatis subtus tomentoso-incanis, amentis cylindraceis laxis. *Smith. fl. brit.* 3. p. 1080. — *Willd. Linn. Sp. pl.* 4. p. 802. — *Lamarck et Decand. fl. franc.* 3. p. 299.

Populus alba β. *Lamarck fl. franc.*
2. p. 235.

Populus alba folio minore. *Bauh. hist.* 1. p. 160.

☉ *n. Lobel. Icon. 2. p. 193. f. 2.*

☿ *m. Weisgraue Pappel.*

♂ *ll. Peuplier grisâtre.*

♂ *b. Passim in dumetis, locis praesertim humidis. Inter Carlsruhe et Mühlberg.*

☉ *r. Martio, Aprili. Semina Majo, Junio maturat. 5*

Affinis praecedenti speciei a qua differt, *Stamina* minore, *Foliis* multo minoribus, subrotundis non lobatis, minus dentatis, subtus non argenteo-niveis, sed griseo tomentosis, *Amentis* majoribus, laxioribus, non ovatis, sed cylindricis.

535. *POPULUS nigra* foliis deltoidibus acuminatis serratis. *Linn. Sp. pl. 2. p. 1464.*
Pollich. fl. pal. n. 930.

Populus foliis glabris cordato-rhomboideis serratis. *Hall. stirp. helv. n. 1632.*

Populus nigra. Bauh. pin. 429. — Tabernaem. 1392. Icon. — Mapp. fl. Als. p. 248.

☉ *n. Pfleger arb. Badens.*

☿ *m. Schwarze Pappel. Schwarze Espe.*

♂ *ll. Peuplier noir.*

♂ *b. Utrinque in sylvis praesertim demissioribus rhenanis et pascuis inundatis.*

☉ *r. Februario, Martio. Semina Majo maturat. 5*

Frutex saepe arbor variae magnitudinis. *Ramuli* teretes, glabri. *Folia* alterna, longe petiolata subcordato-subrhomboidea acuminata, serrata, glabra, laete viridia. *Stamina* 16-18-21.

☿ *m. POPULI Oculi f. Gemmae.*



Populus fastigiata. Lamarc
franc. 3. p. 300.

Icon. Pfleger arb. Badens.

Germ. Italiänische Pappel. Los
pel. Pyramiden - Pappel.

Gall. Peuplier d'Italie ou de La
lier pyramidal.

Hab. In Italia. Pulcherrima et s
bulacra inter Durlach et
non inter Raftadt et Baden
iam in Palatinatu prope Schu
nec non in Alsatia prope Al
bi ad ambulacra.

Flor. Februario, Martio.

Arbor pulcherrima, introducta
num Ducem Badensem, CAROLUM I
nis 1759 - 1760.

Amat sabulosa humidiuscula, fre
nibus rhenanis humidis, inundationil
vaganda. ubi citissime in arborem

Populus foliis glabris orbiculatis acuminatis, laxe ferratis. *Hall. stirp. helv. n. 1633.*

Populus tremula. *Bauh. pin. 429. — Mapp. fl. Alf. p. 248.*

Populus Lybica. *Tabernaem. p. 1392. Icon. n. Pfleger arb. Badensf.*

m. Espe. Zitterespe. Zitterpappel.

ll. Peuplier tremble.

b. Utrinque in sylvis et dumetosis passim frequens.

r. Februario, Martio. Semina Majo matura. b

Frutex seu arbor variae magnitudinis. *Fol-*
alterna, petiolata, subrotunda, acutiuscula,
usa, sinuato-dentata, glabra, laete viridia,
tus reticulato-venosa. *Petioles* longi, biglan-
osi ad basin folii compressi, debiles; hinc folia
le trementia. *Stamina* octo.

Folia juniora ovibus et capris accepta.

Flores melliferi.

Lignum Populi album laeve.

ota. In hortis Carlsruhanis et Schwezingensibus
praesertim adhuc sub dio hospitantur. *POPULUS*
balsamifera. Linn. *P. candicans.* Ait.
P. canadensis. *P. heterophylla* Linn.
P. graeca. *P. cordata.* et *P. grandiden-*
tata.

DIOECIA ENNEANDRIA.

CCCXLIX. MERCURIALIS.

Linn. Gen. pl. ed. Schreb. n. 1534.

Mas. Calyx tripartitus: laciniis ovato-lan-
latis, concavis, patentibus. *Corolla* nulla.

Filamenta 9-12 capillaria. *Antherae* glaberrimae. *Femina*. *Calyx* ut in mare. *Corolla*. *Nectaria*: Acumina duo, subulata ad apicem, lumen latius germinis singula, sulco germinis compressa. *Germen* hispidum, compressum. *Stylus* reflexi, cornuti, hispidi. *Stigmata* acuta. *Capsula* subrotunda, scrotiformis, didyma, locularis. *Semina* solitaria, subrotunda.

1538. *MERCURIALIS perennis*

simplicissimo, foliis scabris. *Lin.*
2. p. 1465. — *Poll. fl. pal. n. 93.*

Mercurialis caule simplici, foliis ovatis, serratis, hirsutis. *Hall. stirp. helv.*

Mercurialis montana spicata. (♂) *Boiss.*
122. — *Mapp. fl. Alf. p. 194.*

Mercurialis montana testiculata. (♀)
pin. 122. — *Mapp. fl. Alf. p. 194.*

Mercurialis sylvestris mas et foemina.

*Radix repens. Caulis palmaris, pedalis et
erectus, sulcatus, scaber, simplex. Fo-
lia opposita, petiolata, ovato-lanceolata, ser-
iceo villosa, scabriuscula, obscure viridia. Sti-
les duae, acutae, parvae, ad petiolorum ba-
ses. Flores axillares, spicati, pedunculati.*

Planta virosa, suspecta, narcotica.

59. MERCURIALIS annua caule bra-
chiato, foliis glabris, floribus spicatis. *Lin.
Sp. pl. 2. p. 1466. — Poll. fl. pal. n. 932.*

*Mercurialis annua brachiata, foliis conjuga-
tis, ovato-lanceolatis glabris. Hall. stirp.
helv. n. 1600.*

Mercurialis spicata f. foemina, dioscoridis et
Plinii. (♂) *Bauh. pin. 121. — Mapp. fl.
Alf. p. 195.*

*Mercurialis testiculata, five mas, Dioscoridis
et Plinii. Bauh. pin. 121. Mapp. l. c.*

*Mercurialis mas et foemina. Tabernaemont.
p. 941. Icon.*

*n. Curtis. fl. Lond. t. 335. — Bulliard. herb.
t. 159. 235. ♂ et ♀ Blackw. t. 162.*

*m. Bingelkraut. Einjähriges Bingelkraut.
41. Mercuriale annuelle.*

*b. Utrinque passim in oleraceis, agris, vineis
et ruderatis,*

*r. Julio, Augusto, Septembri. Semina Oc-
tobri, Novembri matura. ☉*

*Radix cylindrica, fibrosa, descendens, alba.
Caulis pedalis, sesquipedalis et altior, erectus,
angulatus, glaber, striatus, ramosus. Rami op-
positi, brachiati, superne foliosi. Folia opposita,
petiolata, ovata, obtusiuscula, obtuse serrata,*

glabra, sublucida, nervosa. *Stipulae* quatuor genicula, lanceolatae, erectae, submenstruae, parvae. *Pedunculi* oppositi, longi, tetragoni, erecti. *Flores* spicati, nudi, res remoti, superiores conferti, conglomerati.

Pharm. MERCURIALIS Herba.

Planta foetidiuscula, amariuscule, nauseosa, suspecta.

Syrupum longae vitae Zwingeri ingratum.

Obs. Saepius in pharmacis confunditur ob similitudinem tanices ignorantiam cum *CHENOPODIO* rubro et albo, immo cum *IMPATIENTE* solis genere, ut ante aliquot annos vidi.

CCCCL HYDROCHARIS

Linn. Gen. pl. ed Schreb. n. 153.

Mas. Calyx Spatha triflora, diphylla longa. *Calyx* proprius triphyllus: foliolis oblongis, concavis, margine membranaceo. *Corolla* petala tria, subrotunda, plana, magnitudine novem in tres ordines disposita.

DIOECIA ENNEANDEIA. 767

Hydrocharis Morsus ranae. Linn. Sp. pl. 2. p. 1466. — Poll. fl. pal. n. 933. — Hall. stirp. helv. u. 1068.

Nymphaea alba minima. Bauh. pin. 193.

Morsus ranae foliis circinatis, floribus albis. Mapp. fl. Als. p. 199.

Ranae morsus. Tabernaem. p. 1131. Icon. on. Curtis. fl. Lond. t. 167. — Oeder. fl. Dan. t. 878. — Schkuhr. bot. handb. t. 333.

erm. Froschbifs.

all. Morène grenovillete,

ab. Utrinque in aquis rhenanis stagnantibus fossis aquaticis et rivulis segnioribus passim frequens.

or. Julio. Semina Augusto, Septembri matura. 7

Caules longi, radicati, funiformes, sub aqua penses, geniculati. Folia congesta, longe petiolata, orbiculato-reniformia, integerrima, plani, glabra, splendentia, superne obscure viridia, inferne rubella, supra aquam natantia, submersa. Flores mares pedunculati, 2-3-4. Fecundi solitarii. Petala sat magna, nivea, unguiculata. Stamina 9-12.

Nota. Ex Dioecia Dodecandria hospitantur in hortis Carlsruhanis MENISPERMUM virginicum et canadense Linn.

DIOECIA MONADELPHIA.

CCCCLI. JUNIPERUS.

Linn. Gen. pl. ed. Schreb. n. 1552.

Mas. Amentum conicum, singulus flos proli agnoscit Squamam latam, brevem, incur-

bentem, pedicello columnae affixam. *Corolla* nulla. *Filamenta* in flosculo terminali tria inferne coalita, in flosculis lateralibus vix manifesta. *Stamina*. *Calyx* tripartitus, minimus, germini aditus, persistens. *Corolla* petala tria, rigida, acuta, persistentia. *Germen* inferum. *Styli* tres simplices. *Bacca* carnosa, subrotunda, apice tribus denticulis umbilicata. *Semina* ossicula tria, hinc convexa, inde angulata, oblonga.

1541. JUNIPERUS communis foliis ternis patentibus mucronatis bacca longioribus. *Linn. Sp. pl.* 2. p. 1470. — *Pall. fl. pal.* n. 934.

Juniperus foliis convexo-concavis aristatis, baccis alaribus sessilibus. *Hall. stirp. helv.* n. 1661.

Juniperus vulgaris fruticosa. *Bauh. pin. pl. Mapp. fl. Als.* p. 158.

Juniperus baccifera. *Tabernaemont. p.* 1357. *Icon.*

Icon. *Pfleger arb. et frut. Badens.* — *Oeder. fl. Dan.* t. 1119. — *Blackw. t.* 187. — *Regnault. bot.*

Germ. Gemeiner Wachholder.

Gall. Gènevrier commun.

Hab. In sylvis aridis siccioribus, collibus apertis utrinque passim.

Flor. Martio, Aprili. *Baccae* insequente anno Octobri maturae. **5**

Frutex 2-3-5 pedalis, quandoque arbuscula, sempervirens. *Baccae* ad foliorum axillas sessiles, caeruleo-nigricantes.

Pharm.

Pharm. JUNIPERI Lignum; Summitates; Baccæ.

Lignum durum, gratam fragrantem flammam alit. Baccæ dulcescentes, aromaticæ, sudoriferæ, incipientes, fermentatæ spiritum ardentem largiuntur.

Oleum empyrevmaticum fragrans, ex tota planta obtentum, optimo cum successu externe contra ovium scabiem (die Schafräude) adhibetur, idem certe præstat quod in Gallia meridionali Oleum Cati, huil de cade, quod ex Junipero Oxycedro ibi in abundantia rescente; aut Miera hispanorum: certissimum et unicum apud greges ovilas hispanicas remedium, quod est oleum empyrevmaticum ex Junipero Oxycedro, hoenicea, Lycia et thurifera ibi promiscue in abundantia crescentibus.

542. JUNIPERUS Sabina foliis oppositis erectis decurrentibus: oppositionibus pyxidatis. *Linn. Sp. pl. 2. p. 1472.*

Juniperus foliis cauli adpressis, lanceolatis, alterne conjugatis. *Hall. stirp. helv. n. 1662.*

Sabina folio Cupressi. *Bauh. pin. 487.*

Sabina mas et foemina. Tabernaem. p. 1355. Icon.

Icon. Bulliard. herb. t. 139. — Pfleger arb. et frut.

Ger. m. Sadebaum. Sevenbaum. Scvibaum. Stinkender - Wachholder.

Gall. Gênevrier Sabine.

Hab. In Hispania, Gallia meridionali, Sibiria, Oriente. Hospitatur vulgo in hortis rusticorum.

Flor. Martio, Aprili. *Baccæ* Octobri maturæ. **5**

Frutex 2 3-5 pedalis, quandoque in hortis arbuscula. *Rami* diffusi, sparsi, valde ramosi, densi,

recti *Foliis* oppositis, linearibus, acutis, convexo-concavis, basi appressis, brevibus sempervirentibus, glabris.

Pharm. SABINAE Herba.

Herba fragrans, gravis, fortis, amaricans, quinem maxime movens, calida, violenta. In carie cum pulvis Sabinæ a quibusdam laudatur. Decoctum aut infusum Herbae pediculos pellit.

Nota. In hortis rusticorum ob perniciosum suum abusum diminuenda aut plane removenda.

Obs. In hortis Carlsruhanis sub dio hospitantur *JUNIPERUS Oxycedrus virginiana* et *Sabina* fol. variegatis.

CCCCLII. TAXUS.

Linn. Gen. pl. ed. Schreb. n. 1553.

Mas. Calyx nullus: nisi *Gemma* similis perianthio tetraphyllo. *Corolla* nulla. *Filamenta* numerosa, coalita. *Antherae* depressae, octidadae, basi undique dehiscences (et dejecto polline) planae, peltatae, margine octofido notabiles. *Femina* Calyx nullus. *Corolla* nulla. *Bacca* e receptaculo elongato in praeputium globosum, succulentum, apice dehiscens coloratum. Semen unicum, ovato-oblongum, apice extra baccam prominens.

1543. *TAXUS baccata* foliis approximatis.

Linn. Sp. pl. 2. p. 1472. Hall. stirp. helv. n. 1663.

Taxus. *Bauh. hist. 1. p. 241. — Mapp. fl. Alf. p. 300.*

Icon. Pfleger arb. et frut. Badens. — Schkuhr. bot. Handb. t. 339.

Ferm. Eibenbaum. Taxbaum.

Fall. If commun.

I a b. In Marggraviatu superiore in montosis sylvaticis prope *Gernsbach*, ubi hodie rarius. In *Alsatia in dem Niedeker Thal*, juxta rivum *t. Mappo*. In Principatu Fürstenbergenli prope *Heiligenberg*.

Flor. Febuario, Martio, Aprili. *Bacca* Octobri, Novembri maturae. *h*

Frutex saepe arbor mediocris, sempervirens. *Cortex* fuscus. *Folia* approximata, pectinata, linearia, coriacea. *Flores* axillares. *Bacca* viscosa, glutinosa, pulchellae.

Lignum exquisitum rubello - marmoreum, durum, ornatoribus et fabris scriniariis expetitur.

Bacca fatuae dulcescentes vix suspectae aut venetae, avibus acceptae. *Folia* pecoribus, et equis noxia.

b f. Ex hac Classe et ordine hospitantur adhuc in hortis Carlsruhanis *EPHEDRA monostachya* et *distachya* Linn.

CLASSIS XXIII.
POLYGAMIA

MONOECIA.

CCCCLV. HOLCUS. ♀ *Glum.* 1-flora. *Glum.*
2-valvis. *Stam.* 3. *Styl.* 2.
Sem. 1.

♂ *Glum.* 1-flora. *Glum.* 1-
valv. *Stam.* 3.

CCCCLIV. ANDROPOCON. ♀ *Glum.* 1-flora.
Glum. basi aristata. *Stam.* 3.
Styl. 2. *Sem.* 1.

♂ *Glum.* 1-flor. *Glum.* bā
aristata. *Stam.* 3.

CCCCLVI. VALANTIA. ♀ *Cal.* nullus. *Cor.* 4-
partita. *Stam.* 4. *Styl.* 2-fid.
Sem. 1.

COCCLIII. VERATRUM. ♀ *Cal.* nullus. *Cor.* 6-
petala. *Stam.* 6. *Pist.* 3. *Capf.* 3.
♂ *Cal.* nullus. *Cor.* 6-petala.
Stam. 6.

CCCCLIX. ACER. ♀ *Cal.* 5-fidus. *Cor.* 5-petala.
Stam. 8. *Styl.* 2. *Capf.* 2-coo-
ca, alata.
♂ *Cal.* 5-fidus. *Cor.* 5-pe-
tala. *Stam.* 8.

MONOECIA.

773

CCLVII. **PARIETARIA.** ♂ *Cal.* 4-fidus.

Cor. o. *Stam.* 4. *Styl.* 1. *Sem.* 1.

♀ *Cal.* 4-fidus. *Cor.* o. *Styl.* 1.

Sem. 1.

CCLVIII. **ATRIPLEX.** ♂ *Cal.* 5-phyllus.

Cor. o. *Stam.* 5. *Styl.* 2-fid.

Sem. 1.

♀ *Cal.* 2-phyll. *Cor.* o. *Styl.*

1-fid. *Sem.* 1.

DIOECIA.

CCLX. **FRAXINUS.** ♂ *Cal.* o. f. 4-part. *Cor.* o.

f. 4-petala. *Stam.* 2. *Pistill.* 1.

Sem. 1.

♀ *Cal.* o. f. 4-part. *Cor.* o. f.

4-petala. *Pistill.* 1. *Sem.* 1.



CLASSIS XXIII.

POLYGAMI

MONOE.CIA.

CCCCLIII. V E R A T R U M.

Linn. Gen. pl. ed. Schreb. n. 1564.

Flos hermaphroditus. Calyx nullus. Corolla
Petala sex, oblonga, lanceolata, marginem
ora, ferrata, persistentia. Filamenta sex, subula
corolla breviora. Antherae quadrangulae. Ge
mina tria. Styli vix manifesti. Stigmata tria
Capsulae tres, oblongae, erectae, comp
uniloculares, univalves, introsum dehiscen
Semina plura, oblonga, compressa, mem
nacea. Masculus flos in eadem planta, inflo
minos. Calyx ut in masculo. Corolla ut in

POLYGAMIA. MONOECIA. 775

on. *Jacqu. fl. Austr. t. 336. — Schkuhr. bot. Handb. t. 341.*

erm. *Schwarzer Germer. Weisse Nießswurz mit schwarzen Blumen.*

all. *Vératre noir.*

ab. In Alfatiae summis jugis. In monte *Spiesberg* inter *St. Dié* et *St. Marie aux mines*. nec non in monte retro *Blinschweiler* loco dicto *der Kreutzpfad. t. Mappo* et *Tabernaemontano.*

or. Julio, Augusto. *Semina* Septembri, Octobri matura. ☞

Radix oblonga, praemorsa, fibris copiosis side flavescens, crassiusculis, praedita. *Caulis* 3-4 pedalis, erectus, simplex, crassiusculus, obsolete angulatus, villosus, foliosus. *Folia* radicalia, caulinaque inferiora, petiolata, ovato vaginante, ovalia, acuta, integra, firma, nervosa, plicata, glabra, reliqua alterna sessilia, vaginantia, suprema minora, breviora angustiora. *Racemus* terminalis, reliquis racemis superioribus axillis foliorum ortis longior, densior, erectus; cylindrici, multiflori, densi, triplices, infimi ramo uno alterove aucti. *Colla* purpureo-atra, patentissima. *Antherae* prode rubrae.

Planta florens omnino speciosa.

Radix et tota planta venenata, drastica, vomitoria, arrima, in lacte cocta muscas perimit, mures necat.

Ad rariores omnino Florae cives pertinet.

CCCCLIV. ANDROPOGON.

Linn. Gen. pl. ed. Schreb. n. 1566.

Gramen. Hermaphroditi flores sessiles. *Caulis* *Gluma* uniflora, bivalvis, mutica: *Valvula*

exteriore basi cincta lanugine bipartita. *Corolla*.
Gluma bivalvis: *valvula* exteriore minore, aristata: *Arista* terminali, longa, geniculata, inferne tortili. *Nectarium* diphyllum, diaphanum. *Stamina* tria. *Styli* duo capillares. *Stigmata* palmosa. *Semen* solitarium oblongum, gluma *Corollae* et calycis tectum. *Masculi* flores pedunculati, solitarii s. gemini, singulo hermaphrodito adstantes. *Calyx* ut in hermaphrodito. *Corolla* absque arista. *Nectarium* ut in hermaphrodito. *Stamina* ut in hermaphrodito.

1545. **ANDROPOGON** *Ischaemum* spicis digitatis plurimis, flosculis hermaphroditis sessilibus aristatis, masculis pedunculatis, muticis, pedicellis lanatis.

Andropogon Ischaemum. Linn. Sp. pl. 1: 1483. — Poll. fl. pal. n. 935.

Andropogon spicis villosis septenis, loculis bifloris, folliculis alternis sessilibus. Hall. stirp. helv. n. 1414.

Gramen Dactylon spicis villosis. Bauh. pin. 8. — Scheuchz. Agrostr. p. 94. — Murr. fl. Alf. p. 132.

Icon. Schreb. Graef. 2. t. 33. — Jacqu. Austr. t. 384.

Germ. Bartgras.

Gall. Barbon vélue.

Ha b. In collibus graminosis aridis, ad vias, agrorum versuras gramineas passim frequens. Circa Carlsruhe am Landgraben versus Mühlburg, inter Mühlburg et Grünwinkel.

Fr. Julio, Augusto. *Semina* Septembri, Octobri matura. γ

Radix longe lateque repens. *Culmi* sesquiduales et altiores, basi incurvi, abhinc erecti, quandoque inferne ramosi. *Genicula* fusco-purpurea. *Folia* carinata, rariter pilosula. *Ligula* brevissima, villosa-pilosa. *Spicae* 4-5-6-13, erectae, lineares, 2-3 uncias longae, pilosae.

Non confundendum cum *Panico Dactylon*. **nn.**

Sabulum in arenosis figit. **Gramen** adhuc tenellum vix acceptum.

CCCGLV. H O L C U S.

Linn. *Gen. pl. ed. Schreb. n.* 1565.

Gramen. *Hermaphroditi* flores subseffiles. *Calyx* *Gluma* 2-3 flora, bivalvis mutica, compressa, carinata, subciliata. *Corolla* *Gluma* bivalvis, calyce minor, mutica. *Nectaria* foliola duo, lanceolata. *Filamenta* tria. *Antherae* bidae. *Semen* solitarium, ovatum, corollae adnatum, calyce tectum. *Masculi* flores pedunculati, solitarii, seu gemini singulo hermaphrodito adstantes, minores. *Calyx* *Gluma* bivalvis, minor, valvula exterior infra apicem aristata: *Arista* geniculo subreflexa. *Nectarium* ut in hermaphroditis. *Stamina* ut in hermaphroditis. *Germen* parvum abortiens.

1546. **HOLCUS** *lanatus* glumis bifloris villosis: hermaphrodito mutico, masculo

arista recurva. *Linn. Sp. pl.* 2. p. 1485. —
Poll. fl. pal. n. 936.

Avena diantha floribus ovatis, perfecto ma-
 tico, imperfecto aristato. *Hall. stirp. helv.*
 n. 1484.

Aira foliis pubescentibus, panicula contracta,
 flosculo hermaphrodito mutico, masculo
 arista uncinata, calyce brevior. *Linn. fl.*
Suec. p. 67.

Gramen pratense paniculatum molle. *Bauh.*
pin. 2. *prodr.* 3. — *Scheuchz. Agric.*
p. 434. — *Mapp. fl. Alf.* p. 134.

Icon. Schreber. Graefss. t. 20. *f.* 1. — *Curtis.*
fl. Lond. t. 228. — *Oeder. fl. Dan. t.* 170.
Leers. fl. Herb. t. 7. *f.* 6.

Germ. Honiggras. Wolliches Darrgras.
Gall. Houque laineuse.

Ha b. Utrunque in pratis humidis, pomariis et
 ad fossas frequens, nec sicciore respuit.

Flor. Majo, Junio. **Semina** Junio, Julio ma-
 tura. ¶

Radix fibrosa, non repens. **Culmi** plures,
 sesquipedales, cubitales, tripedales et altiores,
 erecti, geniculati, teretes, pubescentes, molles.
Folia mollia, villosa, plana, latiuscula. **Vaginae**
 laxae, striatae, villosae. **Ligula** truncata, crenu-
 lata, externe pilis ciliata. **Panicula** primo spica-
 formis, mollis, laxa, rubella, subnutans, demum
 erecta, diffusa, patens, varie rubelle colorata
 dein pallescens. **Spiculae** biflorae villosulae, rufae
 apicem coloratae.

Pulchrum et egregium in pratis praesertim humidis
 gramen, bonum pecoribus pabulum, etiam ovibus con-
 veniens.

Nota. In pratis Rheno vicinis aliisque humidis omnino studio propagandum. Pro pratis artificialibus in locis humidis peraptum.

47. *HOLCUS mollis* glumis bifloris nudiusculis: flosculo hermaphrodito mutico, masculo arista geniculata. *Linn. Sp. pl.* 2. p. 1485. — *Poll. fl. pal.* n. 937.

Avena *Hall. stirp. helv.* n. 1485.

Gramen paniculatum molle, radice graminis canini repente. *Morif. hist.* 3. p. 202.

Gramen caninum paniculatum molle. *Scheuz. Agrostr.* p. 235.

Gramen caninum longius radicatum. *Bauh. pin.* 1.

n. Schreber. Graef. t. 20. f. 2. — *Oeder. fl. Dan.* t. 1059. — *Curtis. fl. Lond.* t. 223. — *Leer. fl. Herb.* t. 7. f. 7.

rm. Weiches Honiggras.

ll. Houque mollet.

b. Utrinque in cultis, sabulosis agris, fylvis, ad viarum marginalia passim frequens nec prata respuit, in his tamen rarior.

r. Aestate. *Semina* Julio, Augusto matura. ¶

Affinis primo intuitu praecedenti, a qua distinguitur, *Radice* quammaxime repentici canini instar, *Culmis* paucioribus, articulis procumbentibus, *Foliis* angustioribus, vix pubescentibus (*Spiculae* biflorae, quae triflorae), *Arista* calyce longiore.

Bonum pecoribus et ovibus pabulum praebet, in locis ficcis locis magis propagandum, ubi arenam

In horto culta, constantissima permanfit, minime pro variatione praecedentis habenda.

1548. *HOLCUS odoratus* glumis trifloris
muticis glabris: hermaphrodito diandro.
Linn. Sp. pl. 2. p. 1485.

Avena locustis trianthis paniculatis hians
flosculo tertio imperfecto. *Hall. J. helv. n. 1496.*

Avena foliis planis, calyce trifloro, flosculo
intermedio mutico diandro. *Hof. J. helv. 2. p. 152.*

Aira foliis planis, floribus trifloris an-
tis, flosculo intermedio mutico diandro.
Linn. fl. Suec. 70.

Gramen paniculatum odoratum. *Bauh. pin.
7. — Scheuchz. Agrost. 236.*

Icon. Schkuhr. bot. Handb. t. 344. an?

Germ. Geruchvolles Honiggras.

Gall. Houque odorante.

Hab. In Alsatia prope Mulhusiam in pratis
tro dem Thurn, paulo rarius, sed circa Pab-
versheim prata replet t. *Hofero.*

Nota. Hoferi gramen hucusque in terris Baden-
sibus non vidi.

CCCCLVI. VALANTIA

Linn. Gen. pl. ed. Schreb. n. 1575.

Hermaphroditus flos solitarius. *Calyx* vi-
nullus. *Corolla* monopetala, plana, quadri-
tita: laciniis ovatis, acutis. *Stamina* quatuor
Germen magnum, inferum. *Stylus* semibifidus
Stigmata capitata. *Capsula* coriacea, compressa,
reflexa. *Semen* unicum, globosum. *Masculus* flos
hermaphrodito utrinque solitarius adfistens. *Ca-*

vix ullus. *Corolla* monopetala, plana, tripar-
aut quadripartita; *laciniis* ovatis, acutis. *Sta-*
a 4. f. 3. *Germen* parvum, inferum. *Stylus*
stigmata obsoleta, vix notabilia. *Capsulae* ru-
mentum. *Semen* nullum.

49. *VALANTIA Aparine* floribus maf-
culis trifidis pedicellatis, hermaphroditicis
pedunculo infidentibus. *Linn. Sp. pl.*
2. p. 1491. — *Poll. fl. pal. n.* 938.

Galium tricornae. Lamarck et Decand. fl.
franc. 4. p. 262. — *Smith. fl. brit. 1. p.* 176.

Valantia triflora. Lamarck. et franc. 3. p.
384.

Galium foliis fenatis, ferratis, petiolis triflo-
ris, incurvis. *Hall. stirp. helv. n.* 725.

Aparine semine laevi. *Vaill. paris. 18. t. 4.*
f. 3. a. a. — *Mapp. fl. Als. p.* 25. (*sub*
sequente specie.)

r m. Klebkrautartige Vaillantia.

all. Vaillantie gaillet.

z b. In cultis, agris et vineis passim.

o r. Augusto, Septembri. *Semina* Septembri,
Octobri matura. ☉

Medium quasi tenet locum inter *Galium Apa-*
rine et *spurium*. Differt a priori, *Statura* demissi-
s, *fructibus* non pilosis, sed tuberculis minu-
exasperatis, ab altero pedunculis trifloris folia
n excedentibus, *Fructibus* majoribus.

50. *VALANTIA saccharata* caulibus
decumbenti- adscendentibus, foliis sub-
senis linearibus, margine ferrulato aspe-
ris, pedunculis apice curvatis, seminibus
saccharatis.



Gal. valianae anis juccre.

Hab. In cultis inter fegetes p.
Flor. Junio, Julio. *Semina*

A praecedente specie affi
Linnaeum, Hallerum, Mappu
sim botanicis militat, facile fr
latis saccharatis distinguitur.

Obs. Planta repetitis vicibus in hor
non succubuit, hinc omnino 1
declaranda.

1551. *VALANTIA cruciat*
culis quadrifidis, pedu
Linn. Sp. pl. 2. p. 1491.
n. 939.

Galium foliis quaternis, hirsu
floris. *Hall. stirp. helv. 7*

Cruciata hirsuta. Bauh. pin.
fl. Alf. p. 84.

Icon. Blockus herb. 176

Caules plures sesquipedales et altiores, erecti, basi quandoque decumbentes tetragoni, pilosi. *Folia* quaterna, sessilia, ovato-lanceolata, serrata, carina marginibusque pilosa. *Pedunculi* axillares, utrinque tres, diphylli, trifidi, pilosi; *in* cellulo medio simplici, lateralibus ramofo-quadrifidis. *Flores* masculi et hermaphroditi quadrifidi, quandoque quinquefidi, lutei.

var. m. CRUCIATAE f. ASPERULAE AUREAE
Herba.

Olim in vulnerariis et nervinis habebatur, hodie medicina exoleta.

Flores melliferi.

Radix rubro colore tingens.

CCCCLVII. PARIETARIA.

Linn. Gen. pl. ed. Schreb. n. 1576.

Hermaphroditi flores duo continentur *In* involucro plano, hexaphyllo: foliolis duobus oppositis et exterioribus majoribus. *Calyx* monophyllus, quadrifidus, planus, obtusus. *Corolla* nulla. *Stamina* quatuor. *Antherae* didymae. *Stig-* penicilliforme, capitatum. *Calyx* seminiferus integatus, campanulatus, ore clauso. *Semen* unum, ovatum. *Femineus* flos unicus inter hermaphroditos ambos, intra involucrum. *Calyx* in hermaphrodito. *Corolla* nulla. *Pistillum* ut in hermaphrodito. *Calyx* involvens fructum. *Semen* ut in hermaphrodito.

52. *PARIETARIA officinalis* foliis lanceolato-ovatis, alternis. *Linn. Sp. pl. 2. p. 1492. — Poll. fl. pal. n. 940.*

Gall. Pariétaire. Casse pierre.

Hab. In ruderatis, ad sepes, nemum et vinearum muros passim.

Flo r. Junio, Julio. *Semina* Aug.

Caules plures, sesquipedales, ascendentes, erectiusculi, firriati, inferne ramosi. *Folia* alterna, per lanceolata, utrinque attenuata, incisas longa, patentia, subtus nervo goso-pubescentia. *Flores* verticillati conglomerati: *Pedunculi* gemini. *Stamina* calyce longiora, inflexa: resiliencia pollinem antherarum inclinata: sic etiam acus ictu, quod adfirationem de irritabilitate antherarum fit.

Pharm. PARIETARIAE Herba.

Planta subsalsa, aperientes, diuretica.

Ob asperitatem hinc inde ad vitra adhaerenda recipitur.

POLYGAMIA MONOECIA. 785

Parietaria foliis ovato lanceolatis subhirsutis.
Hall. stirp. helv. n. 1613.

Parietaria minor ocymi folio. Bauh. pin. 121?

Parietaria ocymi folio Tragi. Boccon. ficul.
47. t. 24. A.

non. Schkuhr. bot. Handb. t. 344.

erm. Jüdisches Glaskraut.

all. Pariétaire de Judée.

ab. In Sponhemia ad rudera arcium antiqua
prope Winterburg et Martinstein, prope
Winningen ad Mosellam. In regionibus rhe-
nanis ad rudera arcium prope Bingen, Caub,
Bacharach, Weissenfels, Coblenz et An-
dernach frequens.

or. Junio, Julio. Semina Augusto matura.

Affinis praecedenti, a qua differt: *Caulibus*
noribus, digitalibus, palmaribus, vix pedali-
s. Foliis ovatis, brevioribus et latioribus, sub-
hirsutis, saepe rubellis. Florum verticillis mino-
ris, in summo caule magis contiguis, pur-
pureis.

os. Planta in horto culta mutationi non succubuit.

CCCGLVIII A T R I P L E X.

Linn. Gen. pl. ed. Schreb. n. 1577.

Hermaphroditus flos. Calyx pentaphyllus,
concavus, persistens: laciniis margine membra-
ceis. Corolla nulla. Stamina quinque. Anthe-
ae didymae. Stylus bipartitus. Stigmata reflexa.
Calyx fructiferus clausus, pentagonus. Semen
oblongum, orbiculatum, depressum. Feminis flos
eadem planta. Calyx diphyllus: foliolis planis,
Flora Bad. Tom. III.

D d d

erectis, ovatis, acutis, magnis, compressis.
 rolla nulla. *Stylus* bipartitus. *Stigmata* re-
Calycis fructiferi valvae magnae, cordatae
 men includentes unicum, orbiculatum, ap-
 pressum.

1554. *ATRIPLEX hortensis* caule
 herbaceo, foliis triangularibus. *Linn. Sp.*
pl. 2. p. 1493.

Atriplex hortensis alba f. pallide virens. *Bauh.*
pin. 119.

β. *Atriplex hortensis rubra.* *Bauh. pin. 119.*
Icon. Regnault. bot. — Blackw. t. 99. et
Germ. Gartenmelde. β. Rothe Gartenmelde.
Gall. Arroche cultivée.

Hab. In Tartaria. Colitur in hortis et
 ex quibus in ruderata pagorum et
 saepe aberrat.

Flor. Julio, Augusto. *Semina* Septembri
 tura. ☉

Planta pro usu culinario nota.

1555. *ATRIPLEX rosea* caule herbaceo
 foliis serratis. *Linn. Sp. pl. 2. p. 1493.*
Poll. fl. pal. n. 941.

Atriplex sylvestris, fructu, roseo, compresso.
Bauh. pin. 119. — Prodr. p. 58.

Germ. Rosenmelde.

Gall. Arroche rose.

Hab. Circa Klein Riedesheim, Hessen et
 vias, t. Pollichio.

Flor. Septembri. *Semina* Octobri maturantur.

Caulis sesquipedalis, cubitalis, erectus, sub-
gulosus, albus, glaber, ramosissimus, ramis
ernis, denuo ramosis, diffusis. *Folia* alterna,
sessilia, rhombeo-cordata, glabra, ultra pol-
em longa et lata, margine sinuato-denticu-
a, incana, pulvere albido tenuissimo quasi con-
rsa. *Flores* axillares, sessiles, conglomerati,
afi. *Fructus* valvulae ex subrotundo ovatae,
stiusculae, compressae, lateribus denticula-
, margine pulcherrime crenulatae, incanae.
Pollichius.

Nota. Hanc plantam hucusque in terris Badensibus
et affinibus regionibus non vidi.

56. ATRIPLEX hastata caule herba-
ceo, valvulis foemineis magnis deltoidi-
bus sinuatis. *Linn. Sp. pl. 2. p. 639. —*
Pollich. fl. pal. n. 942.

Atriplex foliis triangularibus, basi productis,
valvulis triangularibus, asperis. *Hall. stirg.*
helv. n. 1617.

Atriplex sylvestris annua, folio deltoide tri-
angulari sinuato et mucronato hastae cu-
spidis simili. *Morif. hist. 2. p. 607. Sect.*
5. t. 32. f. 14.

Atriplex sylvestris folio hastato s. deltoide.
Mapp. fl. Alf. p. 35.

rm. Spondonblacttrige Melde.

ll. Arroge en fer de Lance.

b. Utrunque passim in ruderatis, ad vias, mu-
ros, sepes et fossas.

r. Julio, Augusto. *Semina* Septembri ma-
tura. ☉

Caulis cubitalis, tripedalis et altior, erectus
pe flexuosus, sulcatus, ramosus; *Rami* oppo-

lycibus feminum disco
pl. 2. p. 1494. — Poll.

Atriplex foliis imis hamati
latis, masculis seminalib
stirp. helv. n. 1616.

Atriplex angusto oblongo
219. — *Mapp. fl. Alf. 1*

Atriplex sylvestris Polygoni
liis. *Lob. Icon. 257.*

Atriplex sylvestris. *Taberna*
Germ. Ausgebreitete Melde.
Gall. Arroche étalée.

Hab. Utrunque in ruderatis, se
tes pagorum et urbium, n
passim frequens.

Flo r. Julio, Augusto. *Semina*
tobri matura. ☉

Caulis 2-3-5 pedalis, erect
prostratus, angulosus, striatus,
Rami oppositi alternique, dive

ota. Huc sine dubio etiam pertinet *Atriplex angustissima* et longissimo folio. Mapp. fl. Alf. p. 35. ad Argentinam inter portam Cronenburg et Steinstrasser Thor, auf dem Glacis citata. Synonymon alias ad *Atriplicem littoralem* Linn. citatum, quae Argentorati non crescit.

ATRIPLEX Halimus Linn. caule fruticoso, foliis deltoidibus integris in hortis Carlsruhanis sub dio hospitatur.

CCCCLIX. A C E R.

Linn. Gen. pl. ed. Schreb. n. 1590.

Hermaphroditi Calyx monophyllus, quin-
fidus, acutus, coloratus, persistens. *Corolla*
la quinque, ovata, obtusa, calyce vix ma-
, patentia. *Stamina* octo. *Stylus* filiformis,
mata duo, reflexa. *Capsulae* 2-3, basi coa-
, subrotundae, compressae, ala membra-
a maxima singulae terminatae. *Semina* fo-
ia, subrotunda. *Masculi flores Calyx* — Co-
 — *Stamina* ut in hermaphroditis. *Germen*
um. *Stylus* nullus. *Stigma* bifidum.

58 *ACER Pseudo-Platanus* foliis
quinelobis inaequaliter ferratis, flori-
bus racemosis. *Linn. Sp. pl. 2. p. 1495.* —
Poll. fl. pal. n. 944.

Acer foliis quinelobis acute ferratis, race-
mis pendulis. *Hall. stirp. helv. n. 1029.*

Acer montanum candidum. Bauh. pin. p. 430.
n. Pfleger arb. Badensf.
m. Weisser Bergahorn. Grofser Ahorn.

In loco humido densas et durabiles sepes praebet.

Lignum durum venosum exquisitum.

Folia pecoribus pabulum.

Flores apibus melliferi.

Nota. Acer major folio rotundiore minus laciniato, an Opulus Italicorum? Mapp. fl. AlC. p. 5. forsitan mera variatio, crescens secundum Mappum in monte St. Ottiliae et versus Hochfeld. In locis citatis saltem Acerem Opalum et opulifolium frustra repetitis vicibus cum Cl. Hermann o, praeceptori et amico nunquam non tuis venerando quaesivi.

1561. ACER monspessulanum foliis trilobis integerrimis glabris, annuis. *Linn. Sp. pl.* 2. p. 1497. — *Scopol. Carn. ed. 2.* n. 1246.

Acer foliis trilobis cordatis integerrimis, lobis aequalibus divaricatis. *Du Roi. Harbke.* p. 26.

Acer trifolium. *Bauh. pin.* 431.

Icon. Pfleger arb. et frut. Badenf. — *Dummel. arb.* 1. p. 28. t. 10. f. 8.

Germ. Mas holder von Montpellier.

Gall. Erable de Montpellier.

Hab. In Sponhemia prope *Kirn, Naumburg, Fischbach, Oberstein, Idar, Birkenfeld*, et alibi frequens.

Flor. Aprili. Semina Julio matura. **B**

Frutex 4-5-6 pedalis, rarius arbuscula. *Folia* opposita, petiolata, perfecte triloba: lobis aequalibus, integris, obtusis, laete viridibus, glabris, firmis. *Semina* alata, purpureo-rubella, glabra, fere parallela, subdivergentia.

POLYGAMIA MONOECIA. 793

Nota. Omni ex parte cum speciminibus in gallia meridionali et Hispania lectis, observatisque convenit; Semina matura copiosa in Sponhemia fert, que in horto disseminata, egregie vegetabant. Individua ex seminibus in horto obtenta, mutationi non succubuerunt.

Obs. Sine dubio eandem Pollichius in monte Donnersberg observavit, et sub variatione A. campestris, foliis trilobis, integerrimis obtusis, allegavit.

Hucusque a Germaniae Floristis omissa.

In hortis Carlsruhanis sub dio hospitantur: Acer ataricum, rubrum, dasycarpon. Ehrh. — accharinum — Platanoides laciniatum, triatum Du Roi. monspeffulanum, cretium, hybridum et Negundo.

POLYGAMIA DIOECIA.

CCCCLX. FRAXINUS.

Linn. Gen. pl. ed. Schreb. n. 1597.

Hermaphroditus. Calyx nullus, aut monophyllus quatripartitus. Corolla nulla, aut Petala quatuor, linearia, longa, acuta. Stamina duo. Germen ovatum compressum. Stylus cylindraceus. Stigma bifidum. Capsula ovato-oblonga, superne foliacea, bilocularis, altero lobulo abortiente. Semen lanceolatum, compresso-membranaceum, uniloculare. Femina, Calyx, Corolla, Pistillum, Capsula, Semen ut in hermaphrodito.

1562. FRAXINUS excelsior foliolis serratis, floribus apetalis. *Linn. Sp. pl. 2. p. 1509. — Poll. fl. pal. n. 947.*

Fraxinus apetala foliis pinnatis ferratis. *Hall. stirp. helv. n. 528.*

Fraxinus excelsior. *Bauh. pin.* 416. — *Ternaemont. p.* 1428. *Icon.* — *Mapp. fl. Als. p.* 111.

Icon. Pfleger arb. Badens. — *Oeder. fl. Dn.* t. 969. — *Regnault. bot.*

Germ. Esche, Asche, Edel-Esche.

Gall. Frêne élevé.

Hab. Passim in sylvis humidiusculis montosis et demissioribus.

Flor., Aprili. *Semina* Julio matura. 6

Arbor interdum sat procera, longaeva, saepe autem humilis. *Folia* impari-pinnata: foliola ovato-lanceolata, ferrata, glabra, saturate viridia. *Flores* conglomerati, ante folia e gemmis lateralibus erumpentes, aetate laxe racemosi. *Calva* nullus. *Corolla* nulla. *Femina* saepe interspersa habet flores hermaphroditos, et hermaphroditos semineos.

Pharm. FRAXINI Cortex; Folia; Semina.

Adstringentes vires habet. Cortex cortici peruviani a quibusdam substituitur.

Folia siccata ovibus per hyemem gratum pabulum suppeditant.

Fronde arboris saepe Meloë vesicatorius Linn. devassat.

Lignum perdurum, album, luteo-variegatum. fabricis lignariis, tornatoribus, mechanicis, architectonibus et nautis acceptissimum.

Nota. Arbor hujus quammaxime procera, omnium facile speciosissima, trunco longo recto, crasso praedita in Horto Durlacensi summum decus.

1563. *FRAXINUS Ornus* foliolis ferratis, floribus corollatis. *Linn. Sp. pl.* 2. p. 1510.

Fraxinus florifera. *Scopol. fl. Carn.* 2. n. 1250.

Fraxinus paniculata. *Mill. dict. n. 4.*

Fraxinus humilior f. altera Theophrasti, minore et tenuiore folio. *Bauh. pin. 416.* —

Mapp. fl. Alf. p. 112.

Fraxinus tenuiore et minore folio. *Bauh. hist. 1. p. 177.*

Germ. *Blumentragende Esche.*

Gall. *Frêne à fleurs.*

Hab. In Alsatiæ Vogefis ad rivum juxta monasterium *Peris* versus *dem schwarzen See*; prope *Rappolstein* in cacumine juxta trium arcium rudera antiqua. *t. Mappo.*

Flor. Martio, Aprili. *Semina* Junio matura. **h**

Nota. Arbusculam alias in calidioribus regionibus obviam, frustra in Vogefis et terris Badenſibus hucusque quaesivi. Facile autem haec species Floribus paniculatis longe corollatis distinguitur.

Obf. In horto Carlsruhano sub dio hospitat, sed hyemes fortiores aegre fert.

Porro in hortis Carlsruhanis sub dio hospitantur: **FRAXINUS** aurea, horizontalis, pendula, monophylla, americana, nigra, alba, sambucifolia, juglandifolia, novae angliae, caroliniana, pubescens, Lamarck. rotundifolia f. *Ornus mannifera*, mycrophylla lanceifolia et rotundifolia.

Porro ex hac Classe hospitantur in hortis Carlsruhanis et passim Schwezingensibus aliisque sub dio. **CELTIS** australis, occidentalis, orientalis, et cordata. **CLEDITSCHIA** triacanthos, inermis, sinensis, monosperma et horrida. **DIOSPYROS** virginiana. **NISSA** aquatica et sylvatica. **FIGUS** Carica.

Finis tomi tertii.







